



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:


- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



Chem 668.77.3




This volume was a part of a bequest to the  
Medical School by  
**EDWARD M. BARRINGER**  
of Schenectady, N. Y. For a consideration,  
paid to the Medical School out of the  
**MINOT FUND,**  
the books (not medical) so bequeathed have  
been transferred to the Library of the  
University, April 8, 1881.

SCIENCE CENTER LIBRARY






25









AN ELEMENTARY TREATISE  
ON  
PRACTICAL CHEMISTRY

AND  
QUALITATIVE INORGANIC ANALYSIS,  
SPECIALLY ADAPTED FOR USE IN THE LABORATORIES OF SCHOOLS  
AND COLLEGES, AND BY BEGINNERS.

BY  
FRANK CLOWES, D.Sc. Lond.  
FELLOW OF THE CHEMICAL SOCIETIES OF LONDON AND BERLIN; SENIOR SCIENCE  
MASTER AT THE HIGH SCHOOL, NEWCASTLE-UNDER-LYME; LATE  
SCIENCE MASTER AT QUEENWOOD COLLEGE.

*WITH ILLUSTRATIONS.*  
FROM THE SECOND ENGLISH EDITION.



PHILADELPHIA:  
HENRY C. LEA.  
1877.



~~V. 7H~~

Chem 668.77.3

1877  
(Bancroft) Reginald.

## PREFACE TO THE SECOND EDITION.

---

THIS little treatise was commenced to supply a course of Practical Chemistry to my own classes. I was encouraged to proceed with it, by finding that the want of a sufficiently elementary and explanatory Laboratory Text-book was very widely felt.

It has been my aim throughout to give all necessary directions so fully and simply as to reduce to a minimum the amount of assistance required from a teacher. The language employed has been rendered simple and intelligible by avoiding the unnecessary use of scientific terms, and by explaining or paraphrasing in ordinary words any such terms when introduced for the first time. The directions how to work, and the description of the preparation and use of apparatus, have been given more fully than is usual, since my own experience, confirmed by that of other teachers, convinces me that one of the most serious hindrances to the utility of many of the smaller Text-books on Practical Chemistry is the too great conciseness of the language employed, which frequently renders it unintelligible to the student unless supplemented by copious verbal explanation from the teacher.

Whilst making the very desirable amplifications above referred to, the book has been kept within small dimen-

sions, partly by the omission of all such higher instruction as is not required by a student of elementary chemistry, and partly by the insertion of the supplementary or merely explanatory portions in smaller type. I have also thought it best to avoid entering into any lengthy theoretical explanations. The modern teaching of chemistry is in practice very appropriately divided into two departments, namely, theoretical instruction imparted by lectures or by the study of text-books of theoretical chemistry, and practical instruction imparted in a chemical laboratory by working according to the directions of a practical text-book. Since by this system the student has time and opportunity afforded him for the study of the theoretical and descriptive portions of the science, it is as unnecessary as it is undesirable that his Practical Text-book should tempt him to bestow valuable time in the laboratory upon the study of matters of theoretical, not practical, importance.

The analytical reactions and methods have been carefully worked through from the text by myself and by the members of my classes: the accuracy and intelligibility of their descriptions have thus, I hope, been secured; only those reactions and methods which are commonly employed for analytical purposes have been entered. I have naturally, in selecting analytical methods for an elementary treatise, felt it desirable that those chosen should be as simple and easy of execution as possible; in some cases, however, methods which are most eligible on these grounds have proved on trial to be so inferior in accuracy and delicacy, that they have been abandoned in favor of others which are recommended by their reliability rather than by their simplicity. In such cases, however, I have also described

the more simple methods, since they may be employed in analyses, in which minute quantities of a substance have not to be tested for. The reactions given in an elementary text-book have necessarily been limited as to number, and I have felt it in general advisable to introduce such reactions as are useful in general analysis, rather than those which claim to be merely interesting and instructive.

The book has been divided into seven sections, the contents of which are fully stated on pages ix-xv. The first six contain a good practical course for senior students; this may, however, be modified to suit junior students, or those working with a special object, as is shown in the Introduction.

The seventh section contains full lists of all apparatus, reagents, and chemicals required in working through the different sections; there is also added a list of general apparatus, with a description when necessary of its construction and use. In this section there will also be found full and systematic descriptions of the most simple methods for preparing the different solutions required in analysis, with a statement of the strength most appropriate for each. Experience has proved that these are matters which merit more attention than is usually bestowed upon them. The methods of preparing pure chemicals are omitted, since they may now be readily and cheaply purchased; before using purchased chemicals, their purity should, however, always be ascertained by the tests given in this section.

Symbolic notation has been employed, instead of the full chemical names, throughout the sections on analytical chemistry; in its most concise form this chemical shorthand conduces so much to brevity in writing down

results that no other plea is required for its use. The simple plan of labelling each bottle in the laboratory with the chemical formula as well as the name of its contents, will prevent any difficulty arising from this general employment of chemical formulæ.

The "Tables of Differences," which contain for each Analytical Group a summary of the differences of behavior of its members with reagents, are special—being an extension of the system employed in Galloway's *Manual of Qualitative Analysis*.

It is almost superfluous to mention that free use has been made of the standard works of Fresenius and Rose: much valuable information has been introduced from these sources. I have also frequently adopted the very convenient tabular form of entering analytical methods which is employed in Valentin's *Text-book of Practical Chemistry*, and with the author's permission have transcribed, with a few trivial alterations, the excellent Phosphate Table devised by him. My acknowledgments are also due to Dr. W. A. Tilden, of Clifton College, and to the Rev. T. N. Hutchinson, of Rugby, and to many other teachers who have suggested valuable improvements.

The book is especially intended to furnish a course of instruction in practical chemistry in the laboratories of our public and other schools. It will thus supply a demand which is rapidly increasing, as the value of a sound elementary instruction in practical science is becoming more widely appreciated, both as means of mental training and as a preparation for the chemical and medical professions, as well as for many branches of manufacturing industry and enterprise. The fifth section has been inserted for the use of those who are spe-

cially preparing for practical examinations in which proficiency in the analysis of simple salts only is required of the candidate. This is the standard fixed for the Preliminary Scientific (M. B.) Examination of the London University, and for the more elementary examinations in the Oxford University, such as those for school certificates and open scholarships.


The sixth section, however, contains additional details suited for the higher analytical work of advanced students, and will be found sufficient to qualify a student for the higher examinations in analytical chemistry, such as the B. Sc. Honors Examination in the London University.

The introduction as appendices into this edition of the reactions and methods of detection of the rarer elements, and of the use of the spectroscope with a spectrum chart, will, it is believed, render the book more useful to advanced students.

F. C.

NEWCASTLE-UNDER-LYME,  
November, 1876.





# CONTENTS.

SECTION I.		Paragraph.	Page.
EXPERIMENTS ILLUSTRATING THE METHODS OF PREPARATION AND PROPERTIES OF GASES, ETC.			
I.	Oxygen, . . . . .	...	15
II.	Hydrogen, . . . . .	...	22
III.	Carbon dioxide, . . . . .	...	25
III a.	Nitric oxide, . . . . .	...	30
IV.	Ammonia, . . . . .	...	81
V.	Carbon monoxide, . . . . .	...	33
V a.	Chlorine, . . . . .	...	36
V b.	Hydrochloric acid gas, . . . . .	...	36
VI a.	Distillation of water, . . . . .	...	36
VI b.	Preparation of nitric acid, . . . . .	...	38
SECTION II.			
PREPARATION AND USE OF APPARATUS RE- QUIRED FOR CHEMICAL ANALYSIS, . . . . .			
	Bunsen burner, . . . . .	1-19	40
	Spirit-lamp, . . . . .	1, 2	40
	Blowpipe, . . . . .	3	42
	Glass tubes, cutting, . . . . .	4	42
	" bending, . . . . .	5	43
	" drawing out, . . . . .	6	44
	Boring corks, . . . . .	7	44
	Making sulphuretted hydrogen tube, . . . . .	8	45
	Making ignition tubes, . . . . .	9	45
	" stirring-rods, . . . . .	10	46
	Mounting platinum wires, . . . . .	11	46
	Making wash-bottle, . . . . .	12	47
	Cleaning glass and porcelain apparatus, . . . . .	13	47
	Cleaning platinum-foil and wire, . . . . .	14	49
	Putting away apparatus, . . . . .	15	50
	Heating glass and porcelain: . . . . .	16	50
	Porcelain dishes, . . . . .	17	51
	Glass vessels, . . . . .	18	51
		19	51

	Paragraph.	Page.
<b>SECTION III.</b>		
<b>ANALYTICAL OPERATIONS, DESCRIPTIONS ILLUSTRATED BY EXPERIMENTS, . . . . .</b>	<b>20-35b</b>	<b>53</b>
Solution, . . . . .	20	53
Evaporation, . . . . .	21	55
Precipitation, . . . . .	22	56
Filtration and decantation, . . . . .	23, 23a	58
Decantation, . . . . .	23b	60
Washing precipitates on the filter, . . . . .	24, 24a	61
"        "        by decantation, . . . . .	24b	61
Drying precipitates, . . . . .	25	62
Removing precipitates from the filter, . . . . .	26	62
Ignition, . . . . .	27	64
Sublimation, . . . . .	28	65
Fusion, . . . . .	29	66
<i>The Use of the Blowpipe:</i> . . . . .	30	66
Borax-beads, . . . . .	31	67
Flame colorations, . . . . .	32	69
Ignition on charcoal in the blowpipe flame, . . . . .	33	70
Levigation of residue, . . . . .	33a	72
<i>The Use of Test-papers:</i> . . . . .	34	72
Making neutral, acid, and alkaline, . . . . .	35	74
Turmeric-paper, . . . . .	35a, 35b	75
Fractional solution, crystallization and distillation, . . . . .		76
<b>SECTION IV.</b>		
<b>ANALYTICAL REACTIONS, . . . . .</b>	<b>36-329</b>	<b>78</b>
Introductory remarks on analytical classification and on the analytical course which follows. . . . .		
Analytical classification, . . . . .	36	81
Analytical groups, . . . . .	37	82
Method of trying the reactions (rules), . . . . .	38	83
<i>Entry in the Note-book:</i> . . . . .	39	84
Chemical notation, . . . . .	40	84
Chemical equation, . . . . .	41	85
Rules for drawing out an equation, . . . . .	42	85
Contractions to be employed, . . . . .	43	87
Examples of showing method of entry in the note-book, . . . . .	44	87
Rules to be observed whilst working, . . . . .	45	87
<b>Reactions for Metals:</b>		
<i>Group V (Potassium Group):</i> . . . . .	46-47	89
Potassium, . . . . .	47-50	90
Ammonium, . . . . .	51-55	92

	Paragraph.	Page.
Sodium, . . . . .	56, 57	93
Magnesium, . . . . .	58-63	94
Table of Differences and its use, . . . . .	64, 65	96
Principles of separation and detection, . . . . .	66	97
Table for separation and detection, . . . . .	67	99
<i>Group IV (Barium Group):</i> . . . . .	68-89	100
Barium, . . . . .	69-74	100
Strontium, . . . . .	75-79	101
Calcium, . . . . .	80-86	101
Table of Differences, . . . . .	87	102
Principles of separation and detection, . . . . .	88	102
Table for separation and detection, . . . . .	89, 438	103
<i>Group III A (Iron Group):</i> . . . . .	90-109	103
Aluminium, . . . . .	91-94	104
Iron, . . . . .	95-101	104
Chromium, . . . . .	102-106	106
Table of Differences, . . . . .	107	108
Principles of separation and detection, . . . . .	108	109
Table for separation and detection, . . . . .	109, 436	109
<i>Group III B (Zinc Group):</i> . . . . .	110-141	109
Zinc, . . . . .	111-114	110
Manganese, . . . . .	115-120	111
Nickel, . . . . .	121-128	112
Cobalt, . . . . .	130-137	114
Table of Differences, . . . . .	138	116
Detection of members separately, . . . . .	139	117
Principles of separation and detection, . . . . .	140	117
Table for separation and detection, . . . . .	141, 437	118
<i>Group II A (Copper Group):</i> . . . . .	142-176	118
Mercuricum, . . . . .	143-148	118
Lead, . . . . .	149-153	120
Bismuth, . . . . .	154-159	122
Copper, . . . . .	160-168	123
Cadmium, . . . . .	169-172	125
Table of Differences, . . . . .	173	126
Detection of members separately, . . . . .	174	126
Principles of separation and detection, . . . . .	175	126
Table for separation and detection, . . . . .	176-435a	126
<i>Group II B (Arsenic Group):</i> . . . . .	177-210	127
Arsenic, . . . . .	179-190	128
Antimony, . . . . .	191-200	135
Tin, . . . . .	201-205	137
Table of Differences, . . . . .	206	139
Principles of separation and detection, . . . . .	207	139
Table for separation and detection, . . . . .	208-435b	140
A second method of separation and detection, . . . . .	209	140
A third, . . . . .	210	141

	Paragraph.	Page.
<i>Group I (Silver Group):</i> . . . . .	211-222	143
Lead, . . . . .	[149-153]	120
Silver, . . . . .	212-216	143
Mercurous, . . . . .	217-219	143
Table of Differences, . . . . .	220	144
Principles of separation and detection, . . . . .	221	144
Table for separation and detection, . . . . .	222, 433	145
Reactions for gold, . . . . .	223, 224 <sub>a</sub>	145
Reactions for platinum, . . . . .	225, 226	145
<b>Reactions for Acid Radicles:</b> . . . . .	227-329	147
<i>Group I. Sulphate Group:</i> . . . . .	227, 228	147
<i>Group III. Carbonate Group:</i> . . . . .	229-246	148
Carbonates, . . . . .	229	148
Sulphides, . . . . .	230-235	149
Sulphites, . . . . .	236-238	151
Theiosulphates, . . . . .	239, 240 <sub>a</sub>	151
Hypochlorites, . . . . .	241, 242	152
Nitrites, . . . . .	243-245	153
Detection of the members when mixed, . . . . .	246	153
<i>Group III (Nitrate Group):</i> . . . . .	247-256	154
Nitrates, . . . . .	247-251	154
Chlorates, . . . . .	252-255	156
Detection when mixed, . . . . .	256	157
<i>Group IV (Chloride Group):</i> . . . . .	257-274	157
Chlorides, . . . . .	257-260	157
Bromides, . . . . .	261-264	159
Iodides, . . . . .	265-270	161
Detection of separate members, . . . . .	271	163
Separation and detection, . . . . .	272-274	164
<i>Group V (Phosphate Group):</i> . . . . .	275-282	166
Phosphates, . . . . .	275-278 <sub>a</sub>	166
Arsenates, . . . . .	279	167
Detection, . . . . .	280-281 <sub>a</sub>	168
Separation and detection, . . . . .	282	168
<i>Ungrouped Acid Radicles (Inorganic):</i> . . . . .	283-302	169
Borates, . . . . .	283-285	169
Chromates, . . . . .	286-289	170
Silicates, . . . . .	290-293	172
Fluorides, . . . . .	294-298	173
Fluosilicates, . . . . .	299-302	175
<i>Organic Acid Radicles:</i> . . . . .	303-328	177
Cyanides, . . . . .	303-306	177
Ferrocyanides, . . . . .	307-310	179
Ferricyanides, . . . . .	311-314	179
Sulphocyanides, . . . . .	315, 316	180
Oxalates, . . . . .	317-319	180
Tartrates, . . . . .	320-325	181
Acetates, . . . . .	326-328	183

	Paragraph.	Page.
<b>SECTION V.</b>		
<b>ANALYSIS OF SIMPLE SUBSTANCES FOR ONE METAL AND ONE ACID RADICLE,</b>	<b>330-369b</b>	<b>185</b>
Introductory remarks,	330	185
Preliminary examination of a solid for the metal,	331, 333	188
Solution of a solid substance,	332	189
Preliminary examination of a liquid for the metal,	334	194
Actual examination for the metal in a neutral or acid liquid,	335-345	196
Examination of a solid or liquid for the acid radicle:	346-358b	207
Introductory remarks,	346-347	207
Preliminary examination,	348-351	209
Actual examination,	352-358b	211
Examination of an alkaline liquid,	359-360a	214
<i>Examination of a substance with metallic appearance,</i>	366	217
<i>Examination of an insoluble substance,</i>	367-369b	218
<i>Examples of entry of analyses,</i>		222
<b>SECTION VI.</b>		
<b>FULL ANALYTICAL COURSE AND TABLES,</b>	<b>370-493</b>	<b>232</b>
<i>Introductory:</i>		
Separation of metals into groups by group reagents,	370	232
Addition of reagents in excess,	371	233
Evaporation before precipitating Group III,	372	233
General remarks,	373, 374	234
Preliminary examinations for their use,	375-378	234
<i>General analytical course:</i>		
Preliminary examination of a liquid,	380, 381	237
Solution of a non-metallic solid,	382-385	238
Preliminary examination of solids for the metals present,	387-409	240
Preliminary examination of solids for acid radicles,	410-418	246
<i>Actual examination for metals:</i>	419-439	249
General table and notes thereon,	419-432	349
Table I (Silver Group) and Table G,	433, 434	254
Table II (Copper and Arsenic Groups),	435	255
Table III A (Iron Group),	436	257
Table III B (Zinc Group),	437	259
Table IV (Barium Group),	438	261
Table V (Potassium Group),	439	263



	Paragraph.	Page.
<i>Actual examination for acid radicles, . . .</i>	440-454	265
Table showing solubility of substances in water and acids, . . .	455	271
Explanation and use of the table, . . .	456-458	272
Remarks on the precipitation of Group III, Rules for the precipitation and examination of Group III, . . .	459-462	278
Table III c <sub>1</sub> (Iron and Zinc Groups mixed), . . .	463	275
Table III c <sub>2</sub> (Iron and Zinc Groups mixed), . . .	464	276
Table III d (Phosphates in Group III), Examination of substances with metallic appearance, . . .	465	278
Separation and detection of gold and platinum, . . .	466	279
Examination of substances insoluble in water and acids, . . .	467-471	281
Analysis of silicates, . . .	472-474	285
Analysis of substances containing cyanogen, Example showing how to enter analytical results, . . .	475-479	286
	480, 481	290
	482-487	291
	488-498	296
<b>SECTION VII.</b>		
<b>APPARATUS AND CHEMICALS REQUIRED FOR THE PRECEDING COURSE, . . .</b>	494-539	303
<i>List of apparatus for each student, . . .</i>	494	303
<i>List and description of general apparatus: . . .</i>	495-506	306
Indigo-prism, cobalt-glass, cork-borers, . . .	496-498	306
Sulphuretted hydrogen apparatus, . . .	499, 500a	306
Passing sulphuretted hydrogen, . . .	500a	309
Agate mortar, leaden or platinum crucible, . . .	501, 502	310
Steam-oven, water-bath, tubulated flasks, . . .	503-505	310
Distillation of water, . . .	506	318
Recovery of the metals from silver and platinum residues, . . .	507, 508	316
<i>Directions for preparation of reagents and solutions: . . .</i>	509-517	318
Introductory remarks, . . .	509, 510	318
Dilution of liquids, . . .	511	318
Solution of solids, . . .	512-515	319
Preparation of solutions for the reactions, <i>Lists of the reagents, solutions, and chemicals, stating their condition or strength and special processes of preparation: . . .</i>	516, 517	323
Introductory and explanatory remarks, . . .	518, 533	323
Labelling and varnishing, . . .	518-521	323
List of reagents for each bench, . . .	522, 523	324
List of reagents for general use, . . .	524	326
	525, 526	328

	Paragraph.	Page.
Preparation of saturated solutions of solids,	527	332
"                    "                    " of gases,	528	332
List of solutions required for Section IV, .	529, 530	336
List of chemicals required for Section I, .	531	339
List of chemicals required for Section III,	532	340
List of sundry other requisites, . . . . .	533	341
<i>Examples of substances to be given for analysis,</i>	534-539	341
<hr/>		
APPENDIX I.—Reaction and detection of the		
Rarer Elements, . . . . .	540-554	346
Group I. ( <i>Silver Group</i> ): . . . . .	540, 541	346
Thallium, . . . . .	540	346
Tungstates, . . . . .	541	347
Group II A. ( <i>Copper Group</i> ): . . . . .	542	348
Palladium, . . . . .	542	348
Group II B. ( <i>Arsenic Group</i> ): . . . . .	543-545	348
Molybdates, . . . . .	543	348
Selenium, . . . . .	544	349
Tellurium, . . . . .	545	349
Group III. ( <i>Iron and Zinc Groups</i> ): . . . . .	546-550	350
Uranium, . . . . .	546	350
Indium, . . . . .	547	351
Beryllium, . . . . .	548	351
Titanium, . . . . .	549	351
Vanadium, . . . . .	550	352
Group V. ( <i>Potassium Group</i> ): . . . . .	551, 552	352
Lithium, . . . . .	551	352
Rubidium and Cæsium, . . . . .	552	353
General Table for detection of the rarer elements, . . . . .	553	354
Table for examination of Group III, for the rarer elements, . . . . .	554	356
APPENDIX II.—The spectroscope, its use in analysis, . . . . .	555	357
Spectrum-chart, . . . . .	557	359
APPENDIX III.—List of chemical elements with symbols and atomic weights, . . . . .	558	361
Thermometric scales, . . . . .	559	361
Weights and measures, French and English, . . . . .	560, 561	362

- W. L. & J. A. Roberts (1995) *Journal of Animal Ecology*, **64**, 101–112.
- W. L. & J. A. Roberts (1996) *Journal of Animal Ecology*, **65**, 101–112.
- W. L. & J. A. Roberts (1997) *Journal of Animal Ecology*, **66**, 101–112.
- W. L. & J. A. Roberts (1998) *Journal of Animal Ecology*, **67**, 101–112.
- W. L. & J. A. Roberts (1999) *Journal of Animal Ecology*, **68**, 101–112.
- W. L. & J. A. Roberts (2000) *Journal of Animal Ecology*, **69**, 101–112.
- W. L. & J. A. Roberts (2001) *Journal of Animal Ecology*, **70**, 101–112.
- W. L. & J. A. Roberts (2002) *Journal of Animal Ecology*, **71**, 101–112.
- W. L. & J. A. Roberts (2003) *Journal of Animal Ecology*, **72**, 101–112.
- W. L. & J. A. Roberts (2004) *Journal of Animal Ecology*, **73**, 101–112.
- W. L. & J. A. Roberts (2005) *Journal of Animal Ecology*, **74**, 101–112.
- W. L. & J. A. Roberts (2006) *Journal of Animal Ecology*, **75**, 101–112.
- W. L. & J. A. Roberts (2007) *Journal of Animal Ecology*, **76**, 101–112.
- W. L. & J. A. Roberts (2008) *Journal of Animal Ecology*, **77**, 101–112.
- W. L. & J. A. Roberts (2009) *Journal of Animal Ecology*, **78**, 101–112.
- W. L. & J. A. Roberts (2010) *Journal of Animal Ecology*, **79**, 101–112.
- W. L. & J. A. Roberts (2011) *Journal of Animal Ecology*, **80**, 101–112.
- W. L. & J. A. Roberts (2012) *Journal of Animal Ecology*, **81**, 101–112.
- W. L. & J. A. Roberts (2013) *Journal of Animal Ecology*, **82**, 101–112.
- W. L. & J. A. Roberts (2014) *Journal of Animal Ecology*, **83**, 101–112.
- W. L. & J. A. Roberts (2015) *Journal of Animal Ecology*, **84**, 101–112.
- W. L. & J. A. Roberts (2016) *Journal of Animal Ecology*, **85**, 101–112.
- W. L. & J. A. Roberts (2017) *Journal of Animal Ecology*, **86**, 101–112.
- W. L. & J. A. Roberts (2018) *Journal of Animal Ecology*, **87**, 101–112.
- W. L. & J. A. Roberts (2019) *Journal of Animal Ecology*, **88**, 101–112.
- W. L. & J. A. Roberts (2020) *Journal of Animal Ecology*, **89**, 101–112.
- W. L. & J. A. Roberts (2021) *Journal of Animal Ecology*, **90**, 101–112.
- W. L. & J. A. Roberts (2022) *Journal of Animal Ecology*, **91**, 101–112.
- W. L. & J. A. Roberts (2023) *Journal of Animal Ecology*, **92**, 101–112.
- W. L. & J. A. Roberts (2024) *Journal of Animal Ecology*, **93**, 101–112.
- W. L. & J. A. Roberts (2025) *Journal of Animal Ecology*, **94**, 101–112.

# PRACTICAL AND ANALYTICAL CHEMISTRY.

---

## INTRODUCTION.

THE course of practical chemistry which should be pursued by a student depends partly upon his object in studying the science, and partly upon the time and means which he has to devote to the study. The first four sections and the sixth furnish a good general course for a senior student who wishes to obtain a training in practical and analytical chemistry; but for younger students the most suitable analytical course consists in trying through the reactions in Section IV, and, as the reactions for each group are completed, testing several substances containing only one member of the group for the metal present by the table of differences. The analysis of simple substances by Section V may then be worked through, and, after becoming familiar with this, separations of the mixed members in each group are done, leading thus to Section VI. The modification which should be made in the student's course to suit individual cases will, perhaps, be most readily understood by stating the objects of each section in the book:

SECTION I is adapted to impart experience in chemical manipulation, and to afford practical illustrations of the elementary portions of theoretical or descriptive chemistry; it may be omitted when instruction in analysis only is required.

SECTION II is indispensable and must be carefully perused.

SECTION III is also indispensable to the student of analytical chemistry, but if the time at his disposal is *very limited*, he may omit the performance of the experiments described in this section, and merely read through the text.

SECTION IV must be carefully worked through. A student whose time is limited may, however, simply try the reactions and omit the performance of analyses at the end of each group.

SECTION V is intended specially for students whose object is to learn only the analyses of simple salts, such as is required in many modern examinations on practical chemistry.

This section may also be used as an easy beginning in analysis, and as an introduction to more complicated analyses to be made by Section VI, or it may be passed over by the student who is intending to learn general analysis.

SECTION VI is not required by a student who is learning only the analysis of simple salts, but should be carefully worked through by the student of general analysis, who may, after becoming thoroughly conversant with its contents, pass on to quantitative analysis or to any special branch of practical chemistry he may require.

## SECTION I.

## PREPARATION OF GASES, ETC.

IN this section full directions are given for the preparation of five gases (viz., oxygen, hydrogen, carbon dioxide, ammonia, and carbon monoxide), and for certain interesting and instructive experiments which may be made with them. The processes of preparation and manipulation required for these gases are more or less typical of those employed for all other gases, and the student will therefore, from the experience obtained by performing the experiments with the above-named gases, easily prepare and experiment upon the three other gases (nitric oxide, chlorine, and hydrochloric acid) by following the directions inserted in small print, and any other gas by the account given of it in a treatise on chemistry. Those gases which are in small type may be omitted in the practical course, if desired. Two examples of the process of distillation are also appended.

The reference numbers inclosed in brackets refer to the paragraphs which commence with Section II (p. 40); the numbers will be found in thick type in the text, and at the head of each page the numbers of the paragraphs are also placed in square brackets.

A full list of the apparatus required for this section is given in par. 494, and its use is explained in pars. 1-10 and 14-19; a list of chemicals and certain other requisites will be found in par. 531.

The student must carefully read through *the whole description* of each experiment before beginning to perform it, and after its successful performance should enter a brief description of it in his note-book.


**I. OXYGEN GAS.**—When iron is for some time exposed to moist air its surface becomes covered with rust;



many other metals undergo a similar change in moist air, but the alteration produced in their appearance is not usually so noticeable as in the case of iron. The liquid metal mercury does not rust as iron does in moist air, but it becomes slowly covered with red mercury-rust when strongly heated for some time in a flask open to the air; this mercury-rust has received the name of mercuric oxide. The fact that metals become heavier by rusting proves that something is added during the process.


EXP. 1.—Place in a clean and perfectly dry test-tube sufficient mercuric oxide to cover the bottom; heat the

FIG. 1.




powder (1, 19) as shown in Fig. 1, loosely stopping the end of the tube with the thumb. As soon as small drops of mercury form on the sides of the tube, remove the thumb and quickly place inside the top of the tube the burning end of a slip of wood (*e. g.*, the uncoated end of a wooden lucifer match); the flame will be seen to burn more brightly. If after again heating the powder for some time in the way just described, the slip

be introduced into the mouth of the tube immediately after blowing out the flame and whilst there is a spark at its end, the glowing end will be caused to burst into flame. This behavior with a burning or glowing slip of wood is one of the most remarkable properties of oxygen gas, and we frequently make use of this property as a "test" for its presence. The chemical change which has occurred is thus represented by an equation:



Since by heat, then, we can separate from mercuric oxide mercury and oxygen, we learn that the process of rusting consists in the metal taking oxygen gas from the air, and the increase of weight above referred to is thus accounted for; if all metal rusts could be decomposed by

heat we might obtain oxygen from them just as from mercuric oxide. This method of making oxygen is interesting since it was the first means known of preparing the gas; it is never used nowadays to prepare large quantities of oxygen, since other substances are known which contain a large proportion of oxygen, and give it off, when they are heated, more easily than mercuric oxide does—substances which are also preferable on account of their greater cheapness. Potassium chlorate is most frequently employed:



EXP. 2.—Place in a clean dry test-tube a little potassium chlorate, and heat it as in Exp. 1. The white salt, after decrepitating (crackling), fuses (or melts) and when further heated, appears to boil; the small bubbles which are given off consist of oxygen gas, as may readily be proved by holding in the mouth of the test-tube a burning or glowing splinter of wood as described in Exp. 1.

Potassium chlorate gives off oxygen gas much more readily than does mercuric oxide; but if it is mixed with small quantities of certain other substances, which themselves appear to undergo no change, its oxygen is driven off by heat with extreme facility; of these substances manganic oxide (black oxide of manganese) is the one usually chosen.


EXP. 3.—Powder some potassium chlorate (about as much as would fill a watch-glass) finely in a mortar, mix with it, by rubbing them together in the mortar, about one-fifth as much powdered manganic oxide, and heat a small quantity of the mixture in a test-tube; the oxygen will begin to come off as soon as the mixture is heated, and a comparatively gentle heat will cause the gas to be rapidly evolved.

In the preceding experiments the oxygen was detected in the tube in which it was prepared, and was allowed to pass away freely into the air. When the gas has to be collected in a vessel unmixed with air, it is made to pass through a bent glass tube (the delivery-tube), which is fitted by means of a cork air-tight into

the mouth of the test-tube; the end of this tube dips into some water, and the bubbles of gas are allowed to rise into a vessel full of water and inverted over the end of the delivery-tube. This process of "collecting" oxygen is fully described in the following experiment; in the performance of which two students may advantageously work together, one attending to the regulation of heat to the mixture, the other to filling the gas-jars.

EXP. 4.—Select a sound cork, of such a size that, after having been softened by being squeezed or by being rolled with gentle pressure on the floor under the foot, it fits tightly into the mouth of the test-tube to be employed. Then bend (6) a piece of hard glass tubing about fourteen inches in length, into the form shown in the figure; so adapting the bends by trial that when the apparatus is fitted together the bottom of the test-tube

FIG. 2.



may be at a convenient height in the flame, the end of the delivery-tube at the same time dipping about an inch under water. Make a hole through the centre of the cork (8), of such a size that the glass tube fits tightly into it. Then test whether the apparatus is air-tight by fitting the glass tube into the cork, and the cork into the test-tube, and blowing down the open end of the delivery-tube; no air must be heard to escape, or must be seen to bubble out on moistening the cork: if air does escape a fresh cork must be taken. Now pour into the perfectly dry test-tube the oxygen mixture (see Exp. 3)

off a piece of paper folded into a trough, or scoop up the mixture from the mortar with the mouth of the test-tube, until the tube is about one-third full, and fit in the cork and delivery-tube.

Before heating the tube fill the jar in which the oxygen is to be collected with water, close it with a stopper or ground-glass plate (or with the hand), invert its mouth into water three or four inches in depth, contained in an earthenware pan or bowl, and carefully remove the stopper or plate. If this operation has been performed with proper precaution the jar will be entirely filled with water, and no air-bubble will remain. Next proceed to heat the upper part of the oxygen mixture, holding the tube in the right hand; keep the lamp slowly moving with the left, in order to prevent any part of the glass from being suddenly and strongly heated, which would be liable to crack it. Oxygen gas will soon be evolved, but will not at once appear at the end of the delivery-tube, since it has first to drive out the air which filled the apparatus; as soon as a slip of wood glowing at its end is kindled, when held at the mouth of the delivery-tube, the oxygen has driven out the air, and is beginning to escape; the end of the delivery-tube is then at once dipped under water beneath the mouth of the jar, and the stream of bubbles rising into it will rapidly displace the water. As soon as the jar is full of gas, close its mouth under water with the stopper or glass plate, and remove it for experiment. The jar may also be removed by slipping under its mouth a small dish or saucer, the water taken out in the saucer then closes the mouth of the jar air-tight.

*Precautions.*—The water must be removed from the pan, when it rises inconveniently high, by means of a small porcelain dish; if at any time the gas should be given off too rapidly, the flame should be removed until the current slackens; the lower portions of the mixture should be heated only after the upper parts refuse to yield any more gas; when the process is to be stopped, the end of the delivery-tube must be removed from the water before the gas has ceased to bubble out, and the


test-tube must not be allowed to touch cold or wet objects, which would cause the hot glass to crack.

Several bottles filled with oxygen will be required for the following experiments, or the same bottle may if necessary be refilled with the gas according to the above directions, after the completion of each experiment.

*Note.*—The use of ground-glass plates, which must close the mouth of the jar perfectly air-tight, is much easier than that of stoppers. An earthenware "beehive shelf" which may be used in a common earthen pan, or a "pneumatic trough," is also convenient, since it supports the jar during the process of collection.

Oxygen gas is remarkable for the energy with which it combines with or burns many substances; three examples are given of this property in Expts. 5, 6, and 7.


EXP. 5.—Select a splinter of wood-charcoal or a small piece about the size of a nut; the experiment is more brilliant if the surface of the charcoal formed originally part of the bark of the tree. Bind this upon a "deflagrating spoon" with a little fine iron or copper wire; then adjust the wire handle of the spoon in the brass cap, so that when held beside the bottle of oxygen with the cap on a level with the mouth of the jar, the little metal cup is about an inch from the bottom of the bottle. Now heat the charcoal in the Bunsen flame, or better the blowpipe flame (4), until a part of its surface glows when held in the air, and quickly place it into the bottle of oxygen, with the brass plate covering the mouth of the bottle. (See Fig. 7, p. 30.) The charcoal will burn much more brilliantly than in air, throwing off sparks if its surface was "barky:"



When it ceases to burn pour into the bottle a little clear lime-water from a small beaker or test-tube, quickly close the bottle and shake the liquid round inside it; the clear liquid becomes milky, indicating the presence of carbon dioxide gas, as will be hereafter explained.


EXP. 6.—Remove the charcoal from the deflagrating spoon and replace it by a piece of sulphur as large as a pea; heat the spoon in the flame until the sulphur melts

and begins to burn with a pale-blue almost invisible flame. Then place the spoon into a fresh jar of oxygen, the sulphur will at once burn with a much larger flame, which emits a beautiful violet light:




Sulphur dioxide (sulphurous anhydride) gas remains in the bottle; its presence is proved by its suffocating smell, also by pouring a little water into the bottle and shaking it round. Sulphurous acid is thus formed, and is recognized by dropping into the water a piece of blue litmus-paper, which is immediately reddened, and by pouring in a few drops of red potassium dichromate solution, the color of which changes to green.

EXP. 7.—Cleanse the deflagrating spoon from any remaining sulphur, and place into it a small piece of phosphorus no larger than a pea. The phosphorus may be cut with a knife, but it must be touched only with *wet* fingers, and should be handled as little as possible, since it is liable to catch fire by the heat of the hand; it is always kept under water, being dried only immediately before being used by pressing it between filter-paper or blotting-paper, or with a dry cloth. Set fire to the phosphorus by holding the spoon in the flame, and notice how it burns in the air; then place the spoon in a jar of oxygen—the phosphorus will burn most brilliantly, producing a white substance called phosphorus pentoxide (or phosphoric anhydride):



When the phosphorus has ceased to burn pour in a little water and shake it round in the jar, the white substance dissolves, yielding phosphoric acid:



and the water may now be proved to be acid by dropping into it a piece of blue litmus-paper, which will be immediately reddened.


*Test for Oxygen.*—A convenient test for oxygen is to introduce into the gas a slip of wood with a spark at the

end, which is caused to burst into flame. Only one other gas possesses this property, and it is readily distinguished from oxygen by other means. This test only detects oxygen when it is in a pretty pure condition.

II. HYDROGEN GAS.—The liquid substance water consists of oxygen gas, combined with another gas called hydrogen; several processes are known for preparing hydrogen from water. This gas is, however, most readily obtained from another liquid containing it, called hydrochloric acid, by the action upon it of the metal zinc.

EXP. 8.—Fit a two-necked Woulffe's bottle\* (A, Fig. 3), with air-tight perforated corks bearing a thistle fun-


FIG. 3.



nel (B), whose end reaches nearly to the bottom of the bottle, and a tube bent as shown at C and terminating just below the cork: join to this, by means of a short piece of tightly fitting india-rubber tube, a bent delivery-tube D. Pour into the bottle sufficient granulated zinc to cover the bottom, replace the corks, and after ascertaining that the apparatus is air-tight by closing the end of the delivery-tube and blowing down the thistle-fun-

\* A wide-necked bottle may be used instead, being fitted as shown in Fig. 5 (p. 26).

nel, pour in through the funnel sufficient water to cover the zinc and the end of the funnel-tube; then add strong hydrochloric acid gradually until, after mixing the acid and water by shaking the bottle, the hydrogen is seen to rise from the zinc in numerous small bubbles:



Then dip the end of the delivery-tube under water contained in the pan or trough, and allow the gas to bubble out through the water for at least five minutes. This delay is necessary in order to give the hydrogen time to entirely remove the air which filled the bottle, and which when mixed with hydrogen produces a dangerously explosive mixture. Before collecting larger quantities of the gas for experiments, ascertain that the hydrogen is no longer mixed with air by inverting a test-tube filled with water over the end of the delivery-tube; as soon as the tube is full of gas close its mouth with the thumb, and hold it to a flame; if the gas burns with a slight explosion, the tube is again similarly filled with the gas and tried in the same way. As soon as the gas burns quietly with a pale flame, a jar (or a small thick glass cylinder or tube) may be filled with it in the same way as was directed for oxygen.


If during the preparation of gas for the following experiments the gas comes off too slowly, it is only necessary to pour in a little more strong acid through the funnel and mix it with the liquid in the bottle by gently shaking the latter.

EXP. 9.—Hold the jar filled with hydrogen with its mouth open and directed upwards for a short time, the gas will entirely escape; the absence of the gas may be shown by holding a lighted taper in the jar, when no flame will be seen at the mouth. If the jar be refilled with hydrogen and held for a short time mouth downwards, the hydrogen will remain in it, and its presence may be shown by the gas burning with a pale flame when a lighted taper is introduced: these results prove that hydrogen is much lighter than air, since its tendency to rise prevents it from passing out downwards



through the open mouth, whilst it readily escapes upwards from the erect cylinder. Since hydrogen is so much lighter than air, it is possible to collect the gas without using water by a process called "displacement." The delivery-tube of the hydrogen apparatus is passed up to the top of the inverted jar containing air, the hydrogen rises to the upper part of the jar and gradually pushes out the heavier air downwards.

EXP. 10.—Fill a jar with hydrogen by "displacement." For this purpose fit upon the india-rubber joint of the hydrogen apparatus a delivery-tube bent as shown in Fig. 4, and when a brisk effervescence of gas has been caused by pouring in some strong hydrochloric acid, pass




this tube up to the top of an inverted jar, and allow the jar to remain in this position for several minutes; it is best to loosely close the mouth of the jar during this process by letting it rest upon a perforated disk of card-board (or the round brass cap of a deflagrating spoon), supported upon a retort-stand ring, or upon an iron tripod-stand, since the entrance of air by "diffusion" is thus almost entirely prevented.

Remove this jar, keeping it mouth downwards, and push up inside it a burning wax taper five or six inches in length; the hydrogen will be lighted, and will burn with a pale flame at the mouth of the jar, but the flame of the taper will be seen to be extinguished by the gas: the taper may, however, be rekindled by holding it in the hydrogen flame burning at the mouth of the jar.

EXP. 11.—Cover the bottle and funnel-tube with a cloth, to prevent accident in case of an explosion, and light the hydrogen at the end of the delivery-tube used in the last experiment. Hold over the flame a perfectly clean, dry, and cool tumbler or beaker; the inside will

## EXP. 12.] PREPARATION OF CARBON DIOXIDE. 25

become dimmed with moisture, showing that hydrogen gas burning in the air produces water :



*Note.*—Since all gases which have been in contact with water contain more or less vapor of water, or are “moist,” it is usual to employ for this Exp. a stream of hydrogen gas which has been freed from moisture or “dried.” A gas is dried (or desiccated) by passing it through some substance which readily absorbs moisture. Strong sulphuric acid (oil of vitriol), calcium chloride, and quicklime are the desiccating agents most commonly employed. The gas may be made to bubble through strong sulphuric acid contained in a bottle, B (Fig. 10, p. 35), connected, as shown in the figure, with the generating apparatus. It may be also dried by passing it through a tube containing fragments of calcium chloride, or of quicklime (Fig. 10*a*), or pieces of pumice-stone moistened with strong sulphuric acid: the moistened pumice may be contained in a U-tube (Fig. 10*b*), or in the bottle B (Fig. 10, p. 35).

**EXP. 12.**—Measure the height of a short thick glass cylinder, and divide it into three equal parts by small pieces of gum-paper stuck upon the outside. Fill the cylinder with water, and, after inverting it in water, fill one-third with oxygen (Exp. 4), and the remainder with hydrogen (Exp. 8); let the jar stand with its mouth under water for five or six minutes to allow the gases to mix, then apply a lighted taper to the mouth of the jar directed downwards, taking care not to place the fingers beneath it: the gases combine to form water with a loud explosion.

*Test.*—Hydrogen gas is recognized by burning with a pale flame in air or oxygen, the flame depositing water on any cold object held above it.

---

**III. CARBON DIOXIDE GAS (CARBONIC ANHYDRIDE).\***—When carbon was burnt in oxygen (Exp. 5), a gas called carbon dioxide remained in the jar; the gas may be prepared in this way, but a much more easy

---

\* Formerly called carbonic acid, a name objectionable on chemical grounds.

## 26 PREPARATION OF CARBON DIOXIDE. [EXP. 13, 14.]

method consists in pouring hydrochloric acid upon some pieces of marble:



Chalk or limestone may be substituted for marble, but it does not answer so well.

EXP. 13.—Rinse out the apparatus used for preparing hydrogen, and place in it some small pieces of marble; fit into the india-rubber joint a delivery-tube bent at right angles, as shown in Fig. 5, then pour through the funnel-tube sufficient water to cover the marble and the end of the funnel-tube, and then strong hydrochloric acid until gas comes off with brisk effervescence. Place the delivery-tube in a jar with its end nearly touching the bottom, cover the mouth of the jar with a small disk of cardboard which has had a slit cut in it for the delivery-tube (or pass the delivery-tube through the brass cap of a deflagrating spoon), and allow the apparatus to stand for several minutes.

Carbon dioxide being much heavier than air, will soon fill the jar by "displacement," that is to say, it will collect in the lower part of the jar, and, by gradually rising in it, will lift out the air. Since this gas has the property of extinguishing a burning taper, it is easy to ascertain when the jar is full by holding a lighted taper just inside its mouth; if the taper is extinguished, the carbon dioxide has reached the top. Carbon dioxide gas, being very largely dissolved by water, is rarely collected over water, the process of "displacement" being very preferable.

EXP. 14.—Allow this jar of carbon dioxide to stand uncovered, and with its mouth upwards for a few minutes, then place in the jar a burning taper; the carbon dioxide is shown to be still present in the vessel by the immediate extinction of the taper. Then hold the jar

FIG. 5.




## EXP. 15-17.] PROPERTIES OF CARBON DIOXIDE. 27


for several minutes with its mouth downwards; on testing with a lighted taper, only air will be found in the vessel. These experiments prove that carbon dioxide is heavier than air, since it remains in a vessel which is open only above, and falls out of one which is open below.

EXP. 15.—Since this gas is so much heavier than air, it can be poured from vessel to vessel like water. This may be shown by pouring carbon dioxide from a bottle filled with the gas into a jar full of air, the latter being somewhat the smaller. The bottle is gradually tilted a little beyond the horizontal position, with its mouth over that of the jar. After holding it in this position for a short time, it may be proved by a lighted taper that the gas has left the bottle and is present in the jar.


FIG. 6.



EXP. 16.—Pour a little lime-water from a test-tube or small beaker into a jar of carbon dioxide, and shake the liquid round in the jar. The lime-water will at once become milky, owing to the lime which is dissolved in the water being converted by the carbon dioxide into common chalk (calcium carbonate), and this, being an insoluble substance, remains mixed as a white powder (“precipitate”) with the water:



EXP. 17.—Carbon dioxide gas readily dissolves in cold water, forming a liquid which probably contains carbonic acid:



The solubility of the gas in water may be proved by displacing the air from a bottle, previously half filled with cold water, by carbon dioxide; then tightly closing the mouth of the bottle with the wet hand, and shaking vigorously for a short time: the bottle will adhere to

the hand, owing to a partial vacuum being produced by the absorption of the gas by the water. A further proof is afforded by dipping the end of the delivery-tube employed in Exp. 13 to the bottom of a beaker containing water, so as to cause the gas to bubble up through the liquid. After the bubbles have passed for several minutes, the water may be shown to contain carbonic acid by pouring some of it into a test-tube and adding a little lime-water, which will cause a milkiness;\* also, by adding to another part of the carbonic acid solution several drops of blue litmus solution, or dipping into it a piece of blue litmus-paper, which will become red, indicating the presence of an acid. If this liquid containing carbonic acid be tasted, it will be found to possess a taste resembling that of soda-water, and in fact soda-water is merely water which contains a large quantity of carbonic acid, as may be proved by examining it with lime-water and litmus. If some of the water containing carbonic acid be boiled in a test-tube, the carbon dioxide gas is driven off again: the bubbles of gas are seen rising in the water long before the latter boils, and after the liquid has been boiled briskly for several minutes, it may be proved to be free from carbonic acid by giving no milkiness on addition of lime-water, and by not changing the color of blue litmus-paper or solution.

Exp. 18.—Dilute some lime-water, contained in a small beaker, with an equal quantity of distilled water, and allow the carbon dioxide gas to bubble through it as in Exp. 17; a milkiness will be produced owing to the formation of chalk (Exp. 16); but if the gas is allowed to bubble for several minutes through the liquid; the milkiness will gradually disappear, since the chalk dissolves entirely in the carbonic acid which is formed by the carbon dioxide dissolving in the water. On boiling some of this clear liquid it again becomes milky, since the carbon dioxide is driven out of the water by

---

\* Sometimes the milkiness disappears, for reasons explained in Exp. 18, unless much lime-water is added.

heat, and therefore the chalk can no longer remain dissolved. It will be found, on pouring out the water, that part of the chalk remains adhering to the inside of the tube, whence it may be removed by pouring in a few drops of hydrochloric acid. The above experiment explains the origin of the coating or "incrustation" of chalk inside kettles and steam-boilers in which chalk-water is boiled. Such water contains chalk dissolved by carbon dioxide gas present in the water; this gas is driven off when the water is boiled, and the greater part of the chalk separates upon the sides of the vessels.


*Tests for Carbon Dioxide.*—It is evident that the presence of carbon dioxide gas is shown by its properties of extinguishing a burning taper, and turning lime-water milky: these are the ordinary "tests" for carbon dioxide. We may now proceed to employ them, to prove that carbon dioxide is evolved from our lungs during the process of respiration, and also that it is produced by a burning candle.

EXP. 19.—Invert a bottle full of water in a pan of water, and fill it with air from the lungs by blowing the breath out through a glass tube, one end of which is dipped into the water and held beneath the mouth of the bottle. In order to obtain air *from the lungs*, a full breath should be drawn, and the nose then closed by pinching it with the finger and thumb. Before allowing any air to pass up into the bottle, a portion should be breathed out through the tube, so as to replace the air contained in the mouth and in the tube by air from the lungs; the remainder of the breath is then allowed to bubble up into the bottle. Close the bottle, remove it from the pan, and introduce into it a lighted taper, the flame will be immediately extinguished.

Now blow air *from the lungs* (obtained as just described) through a glass tube into lime-water contained in a small beaker, the lime-water will become milky.

EXP. 20.—Fasten a small piece of candle or wax taper upon the deflagrating spoon, and place it alight in a bottle of air the mouth of which is closed by the brass plate (Fig. 7). After burning for a short time the flame

will be extinguished, and if relighted will be again extinguished when placed in the jar. Now pour in some lime-water from a small beaker, and shake it round in the bottle, the liquor will be rendered milky.




A similar experiment may be performed, substituting the flame of coal-gas which is burnt from a bent tube for that of a candle, and closing the mouth of the bottle with a piece of cardboard. The extinction of the flame, and the milkiness then produced by shaking lime-water in the bottle, will prove the production of carbon dioxide by the combustion.

Hence carbon dioxide gas is constantly being introduced into the air by respiration and combustion, and we should therefore expect to be able to detect its presence in air by the above tests. It is manifestly not present in sufficient quantity to extinguish a burning taper, but the presence of carbon dioxide in air may be shown by lime-water in the following way:

EXP. 21.—Pour some clear lime-water into a watch-glass (or better a clock-glass), and allow it to stand for a few minutes in the air; a film of chalk will gradually form on the surface, and will be seen as white flakes when the water is stirred.

III. *a.*—*Nitric oxide gas* may be prepared in the apparatus used for making carbon dioxide (Fig. 5, p. 26).

EXP. 22.—The pieces of marble are removed and the apparatus washed out. Some scraps of copper (copper clippings or turnings) are then placed in the flask, and nitric acid diluted with an equal measure of water poured in:



A reddish-brown gas soon fills the inside of the vessel, and should be allowed to bubble off for a time through water; it may then be collected over water as directed for hydrogen gas. Nitric oxide is colorless, but it forms a reddish-brown gas (chiefly nitrogen tetroxide) when mixed with free oxygen. This is shown by filling a jar with the gas over water, and then allowing it to stand with its

mouth open in the air; the entrance of the oxygen of the air at once produces red fumes. The appearance of red fumes in the preparation vessel is thus explained, since the vessel is at first filled with air with which the first portions of the gas mingle.

IV. AMMONIA GAS.—The familiar smell of common “smelling salts” is due to ammonia gas which is constantly being given off from the solid “carbonate of ammonia” contained in the bottle. The gas is thus evolved only very slowly; it may be made to come off much more rapidly if the “carbonate of ammonia” is mixed with lime and the mixture is then gently heated; sal ammoniac is usually employed instead of the carbonate of ammonia:




FIG. 8.



EXP. 23.—Powder some ammonium chloride (sal ammoniac) in a mortar, and mix with it thoroughly on a sheet of paper about an equal quantity of slaked lime in fine powder. Pour some of this mixture into a small flask (Fig. 8) until it is about one-third filled, and close the neck of the flask with a tightly fitting perforated cork, into which is inserted a straight piece of glass tube 8 or 9 inches long. Heat the mixture gently by placing the flask upon a piece of wire-gauze on a tripod-stand and putting underneath it a lighted rose-burner. Ammonia gas will soon be smelt issuing from the end of the tube, and, since it is much lighter than air, may be collected by “displacement,” as described in Exp. 10. To ascertain when the vessel is filled with the gas it is only necessary to hold at the mouth of the jar a piece of moist red litmus or yellow turmeric paper; since ammonia gas changes the color of the former to blue and of the latter to reddish-brown, it is easy to see whether



it has reached the mouth of the jar by observing whether any change is produced in the color of the paper.

Ammonia gas does not burn continuously in air at the ordinary temperature, but it burns readily either in strongly heated air or when lighted in oxygen gas. Show this by holding the end of the delivery-tube, from which a stream of the gas is issuing, in the top of a Bunsen-flame; a pale yellowish-green flame of burning ammonia will be seen; and if the end of the tube is dipped into a jar of oxygen, the  $\text{NH}_3$  may be inflamed as it issues into the oxygen gas.

Ammonia should always be collected by displacement, since it is extremely soluble in water, and therefore its collection over that liquid would lead to great waste of the gas. For this reason, also, the vessels in which ammonia is to be collected should always be perfectly dry inside.

EXP. 24.—Place a jar full of ammonia mouth downwards in a vessel of water, and gently shake the jar so as to agitate the water at its mouth; the water rapidly absorbs the gas and rises in the jar to fill the space formerly occupied by the gas.

If a little water be rapidly poured into a jar of ammonia by momentarily partly opening it and at once covering it again with a glass plate or with the hand, and the water be then shaken in the jar, the water, owing to its having absorbed the gas, will, when poured out, be found to have acquired the smell and behavior with litmus and turmeric papers which characterize the gas. This liquid is in fact weak "*Liquor Ammoniae*," a solution prepared in large quantities by letting ammonia gas bubble for some time through cold water.

EXP. 25.—Pour into a glass jar a little strong hydrochloric acid, close the jar with a glass plate, and shake the acid about inside the jar, hydrochloric acid gas will thus be liberated; the liquid may then be allowed to run out by slipping aside the glass plate for a moment. Place this jar in an inverted position over a jar containing ammonia gas, and covered with a glass plate (Fig. 9,

## EXP. 26.] PREPARATION OF CARBON MONOXIDE. 33

a), then withdraw the glass plates, so that the mouths of the jars are in contact (Fig. 9, b), and the hydrochloric acid and ammonia gases can freely intermingle. Dense white fumes of solid ammonium chloride will immediately be formed :




This experiment may also be performed by dipping a glass rod into some strong hydrochloric acid, and holding it in ammonia gas as it issues from the delivery-tube of the apparatus, or in a jar previously filled with the gas ; the same white fumes will at once appear.


*Tests for Ammonia Gas.*—Ammonia gas may be readily recognized by its pungent smell ; by turning moistened red litmus-paper blue, and moistened turmeric-paper reddish-brown ; and also by giving white fumes with a glass rod moistened with *strong* hydrochloric acid.

*Gases are frequently made to pass through certain liquids, in order to free them from impurities before they are collected ; this is termed "washing" a gas. The preparation of carbon monoxide gas from oxalic acid will serve to show how this process is performed.*


**V. CARBON MONOXIDE.**—This gas is produced when carbon dioxide gas is made to pass over red-hot charcoal :



It is often thus formed in open grates, and is seen burning at the top with its characteristic blue flame. Carbon monoxide is usually prepared by heating solid oxalic acid with strong sulphuric acid, when a mixture of carbon monoxide and carbon dioxide is given off :



**EXP. 26.**—Place a little solid oxalic acid in a test-



#### 34 PROPERTIES OF CARBON MONOXIDE. [EXP. 27.]

tube; pour upon it strong sulphuric acid\* sufficient to cover it to a depth of at least half an inch, and heat the mixture. After a short time effervescence will be noticed, owing to gases being evolved; hold in the mouth of the tube a glass-rod freshly dipped into lime-water, the drop of lime-water hanging upon its end will become milky, showing that carbon dioxide is one of the gases evolved. Hold a burning taper to the mouth of the test-tube, a blue flame will be seen caused by the carbon monoxide gas burning in the air.

In order to get rid of the carbon dioxide gas which is mixed with the carbon monoxide, the mixed gases are "washed" with solution of caustic soda; this liquid absorbs the carbon dioxide readily, but allows the carbon monoxide to pass on:



The way in which this washing is effected is explained in Exp. 27.

EXP. 27.—Heat the mixture of oxalic acid and strong sulphuric acid in a flask (A) fitted as shown in Fig. 10, and pass the gases either into a wash-bottle (a small Woulffe's bottle, or a broad-necked bottle), fitted as shown in B, and containing caustic soda solution; or through a tube (a) containing fragments of quicklime, or a U-tube (b) filled with fragments of caustic soda or with fragments of pumice-stone moistened with strong caustic soda solution. The carbon monoxide gas thus more or less perfectly freed from carbon dioxide may be collected over water, and will be found to give either no milkiness with lime-water, or a slight milkiness if the stream of gas has been so rapid that the caustic soda has


---

\* Strong sulphuric acid is a very corrosive liquid, and great care must be taken not to get any upon the skin or clothes; should any of this acid or of any other acid get upon the skin it must be at once washed off; if it should accidentally be spilt upon the clothes, the part must be rubbed with ammonia solution. If the acid has remained for some time on the clothes it will produce a red stain, which will be removed by ammonia solution unless caused by nitric acid.

not been able to absorb the carbon dioxide completely. By heating a formate with strong sulphuric acid carbon monoxide alone is given off, and is thus readily obtained free from carbon dioxide.


Carbon monoxide resembles hydrogen in being inflammable and in extinguishing a burning taper; it also explodes when mixed with oxygen or air, hence before

FIG. 10.



collecting a cylinder of the gas for experiment, ascertain that the gas coming off from the apparatus is free from air by collecting a small test-tube full and proving that it burns quietly.


EXP. 28.—Push a burning taper up into a cylinder filled with carbon monoxide; the gas will burn with a blue flame at the mouth of the jar, but the taper will be extinguished. As soon as the gas has ceased to burn inside the cylinder, pour in a little lime-water and shake it about; the liquid becomes milky, showing that by the combustion of carbon monoxide in the oxygen of the air carbon dioxide gas is produced:



*Tests for Carbon Monoxide.*—Carbon monoxide is recognized by burning with a pale-blue flame in the air, producing carbon dioxide, which renders lime-water milky.

V. *a. Chlorine Gas* may be made in the apparatus employed for the preparation of carbon monoxide; the washing-bottle may either be dispensed with or may be used containing a little water.

Exp. 29.—Place some manganic oxide, powdered, or better in small lumps, into the flask A (Fig. 10, p. 35), pour upon it some strong hydrochloric acid mixed with about one-third its measure of water, and heat gently in a draught-cupboard or out of doors in the open air. A greenish-yellow gas is evolved, which may be collected by displacement like carbon dioxide, since it is much heavier than air:




The gas has a very destructive action on the lungs, and must on no account be inhaled; it is usually recognized by its yellowish-green color, its peculiar smell, and by its property of bleaching moist vegetable colors. This last property is shown by placing in a jar of the gas a piece of moistened litmus-paper or fabric dyed with madder ("Turkey red"); the colors of both will be destroyed. A burning wax taper plunged into a jar containing chlorine continues to burn with a very smoky flame; oil of turpentine, introduced into the gas by moistening a strip of filter-paper with the warm liquid, catches fire of itself and gives rise to dense smoke. A piece of Dutch-foil or copper-leaf also burns when dropped into chlorine.

*Tests for Chlorine.*—Chlorine gas is recognized by its yellow color, its smell, and its power of bleaching moistened litmus-paper.

V. *b. Hydrogen Chloride, or Hydrochloric Acid Gas.*

Exp. 30.—Place in the cleansed flask A (Fig. 10), which was used for the preparation of carbon monoxide, some lumps of sodium chloride (common salt), obtained by breaking up a mass of the melted powder, or by breaking a piece of rock-salt; pour upon it strong sulphuric acid and heat gently. Hydrochloric acid gas is evolved:



and being heavier than air, may be collected by displacement in the same way as carbon dioxide. The gas fumes strongly in moist air, turns moist blue litmus-paper red, dissolves easily in water (Exp. 24) giving an "acid" liquid (hydrochloric acid), which, like the gas, turns blue litmus red.


*Tests for Hydrochloric Acid.*—This gas is known by fuming in the air, turning moist blue litmus red, giving white fumes with ammonia gas, and yielding when dissolved in water a milky liquid on addition of silver nitrate solution, which does not become clear on adding nitric acid.

VI. DISTILLATION.—This process is employed to separate liquids which boil at a comparatively low tem-

perature, either from solids, or from other liquids which are not converted into vapor at all or only at much higher temperatures. It consists in boiling the liquid and cooling ("condensing") the vapor, which is thus given off, again into a liquid (the "distillate"), the non-volatile solid or liquid substances present being thus left behind in the vessel in which the liquid is boiled. As examples of this process, the purification of common spring-water from the solid substances dissolved in it, and the preparation of nitric acid, may be performed.

A. *Distillation of Water.*—EXP. 31.—Pour into a clean retort *a* (Fig. 11) some tap-water through a funnel placed in the mouth, or in the tubulure (*i. e.*, the opening for the stopper or cork) if, as is better, a tubulated

FIG. 11.




retort is employed. Support the retort, whose bulb has been about half-filled with water, in a retort-stand or upon a tripod, with its neck sloping downwards and dipping into a small, clean flask or bottle *b*, which is partly immersed in cold water contained in an evaporating basin. Cover the bulb of the flask with a broad strip of filter-paper, whose ends dip into the water contained in the dish; or instead of cooling the flask cool the neck

of the retort, by wrapping round the lower part of it a piece of filter-paper *c*, and round this a piece of wet string or tow *d*; then arrange a funnel *e*, with its mouth partly stopped, so as to drop cold water on the upper part of the filter-paper; this water will be drained off by the string *d*. On carefully boiling the water in the retort steam passes into the flask, and is there condensed to "distilled water." The first few drops should be thrown away, as they are apt to be impure from rinsing the retort neck and flask. Whilst this process of distillation is going on, add to some tap-water contained in a test-tube several drops of nitric acid and some silver nitrate solution, and notice that the water becomes milky; a separate portion in another test-tube will also be found to become milky on addition of some ammonia and ammonium oxalate solution. These changes are due to the presence in the water of certain solid substances dissolved in it. If the distilled water from the flask be examined in the same way it will remain clear, showing that these substances have been removed by distillation.

*Tests for Distilled Water.*—Good distilled water should remain quite clear when to separate portions of it are added solutions of ammonium oxalate, silver nitrate, barium chloride, and ammonium sulphide; these tests prove the absence of calcium, chlorides, sulphates, lead, and iron respectively. It should also leave no residue when evaporated.

B. *Nitric Acid.*—EXP. 32.—Clean the retort from Exp. 31 by rinsing out the bulb with a little dilute hydrochloric acid, and then thoroughly with water; let it drain for a few minutes, and place in it some solid potassium nitrate (nitre); then pour upon this, through a funnel placed in the neck (or tubulure), a quantity of strong sulphuric acid about equal in weight to the nitre used, and allow the acid to drain off the neck (in case a non-tubulated retort is employed), by supporting it for some time in an upright position. Proceed to distil just as in the preceding experiment; a thick yellow oily

liquid will trickle down the neck of the retort into the flask; this liquid is somewhat impure nitric acid, and will be found to fume strongly in the air:



*Test for Nitric Acid.*—Place some small pieces of copper in a test-tube, pour upon them a little of this nitric acid; on warming gently reddish-brown fumes will appear in the tube. (See Exp. 22, p. 30.) This property of giving red fumes when treated with copper is often used as a test for nitric acid.




## SECTION II.

**PREPARATION AND USE OF APPARATUS REQUIRED  
IN ANALYSIS.**

*Notes.*—Refer to the list in paragraph (494),\* which shows what apparatus is required for each student, and mark all apparatus as directed in the note at the end of the list.

1. *The Bunsen Burner.*—By far the most useful lamp for general heating purposes is the Bunsen burner (Fig. 12). It is a gas lamp so constructed that coal gas entering through the tube *a* may be burnt mixed with a proper proportion of air which flows in through the holes at the foot of the burner; the oxygen of the air

FIG. 12.



which is thus mixed with the gas burns the carbon in the interior of the flame. Accordingly, the luminosity, which is believed to depend upon the existence in the flame of unburnt carbon or carbon compounds, is destroyed. The Bunsen flame is therefore useful, because it deposits no soot upon any cool object which is being heated in it, and also, because it

is much hotter, owing to the more complete combustion of the gas, than any of the ordinary flames. Its high temperature, non-luminosity, and colorless appearance also render it very valuable for producing flame colorations, as will be seen hereafter. The lamp should be provided with some means for partly or entirely closing the air-holes when requisite; this is usually effected

---

\* Numbers occurring in the text which are inclosed in brackets refer to the paragraphs commencing in this section. For convenience in reference, paragraph numbers are printed in thick type, and at the head of each page will be found, inclosed in square brackets, the numbers of the paragraphs it contains.

either by a loose perforated ring which is slipped over them, or by making the tube *b* turn round on its long axis. When a small flame is being employed, the air must be partly shut off, else the flame "burns below."

When the lamp is to be used, a piece of tightly fitting india-rubber tubing  $\frac{1}{8}$ ths of an inch in diameter is pushed over the end of the tube *a*, and the other end of the tubing is slipped over the tube which supplies gas to the working bench, the gas-tap is turned on, and as soon as the gas is distinctly smelt issuing from the tube *b*, it is lighted. The flame should be almost perfectly colorless, and give scarcely any light. Occasionally the gas will burn at the bottom instead of the top of the tube *b*; this usually happens when the flame is turned down very low, or the supply of gas is insufficient; also when the quantity of air admitted through the holes is too great, or the burner is lighted too soon after turning on the gas-tap. It may be at once detected by looking through the holes in the base of the lamp, when the luminous flame will be seen burning from the small gas jet inside; the flame produced whilst the gas is "burning below" is also characterized by being long and somewhat luminous, and by emitting a very unpleasant smell. In such a case the flame should be at once extinguished by pinching the india-rubber tubing close to the burner, and the gas should be relighted after escaping for a few seconds: the momentary stoppage of the gas-stream by suddenly striking the india-rubber tube upon the bench with the hand will also frequently cause the flame to rise to the top of the burner without extinguishing it. Closure of the air-holes serves the same purpose, but this must be done with care, as after the flame has burnt below for a short time the metal becomes too hot to be touched. The risk of the gas burning below is prevented by slipping upon the top of the burner a tightly fitting thimble, with a hole in its top somewhat less in diameter than that of the burner.

2. *For diffusing heat over a large surface, the "rose-burner" is very useful; it is a small perforated cap c (Fig. 12) made of iron, which, when placed upon the*

top of the tube *b*, yields a small circle of flames; the rose-burner is put on and removed by grasping a small iron peg in its centre with crucible-tongs; it is of course very hot after being removed, and should never be handled, or placed upon wood, until it is cool.

3. *The flame of a spirit-lamp* is occasionally employed instead of the Bunsen flame, but for general purposes the spirit-lamp should only replace the Bunsen burner where coal-gas cannot be obtained.




FIG. 13.

The spirit-lamp (Fig. 13) consists of a glass vessel containing methylated spirit, into which dips a cotton wick supported by means of a brass, or better, a stoneware wick-holder. When not in use the wick should be always covered

with the glass cap to prevent evaporation of the spirit. If the spirit is tolerably free from resinous matter, its flame will be non-luminous, and will deposit no soot upon a cold object.

4. *The blowpipe* is frequently used to produce a small but very hot flame from the flame of a gas-lamp, spirit-lamp, or candle, by blowing through it a fine stream of air from the mouth.

In order to obtain the "blowpipe flame" from the flame of a Bunsen burner, the blowpipe is held by the




FIG. 15.

FIG. 14.

right hand, with its finely pierced tip *a* (Fig. 14), resting on the edge of the burner, and just inside the flame (Fig. 15); the mouth-piece *b* is then taken between the lips, and after blowing out the cheeks to their full extent, the air contained in them is

forced out through the jet *a*, and causes a small-pointed tongue of flame to issue from the side of the gas-flame. The chief difficulty in learning to use the blowpipe properly is experienced in acquiring the habit of keeping up the blast of air for some time uninterrupted by the breathing. A little patient trial will, however, soon remove this difficulty, if it is borne in mind that the cheeks must be kept constantly inflated with air, and that the air must be forced through the blowpipe *by the pressure of the cheeks alone and not by the action of the lungs*, breathing being carried on meanwhile through the nose, and the mouth being occasionally replenished with air from the throat just before breathing out the air from the lungs through the nose.

It is frequently necessary to have both hands free whilst using the blowpipe; this may be secured by resting the jet *a* on the top of the burner, and supporting the other end *b*, by the lips alone, as shown in Fig. 15; or the blowpipe may be rested on a support of convenient height, such as the ring of a retort-stand properly adjusted.

The bright flame obtained by closing the air-holes of the burner is much better suited for use with the blowpipe than the ordinary non-luminous flame. A burner with an elliptical orifice which gives a flat flame is commonly substituted for the ordinary round flame of the Bunsen burner; it is easily obtained by slipping a brass tube down the tube of the burner.

A further account of the uses of the blowpipe is given in par. 30.

5. *Glass tube or rod is cut* by laying it upon a flat surface, and making a deep scratch with the edge of a three-cornered file at the point to be cut. The glass is then held with both hands, one on either side of the scratch and close to it, and a gentle pressure is

FIG. 16.



exerted upon the glass as if trying to break it across (Fig. 16). If the file-mark has been made sufficiently deep, the glass will readily break at the scratch; the sharp edges of a rod or tube should always be at once rounded by holding them in the Bunsen or blowpipe flame until they are partly melted, or by rubbing them with the face of a file.

6. *Glass tube is bent* by holding the part to be bent in the upper edge of a common fish-tail gas-flame, so as




FIG. 17.

to heat at least two inches of the glass (Fig. 17). The Bunsen flame must never be employed for bending glass-tubing; it produces a most unsatisfactory bend. The tube is supported by holding it with both hands, one on either side of the flame, and

whilst being heated it is constantly turned slowly round on its axis so as to heat all sides equally. As soon as the glass is felt to be soft and pliable, it is taken out of the flame and quickly bent to the required angle. The heated part must not be allowed to touch anything until it is cold; the soot is then removed from it by a cloth or piece of paper. A bend, if properly made, should be a curve, and should not alter the bore of the

FIG. 17 a.




FIG. 17 b.

tube (Fig. 17 a); if a sharp angle is made, the bore will be narrowed, and the bend, besides being unsightly, is very liable to break under a small strain (Fig. 17 b).

*Glass rod may be bent in the Bunsen or blowpipe flame.*

7. *Glass tube is drawn out* by holding it with both hands as for bending; and whilst turning it constantly round on its long axis, the part to be drawn out is strongly heated in the Bunsen, or better, in the blowpipe

flame (see Fig. 15); when the glass is well softened, the two ends are slowly pulled asunder in opposite directions.

8. *Corks are bored* usually by means of brass cork-borers, which are used as punches. A borer is selected of the same size as the glass tube which is to be inserted into the cork, rather less than greater; the cork is then pressed against a wooden surface (best against the upright edge of a thick bench or table), and the perforation made in it by pushing the borer through whilst constantly turning it upon its long axis

FIG. 18.



(Fig. 18). Caution and practice will enable the student to make a clean straight hole without damaging the surrounding parts of the cork. In boring a single hole through a cork, the easiest way to make it straight is to bore from the centre of one end halfway towards the other, then reverse the cork and bore a hole to meet this from the centre of the opposite end.

A perforation may also be made by pushing a sharply pointed round file carefully through the cork.

The round file is required for smoothing the interior of holes made by the cork-borer, or for slightly enlarging them when they are too small; whilst doing this, great care must be taken to leave the hole round in shape, and not to enlarge it so much that the glass tubing when inserted fits loosely.

In perforating india-rubber stoppers, the borer used must be sharp, and must be kept well wetted with water, or better with solution of caustic soda or potash.


The edge of the cork-borer is sharpened when necessary by rubbing the outer part of the edge obliquely with the face of a fine-toothed three-cornered or flat file.

9. *Sulphuretted Hydrogen Tube*.—A tube of the requisite shape will have been used for the experiments in Section I; it is seen as a bent delivery-tube in Figs. 4


and 5 (pp. 24, 26). If not already made, bend a piece of glass tubing (6) ten or twelve inches in length at right angles into the shape shown in Fig. 17 *a*, the shorter limb being about two inches long. This tube is to be kept for passing sulphuretted hydrogen gas through liquids.

10. *Small Ignition-Tubes*.—A piece of hard glass tubing,\* perfectly dry inside and out, not much less than the eighth of an inch in internal diameter, is drawn out at its middle point by heating it strongly in the blowpipe flame, as shown in Fig. 15, p. 42. The narrowed portion of the tube (Fig. 19) is then cut across (5) at its

FIG. 19.



middle point, and by heating the conical part *a*, the narrow tube may be drawn off, and a small closed tube is obtained (Fig. 19 *a*). If the closed end is strongly heated in the blowpipe flame and blown into whilst hot, it may be expanded into a small bulb. (See Fig. 36, p. 133.)

FIG. 19 *a*.

11. *Glass Stirring Rods*.—Three or four glass rods are made by cutting (5) a length of solid glass rod into

FIG. 20.



pieces, some about seven inches long, others of about half that length. Any small projections are filed or chipped off, and both ends of each rod are then rounded by holding them in the upper part of the Bunsen flame, or better in the tip of the blowpipe flame, and turning the rod constantly round on its long axis until it

\* To distinguish hard glass from soft (lead) glass, hold the tube toward the light with one end near the eye; with hard glass the ring of glass will appear green or colorless, with soft glass black or dark.

becomes red hot at the end (see Fig. 20); the sharp edges are thus partially melted. The end of the rod must not be allowed to touch anything until it is cool.

11 a. Sometimes a very thin glass rod is required; this may be made from a stouter rod by softening it in the blowpipe flame (Fig. 15, p. 42), and then quickly drawing it out until the required fineness is produced.

12. *Mounted Platinum Wires*.—The platinum wire used here must not persistently color the flame green.


FIG. 21.



Draw out a piece of glass tube six or seven inches in length at its middle point (Fig. 15, p. 42) into the shape shown in Fig. 19. Cut it across (5) at the middle of the narrow portion. Each piece of glass thus obtained serves for the handle to a wire. Break off the narrow part of the tube until it extends only about a quarter of an inch from the shoulder (a, Fig. 19), and into the narrow opening insert the end of a piece of platinum wire about two inches in length and bent round at its other end into a circle about the eighth of an inch across by rolling it round a piece of thick wire; then hold the end of the wire surrounded by glass in the Bunsen or blowpipe flame until the glass melts and thickens around it, fixing it firmly when cold (Fig. 21). At least two wires should be mounted in this way. They are then rendered less liable to be lost and much more easy to handle. The wire may also be mounted by softening the end of a short piece of *glass-rod in the blowpipe flame*, and sticking the wire into it.

13. *Wash-bottle*.—A thin flat-bottomed flask is chosen (best of the triangular shape shown in Fig. 22) and of 16 or 18 ounces capacity; the neck must not be less than an inch in diameter. Procure a sound cork which is slightly too large to enter

FIG. 22.





the neck, soften the cork by placing it upon the floor and rolling it backwards and forwards under the foot with gentle pressure; when thus softened the cork must fit tightly into the flask. A vulcanized india-rubber stopper is much more durable for this and most other chemical purposes; it is perforated by a sharp well-wetted cork-borer, or by a wetted round file. Two pieces of glass-tubing rather longer than would be required for the tubes *a* and *b* are then bent (6) into the form shown in Fig. 22. The ends of the tubes are, if necessary, cut off (5) to the right length, and their sharp edges rounded by holding them in the Bunsen flame or in the tip of the blowpipe flame, or by rubbing them with the face of the file.

Two parallel holes are then bored in the cork by means of a round file or with a proper sized cork-borer (8); the holes must be rather smaller than the glass tubes, and must not run into one another or to the outside of the cork. They are slightly enlarged, if necessary, by the round file. Into these holes the tubes *a* and *b* are then pushed with a twisting motion; if the holes have been made of the proper size the tubes must enter somewhat stiffly, but without requiring much pressure. If the holes have been carelessly made too large, the tubes may often be made to fit by slipping upon them little pieces of narrow india-rubber tubing, or by putting the pieces of india-rubber tubing into the holes in the cork before fitting in the glass tubes.

Upon the upper end of *a* is fitted a piece of small india-rubber tubing about an inch and a half in length, into the other end of which is pushed a short jet (*c*) made by drawing out a piece of glass tubing in the flame (Fig. 15, p. 42); its nozzle may be contracted, if necessary, by holding it *perfectly dry* in the flame for a short time. The neck of the flask should then be bound round with twine like the handle of a cricket bat, or tightly covered with a piece of flannel; this prevents the fingers from being burnt when water has been boiled in the flask and its neck has become heated by the steam.

The wash-bottle, when thus fitted up, is filled about

two-thirds with distilled water. Tap-water should never be kept in the wash-bottle.

A fine stream of water may then be obtained from the jet (c) by blowing down the tube (b); this stream serves for washing precipitates and for other purposes. A larger stream is obtained by inverting the bottle, when the water will flow out from the end of the tube (b), air entering meanwhile by the tube (a).

Hot water is frequently required for washing precipitates; to obtain this the wash-bottle is placed upon a piece of wire gauze supported on a tripod-stand, and is heated by the Bunsen flame issuing from a rose-burner, as shown in Fig. 10 (p. 35).

**14. Cleaning Apparatus.**—It is indispensable to the success of an analyst that all glass and porcelain apparatus should be scrupulously clean before being used.

Test-tubes, beakers, and porcelain dishes are washed in a stream of tap-water by using the test-tube cleaner (Fig. 23 b). Sometimes a little hot hydrochloric acid is requisite to remove stains, and occasionally it is necessary to heat a little strong sulphuric acid in a vessel in order to cleanse it from grease or other impurities; a little hot caustic potash or ammonia may also often be used with advantage. In fact in removing a substance from a vessel to which it strongly adheres, the student should always consider what the substance is and then remove it by a liquid in which it is easily soluble. Each article, after having been well washed and rinsed with tap-water, should be placed upside down in the small wicker basket to drain, or, if required for immediate use, should be rinsed out with a little distilled water. It should be borne in mind that apparatus must be washed as soon as possible after use, as after standing the surface is usually much more difficult to cleanse. The brush must be cautiously moved as it reaches the bottom of a test-tube, since as the glass is very thin the brush is easily pushed through it.

**Test-tube Brush.**—It will be found that the piece of sponge at the end of the test-tube

FIG. 23.



cleaner (Fig. 23 *a*) is not well adapted to cleanse the bottom of test-tubes and boiling tubes. A much more efficient end is given to the brush by removing the sponge and bending back the end of the wire stem upon itself at a point just above where the hairs commence (Fig. 23 *b*). By slightly curving the part of the stem carrying hairs, the brush will better adapt itself to curved surfaces such as those of porcelain dishes.

*Test-tubes*, whilst in use, are placed in the test-tube stand; boiling tubes may be supported in the wicker basket, or in a specially made stand. When washed they should be placed to drain mouth downwards in the wicker basket.

*Glass funnels* should have their narrow stems cut off to within about half an inch from the shoulder; the sharp outer edge must then be removed by rubbing it with the face of a triangular file. The inside of the narrow neck is best cleaned by washing it with a common tobacco-pipe cleaner, inserted from the shoulder of the funnel so as not to cut the hairs by the edge at the other end of the neck.

15. *Platinum foil and wire are cleansed* by boiling them in hydrochloric acid, and rinsing off the acid with water; the wire should then be strongly heated for some time in the blowpipe flame until, on being dipped into pure strong hydrochloric acid, it no longer colors the Bunsen flame. If the tip of the wire cannot be thus cleansed it should be cut off.

Commercial platinum often contains Barium, and the wire made from it therefore gives a green color to the flame; such wire is useless for flame coloration tests and for spectrum analysis.

It is best to keep the foil or wire in a small beaker or dish, containing strong hydrochloric acid diluted with sufficient water to prevent it from fuming; the platinum will thus be ready for use after having been rinsed with water.

16. *Before putting by apparatus* it should be made a rule to wash all glass and porcelain which is not in

actual use, and place it in the wicker basket to drain; the basket is put away with its contents.

All iron apparatus should be carefully dried, and must be kept in a dry place to prevent rusting. *On no account must metal apparatus be kept in the wicker draining basket.*

**17. Heating Glass and Porcelain.**—A few general precautions should be observed in heating glass and porcelain vessels, to guard against cracking them.

A vessel containing a liquid must never be heated by the flame above the level of the liquid inside.

A dry hot vessel must be allowed to cool before pouring in any liquid, or placing it on a cold surface. See also (18, 19).

**18. Porcelain dishes** are generally used for boiling liquids; they are supported on a tripod or retort-stand, and may be safely heated by the naked flame. *Porcelain crucibles* are used for containing solid bodies which are to be strongly heated; they are supported on a pipe-clay triangle placed upon the ring of a retort-stand or upon a tripod-stand: the flame should not be allowed to play steadily at once upon the bottom of the crucible so as to heat it suddenly, but should be constantly shifted by moving the burner, until the porcelain is hot. The crucible should also be allowed to cool slowly on the triangle, as contact with a cold body is very apt to crack it: the crucible and its cover whilst hot are handled by the crucible tongs.

**19. Glass vessels** require to be heated more cautiously than porcelain: a large naked flame must never be allowed to play for any length of time on any one part of the glass surface. In heating a test-tube or boiling tube, this "local heating" is prevented by holding the tube obliquely with the lower part in the flame (Fig. 1) and moving it gently up and down, or by constantly turning it round on its axis. Test-tubes are too narrow for boiling liquids in, the liquid being very apt to boil over. Small quantities may be boiled and larger quantities heated short of boiling in a test-tube, but the broader "boiling tubes" are best suited for this purpose.

Large glass flasks, such as the wash-bottle, are most safely heated by placing them on a piece of wire gauze on a tripod-stand (Fig. 10, p. 35), and heating with a rose-burner; in some laboratories a sand-bath is available, the flask is then heated by being placed on the surface of hot sand. Test-tubes, if not full of liquid, can be held by the neck whilst being heated, if the tube be held obliquely so that the fingers are not over the flame. All risk of burning the fingers is avoided by bending round the neck of the tube a slip of folded paper or of leather, and pinching the ends together close to the tube.

## SECTION III.

## ANALYTICAL OPERATIONS.

THE student, before commencing the analytical reactions, must become familiar with the operations which are constantly employed in Analytical Chemistry. The processes will be most readily and perfectly understood by reading the following descriptions, and performing the experiments given in illustration of them.

**20. Solution.**—Many solid substances, if stirred or shaken with water, gradually “dissolve” in that liquid; salt and sugar may be mentioned as examples. Other liquids may be employed instead of water, and if they cause the solid substance immersed in them to partially or entirely become liquid and mingle uniformly with the liquid, they are said to “dissolve” the solid, the liquid thus obtained being called a “solution” of the solid, and the liquid which dissolves the solid being termed the “solvent.” Thus water, when shaken with sodium chloride (common salt), dissolves it, yielding solution of sodium chloride, and water is therefore called a solvent for sodium chloride. Further, a solid which dissolves in a liquid is said to be “soluble” in that liquid; if it does not dissolve it is said to be “insoluble.”

The process of solution is more rapid when the solid substance is employed in the state of powder than when it is in large lumps, since a larger surface of the substance is thus exposed to the solvent by the powder; it is also much hastened by heating the solvent, which causes a more rapid circulation of the liquid over the solid, and also frequently increases the solubility of the substance.

Two kinds of solution may be distinguished. “Sim-

ple solution" occurs when a substance dissolves in a liquid without alteration in composition, the solution possessing therefore the taste, color, and other general properties of the solid, and yielding again the solid substance when the solvent is removed by "evaporation." Solution of sodium chloride in water is an example of a "simple solution."

"Chemical solution," on the other hand, is always attended by a chemical change in the substance to be dissolved, and since the solution therefore contains a substance differing in composition from the undissolved solid, the properties of the solution are usually not the same as those of the undissolved solid, and on removing the solvent by evaporation, the original substance is not obtained.

In dissolving solid substances in liquids, test-tubes, porcelain basins, or small glass flasks are generally employed.

EXP. 33.—Place a piece of potassium nitrate (nitre) in a small clean beaker, fill the beaker one-third with water,\* and stir the solid about in the water with a glass rod; the potassium nitrate will *slowly* dissolve in the water; if the liquid be heated in the Bunsen flame, the solution will be much more rapidly effected. Powder another piece of potassium nitrate by crushing it and then rubbing it in a mortar with the pestle; place this powder in a test-tube, pour in water, and heat the bottom of the test-tube; the potassium nitrate will dissolve much more rapidly than before, showing that solution is accelerated by powdering the solid and employing heat. Keep these solutions.

EXP. 34.—Powder a little copper sulphate (blue vitriol) in a mortar, place it in a small porcelain dish, half fill the dish with water, and heat it on a tripod or retort-stand. The *blue* copper sulphate dissolves, yielding a *blue* solution. Keep this solution.

These two experiments are both examples of "simple

---

\* Here, and in all subsequent cases, water must be understood to signify *distilled water*.

solution." Exp. 33 shows that a colorless solid gives a colorless solution. Exp. 34 proves that a colored solid gives a colored solution. This is generally true, and hence we can often infer the presence or absence of a colored substance in a solution by merely noting the color of the liquid. Moreover, if a drop of the potassium nitrate solution be tasted, it will be found to possess the same taste as the solid; chemists occasionally rely upon taste, as upon color, when examining "simple solutions."

EXP. 35.—Place in a test-tube a few small pieces of calcium carbonate (marble or chalk), pour upon them a little water and heat the tube; the calcium carbonate will be found to be "insoluble" in the water. Add to the water some hydrochloric acid, "effervescence," or escape of numerous small bubbles of gas, will occur; the pieces of calcium carbonate will meanwhile slowly diminish in size, and will at last entirely disappear in the liquid if sufficient acid is added.

EXP. 36.—Place in a test-tube a few small pieces of copper; on warming these with a little water they remain undissolved, but on adding to the water some nitric acid and heating, the copper slowly dissolves, giving off red fumes, and will be entirely dissolved if sufficient nitric acid is employed.

Experiments 35 and 36 are examples of "chemical solution;" the calcium carbonate is changed by the hydrochloric acid into calcium chloride, and this substance, *not the calcium carbonate*, remains in solution. The copper is changed into copper nitrate, which is then dissolved by the water. It will be noticed that in both these cases a gas is given off; this is a very usual, but not a universal, effect during "chemical solution;" the distinguishing fact is that the solid substance which is to be dissolved has undergone a chemical change in the act of passing into solution.

21. *Evaporation*.—When it is wished to obtain a substance, which is dissolved in a liquid, in the solid condition, the liquid is boiled away as vapor, or "evap-



orated;" the solid substance is then left behind in the vessel.

EXP. 37.—Pour the potassium nitrate solution from Exp. 33 into a porcelain evaporating basin, place the basin on a tripod-stand, and boil it over the Bunsen flame until the water has been nearly boiled away, then make the flame smaller, and heat until the water has quite disappeared; the solid potassium nitrate is left in the dish.

In evaporating a solution a small flame or a rose-burner should always be used towards the end of the process, and the flame should be moved about if necessary, else some of the substance will spirt out of the dish.

Often a part only of the liquid is evaporated for the purpose of "concentrating" the solution (*i.e.*, making it stronger by removing a portion of the solvent). A solution, when sufficiently concentrated, will often form "crystals" of the dissolved substance if allowed to stand till cold.

EXP. 37 *a*.—Concentrate the copper sulphate solution made in Exp. 34, and allow it to cool; if sufficient water has been evaporated, it will form crystals. Keep the copper sulphate in the dish.

22. *Precipitation*.—Two perfectly clear and transparent solutions, on being mixed together, often become more or less turbid and opaque, owing to a solid insoluble substance being formed in the liquid. A solid substance so formed in a liquid is called a "precipitate."

EXP. 38.—Pour into a test-tube some barium chloride solution, and then some ammonium carbonate solution; a white precipitate is formed owing to the production of insoluble solid barium carbonate. Keep this test-tube with the precipitate.

In the above instance barium chloride and ammonium carbonate separately dissolve easily in water, but if their solutions are mixed they yield by a chemical change two different substances,—ammonium chloride and barium carbonate; the former of these dissolves in water, but

the latter (like common marble) is insoluble, and therefore remains as a fine powder in the liquid.

Substances are very frequently removed or separated in analysis by thus causing them to form insoluble compounds or precipitates with other substances which are added for the purpose.

In producing a precipitate care must be taken that the two solutions are well mixed; this mixture may be effected by closing the mouth of the test-tube with the thumb and several times inverting it, or often by simply warming the bottom of the test-tube in the flame; mixture may also be effected by stirring with a glass rod or by pouring the liquid from one vessel to another; the last three methods are to be used when the liquid present is corrosive, and therefore must not be allowed to touch the skin. A precipitate which does not form readily is often caused to appear more rapidly on thoroughly mixing or agitating the liquid by the above means, or by heating it.

Precipitates produced with different substances differ much in appearance and properties; hence we frequently form a precipitate to show the presence of a substance or to separate one substance from another. Substances thus added to produce precipitates are called "reagents."

The appearance of a precipitate is usually described by its color and its condition; it is "flocculent" if it forms in flocklike masses, "crystalline" if in small particles which are seen to be crystals under a lens or microscope, "gelatinous" if jelly-like, etc. A slight precipitate causes only a "turbidity" in a liquid.

The color of a precipitate or liquid is often invisible or falsified by gaslight; if the color is to be seen at night, it should be examined with the light produced by a piece of burning magnesium ribbon.

Occasionally solid substances are used to produce precipitates; thus one metal is not unfrequently precipitated from its solution by immersing in the liquid another metal.

EXP. 39.—Dip a clean penknife blade into some of

the copper sulphate solution left from Exp. 37 a, to which a few drops of sulphuric acid have been added; after a short time the metal copper is precipitated from its solution, and covers the iron as a red film.

FIG. 24.




**23. Filtration and Decantation.**—It is frequently necessary to separate a precipitate from the liquid in which it is suspended. This is effected either by filtration or decantation.

**23 a. Filtration** is performed by pouring the liquid containing the precipitate upon some porous paper; the liquid itself runs through the pores of the paper, but the solid particles of the precipitate cannot pass through, they remain on the surface of the paper. The paper employed is called “filtering-paper,” and the liquid which runs through is termed the “filtrate.” A filtrate may frequently be colored by some substance *dissolved* in it, but it must always be perfectly clear and free from turbidity caused by solid particles *suspended* in it.

**EXP. 40.**—Measure a glass funnel from its shoulder to its edge along the sloping side (Fig. 24), cut a square

FIG. 25.



of filter-paper whose edge is rather less than double this length, fold it across first along the dotted line (Fig. 25 a), then again as shown by the dotted line in Fig.

25 *b*; this gives a square (Fig. 26 *c*) which at one angle has four free corners, these are removed by cutting with a pair of scissors along the curved dotted line shown in Fig. 25 *c*; the "filter" is now made and merely requires to be opened; by separating the edges, formed by cutting off the corners, so that they form a circle, three remaining on one side of the circle and one on the other, a little closed pointed paper bag is formed (Fig. 25 *d*); this is gently pressed *with dry fingers* into the dry funnel, and the folding altered if necessary till it fits tightly into the glass; it is then moistened all over with a little water from the wash-bottle, and is ready for use.

The preliminary moistening of the filter-paper must not be neglected, since if the liquid and precipitate are poured upon a dry filter, some particles of the precipitate get between the fibres, which then swell by the wetting, inclosing the solid particles and stopping the pores of the filter.

As an example of the process of filtration, the barium carbonate precipitate obtained in Exp. 38 may be filtered off. Pour the liquid containing the precipitate upon the filter, the funnel being placed with its neck in a test-tube, and the test-tube being supported in a test-tube stand (Fig. 26); care must be taken that the inside of the upper part of the test-tube is not wetted, else the space between the neck of the funnel and the inside of the tube is liable to become filled with liquid, and the egress of air being thus prevented the filtration is stopped. The funnel might also be supported upon the ring of a filter-stand, and the filtrate be received in a small beaker instead of in a test-tube. Take great care not to let the level of the liquid in the funnel reach the upper edge of the filter-paper. If the above directions have been carried out, and no hole has been made in the paper whilst preparing the filter, the filtrate will run

FIG. 26.



through perfectly clear, leaving the barium carbonate on the filter. Keep the funnel, filter, and precipitate.


Circular filters can be purchased ready cut, and only requiring to be folded twice to be ready for use. The filter must always be of such size that when placed in the funnel its edge is below the rim. Sometimes the filtrate runs through turbid; it can then usually be made quite clear by passing the whole, or the first portions, of the filtrate once or twice again through the same filter; but with many precipitates a double filter should be used, a double filtration being thus performed in one operation.

It should be borne in mind that hot liquids run through a filter much more quickly than cold, hence, if not inadmissible for other reasons, a solution should always be boiled before being filtered.

*Note.*—Common filter-paper contains a little calcium-carbonate, which is of no importance in ordinary analyses, and is only objectionable when a substance has to be examined for traces of calcium. To remove this impurity in special analyses, the filters are moistened in a funnel with dilute hydrochloric acid, and then well washed with distilled water.

**23 b. Decantation.**—Precipitates which settle rapidly and completely in the liquid in which they are formed may frequently be separated from that liquid without

the use of a filter. The vessel is allowed to stand at rest until the precipitate has completely settled, and the liquid is then carefully poured off ("decanted") by allowing it to run gently down a wetted glass rod which is pressed against the



edge or lip of the vessel (Fig. 27), the precipitate being left in the vessel. This process, if carefully performed, effects a fair separation of heavy precipitates.

The precipitate obtained by adding sulphuric acid to some boiling solution of barium chloride may be separated by decantation. Keep the precipitate in the test-tube.

**24. *Washing Precipitates.***—A precipitate which has been filtered from the liquid in which it was suspended has often to be washed until perfectly free from the liquid adhering to it; this “washing” is performed as described under 24 a if the precipitate has been separated by filtration, and by 24 b if by decantation.

**24 a. EXP. 41.**—Support the funnel, which contains the filter and barium-carbonate precipitate from Exp. 40, with its neck in a beaker or flask, and with the wash-bottle blow in a fine stream of hot distilled water, so directed by moving the jet with the fingers as to stir up the precipitate well; in this way fill the filter to within a short distance from its edge; let this water run through perfectly, then nearly fill the filter again in the manner just described, and repeat this process two or three times, letting the liquid run through perfectly each time before putting in a fresh quantity; the water running through from the third or fourth washing will usually be quite tasteless, and the precipitate and filter will be freed from everything which dissolves in water. Boiling, or at least hot, water should always be used for washing purposes, as the process is thus rendered much more rapid.

**24 b. *Washing by Decantation.***—Heavy precipitates, after having been separated from liquids by decantation, are washed also by the same process. The precipitate is shaken or stirred with distilled water, then allowed to settle; when the water has become quite clear it is poured off (23 b). By several times repeating this washing process with addition of fresh portions of distilled water, the precipitate may be entirely freed from adhering solutions.

The precipitate of barium sulphate remaining in the test-tube from the experiment at the end of 23 b may be thus washed by decantation.

25. *Drying Precipitates.*—A precipitate is often required in a dry condition after it has been filtered off and washed. It is dried on the filter and in the funnel by placing the funnel in a hollow tin cone or cylinder (filter-dryer, Fig. 28), and supporting this on a piece of wire gauze upon a tripod-stand over the flame of a rose-burner turned very low; or the filter-dryer may be placed on a gently heated sand-bath. The funnel is thus heated by the current of hot air, and rapidly dries the filter and

FIG. 28.



precipitate. Great care must be taken to regulate the heat and watch the process, so as not to burn the filter.

A more rapid method of drying the precipitate consists in spreading the filter, after having allowed it to drain for a time, upon a piece of wire gauze supported on a tripod-stand; a small flame from a rose-burner is then placed beneath the gauze, and the filter carefully watched to avoid burning it.

26. *Removing Precipitates from the Filter.*—When a small quantity of a moist precipitate has to be taken from a filter to test its behavior or closely examine its appearance, it is most readily removed by dipping the end of a glass rod into the precipitate; by touching a watch-glass or the interior of a test-tube with the end of the rod, a small quantity of the precipitate is deposited for examination.

If the precipitate is to be removed from the filter as completely as possible several methods are available; one or other must be chosen according as circumstances render it suitable.

a. The bottom of the filter may be pushed out through the neck of the funnel with a glass rod, which is small enough to pass easily through the neck (11 a), and the precipitate may then be washed down with a fine stream of water or other liquid from the wash-bottle into a vessel beneath.

b. Without breaking the filter the funnel may be held

with its neck horizontal, and the rim just inside the edge of a porcelain dish (Fig. 29), the precipitate is then washed out by directing a fine stream of water from a wash-bottle against the sides of the filter.

c. The filter and precipitate are allowed to stand for some time, so as to drain off as much water as possible; the filter is then carefully taken out of the funnel, partially dried if necessary by laying it upon several folds of filter-paper, and after removing the portions of the paper which contain no precipitate (more particularly the empty fold)\* it is spread out inside a porcelain dish; the liquid with which it is to be treated is poured upon it, and by shaking the dish so as to cause the liquid to move round and round, and occasionally carefully stirring the precipitate with a glass rod, the precipitate is washed off the paper without tearing the latter to pieces. The paper is then carefully removed by a glass rod.

FIG. 29.



d. If it is undesirable to add a liquid to the precipitate upon the filter, the filter and precipitate, after draining for a short time, are removed from the funnel, the filter is spread out upon a flat piece of glass and the precipitate carefully scraped off with a knife, or better with a small spatula. If the precipitate is required dry, the filter after removal from the funnel may be carefully opened and spread upon several thicknesses of filter-paper to drain. When there is a large quantity of the precipitate a sufficient quantity may be removed on the end of a spatula or knife without taking the filter out of the funnel. This method is usually the most imperfect, but is frequently the best for other reasons.

e. A precipitate has sometimes to be dissolved off the filter; the hot liquid used as a solvent may then be

\* Filters may be made without this fold, by cutting one of the round papers (Fig. 25d) through across one of the folded diameters, bringing the cut edges together, and sharply folding them over several times, pressing them with the nail; if properly made, such a filter is perfectly impervious to the precipitate.



poured upon the precipitate; it will run through the filter into a vessel below, taking with it the precipitate in solution: the liquid should be heated again and once more poured upon the precipitate, if the latter is not entirely dissolved; this reheating and returning of the liquid to the filter should be continued as long as anything is dissolved; any remaining portion of the precipitate must then be removed by a little fresh solvent.

*f.* A precipitate, if in small quantity, may also be rinsed off the filter with the liquid with which it is to be treated or dissolved. The funnel is placed with its neck in a test-tube, the liquid is poured into the filter from another test-tube, and the precipitate is quickly stirred up with the liquid with a glass rod thin enough to pass down through the neck of the funnel; the bottom of the filter is then pushed out through the neck by the glass rod, and the liquid carrying most of the precipitate will run through; if some of the precipitate remains on the filter, the same liquid is poured through the filter again into the other tube, and by thus pouring backwards from one tube to another all the precipitate may be removed.

*Note.*—The greater part of the precipitate may often be removed by a clean knife or spatula, and the remainder by some one of the foregoing methods.

EXP. 42.—Remove the precipitate of barium carbonate remaining from Exp. 41 from the filter by one of the above processes, using hydrochloric acid as a solvent for method *e*.

27. *Ignition.*—Solid substances occasionally require to be heated strongly or “ignited.” This process of ignition is employed with various objects.

1st. In order to see what effect strong heat has upon a substance, since by this means we can frequently detect its composition. For this purpose we use small tubes made from hard glass tubing (10).

EXP. 43.—Heat a little mercuric oxide in a small tube of hard glass; after a short time minute shining drops of the metal mercury will be seen on the upper part of the tube, and if then a small splinter of wood with a spark at its end be held in the tube, it will burst

into flame. The behavior of this red powder on being "ignited," therefore, shows it to contain the elements of mercury and oxygen.

Substances are often heated on platinum foil with the same object.

2d.—Ignition is also used as a means of separating from a solid mixture certain substances which are "volatile," or which, when heated, change entirely into vapor; these are driven off by heat from other substances, which are "non-volatile" (*i. e.*, can be strongly heated without changing into vapor), and which, therefore, remain behind.

EXP. 44.—Heat a small piece of ammonium chloride on a strip of platinum foil held in the flame by crucible tongs; the ammonium chloride, being volatile, will give off white fumes, which are best seen by removing it from the flame for an instant, and the solid substance will slowly disappear or "volatilize."

Heat in the same way a small piece of potassium chloride; this will melt, but will not disappear, since it is only slightly volatile. It will become solid again on cooling.

Hence, if we had a mixture of ammonium chloride and potassium chloride, and wanted to obtain the potassium chloride alone, we should strongly heat the mixture for some time, when the ammonium chloride would be removed as vapor, leaving the potassium chloride behind.

28. *Sublimation* is the process of heating solid substances and converting them into vapor, then cooling this vapor so as to "condense" it, or render it solid again. A volatile substance may thus be obtained free from other substances which are not volatile or only slightly so.

EXP. 45.—Heat a small piece of ammonium chloride in a tube of hard glass closed at one end and two or three inches in length; it changes into vapor, which again cools into solid ammonium chloride on the upper part of the tube. This is an example of the process

called "sublimation," and the solid substance at the upper part of the tube is a "sublimate."

**29. Fusion.**—Substances are often melted or "fused" together to cause certain chemical changes to occur between them. This melting or "fusion" is performed sometimes on platinum foil, but often in porcelain or platinum crucibles. For fusion in a crucible an intense heat is often requisite; this may be obtained by using the blowpipe flame, but the Bunsen flame will usually suffice if the crucible be surrounded by the conical filter-dryer (Fig. 28, p. 62) which steadies the flame and creates a draught.

**Exp. 46.**—Place upon a slip of platinum foil with bent-up edges, a mixture of powdered sodium carbonate and potassium nitrate, then a very small quantity of manganese chloride; hold the foil with crucible-tongs in the top of the Bunsen flame, or heat the under surface of the foil with the blowpipe flame. The mass will melt or fuse; it is kept in a melted state for a short time, and then allowed to cool, when a bluish-green mass is obtained. The production of this color is due to the presence of sodium manganate formed by the fusion; this peculiar color on fusion with sodium carbonate and potassium nitrate is produced only when manganese is present, and is, therefore, a test for that element.

**30. Use of the Blowpipe.**—The blowpipe is used, as already explained (4), to obtain an intensely hot flame, which may be made to play in a convenient direction upon any substance to be heated; it is very useful for analytical purposes, since by its skilful use the composition of an unknown substance, when ascertained by other means, can usually be confirmed; and with many substances (especially minerals) a complete, or at any rate sufficient, analysis can be made by the blowpipe alone. The way in which the flame is obtained has been already explained (4).

In a well-formed blowpipe flame two parts may be distinguished; the "inner flame," which is blue and pointed, and around and beyond this an almost colorless flame, "the outer flame." The inner flame is often called

the "reducing or deoxidizing flame," because it separates or "reduces" metals from their oxides and compounds, and in general "deoxidizes" substances; the outer flame is also known as the "oxidizing flame," because it changes metals into their oxides, and "oxidizes" many other substances. These two flames also produce other distinctive effects.

The "inner flame" is best obtained of large size by placing the tip of the blowpipe nozzle only in the very edge of the flame, or even just outside it; the "outer flame" is best obtained by putting the nozzle much further into the gas flame. It may also be noted that the "reducing flame" is furnished most readily by the luminous gas flame; the "oxidizing flame" is most easily and perfectly formed from a Bunsen flame which contains some air, and which is obtained by either leaving the air-holes at the foot of the burner entirely open or by only partially closing them.

There are several purposes for which the blowpipe is employed in analytical chemistry with which the student should become quite familiar; the following experiments will exemplify the most important of them.

31. EXP. 47.—*Borax Bead*.—Place a little borax, which has been finely powdered in a mortar, upon a watch-glass; moisten the loop of a piece of mounted platinum-wire (12) with water or in the mouth, and dip it into the powdered substance, then hold the loop with the adhering powder in the blowpipe flame until the borax melts; if this is insufficient when melted to form a bead which fills the loop, dip the loop with the melted substance upon it again into the powder, more powder will adhere and may be melted in the flame; in this way powder is to be constantly added until when held in the flame a liquid drop or bead is produced which completely fills the loop. This bead must be perfectly colorless and transparent on cooling. Keep the bead on the wire.

If the melted bead easily drops from the wire, either the loop or the bead is too large and must be lessened in size. The most convenient sized loop is one which is

rather less than one-eighth of an inch across; it is best made by rolling the tip of the wire round a piece of stout iron wire. If the bead is too large, remove a portion whilst it is in a fused condition by giving the wire a sudden jerk.

Certain substances when melted in this bead color it, and as a rule the color produced by each of these substances has some characteristic tint or peculiarity. The color of the bead should be examined after it has been heated, first in the outer and then in the inner flame, and must be noted both hot and cold. If the bead appears opaque, on account of too much of the coloring substance having been introduced, it can generally be made transparent by shaking off some of the fused borax, or by flattening the fused bead by gently squeezing it between a small pair of pincers.

EXP. 48.—Moisten the clear cold borax bead and dip it into some powdered ferrous sulphate, thus taking up *very little* of the powder upon it; then heat the bead and substance for a short time in the tip of the outer blowpipe flame, and examine the color of the bead by looking through it at a piece of white paper or at a window light; it will be reddish-brown whilst hot, but the color will become lighter on cooling; now heat the bead again for some time in the tip of the inner blowpipe flame, its color will be changed to green. These two colors, shown by the borax bead in the outer and inner blowpipe flames respectively, characterize the metal iron.

The borax bead is best detached from the wire whilst it is melted by giving the wire a sudden jerk; it may also be removed after cooling by crushing it on a hard flat surface by a sharp blow with the pestle. The wire when not in use should be kept in dilute hydrochloric acid. Should it not be perfectly clean after washing off the acid, the simplest method of cleansing it is to make a fresh bead upon the loop and jerk it off whilst melted, this will usually remove all impurity. A borax bead must never be used for these purposes until it has been found to be perfectly colorless, both hot and cold, when heated in the outer blowpipe flame. If the first bead

which is made in the loop is tinted with any color, it must be jerked off whilst fused and another bead formed with fresh borax, this being also thrown off and replaced if not quite free from color; this process is to be repeated until the bead is obtained perfectly clear and colorless.

**32. Flame Colorations.\***—Many substances give a characteristic color to the flame. The blowpipe flame is often employed to obtain these colors; an ordinary Bunsen flame, however, although it does not give the colors quite so promptly and intensely, answers the purpose well.

The flame best suited for this purpose is obtained by surrounding the Bunsen flame with a conical chimney, reaching about two-thirds of the height of the flame (Fig. 30), and then so regulating the supply of air entering into the air-holes that a small luminous point appears near the top of the flame; the platinum wire is held either in or just above this luminous tip. A filter-dryer also forms a very good chimney, being supported just above the foot of the burner on a wire tripod, which may be made by bending down the ends of a pipeclay triangle.

FIG. 30.



**EXP. 49.**—Hold a loop of platinum-wire in the inner blowpipe flame; if the wire is clean it will become red-hot without coloring the flame. If it colors the flame, clean the wire by boiling it with hydrochloric acid in a test-tube and then rinsing well with water; or by moistening with strong hydrochloric acid, and heating strongly in the blowpipe flame until it no longer colors the flame. If it cannot be thus cleansed, cut off the end. Now moisten the wire and dip it into a little powdered sodium chloride, or dip the loop into some sodium chloride dissolved in water so that when removed a drop of the solution remains in the loop; then hold the loop at the tip

\* For a more detailed and exact description of flame-tests, see Bunsen's paper in the *Philosophical Magazine* for 1867.

of the inner blowpipe flame, the outer flame will be colored bright yellow. Repeat the experiment, placing the moistened loop in the upper and outer part of the Bunsen flame, the same yellow tint will be obtained. This yellow color is given only by sodium and its compounds.

Remove all sodium chloride from the platinum-wire loop, by heating it strongly in the tip of the blowpipe flame until it ceases to impart any color to the flame. Then dip the wire into some potassium chloride solution; this will give to the flame a pale violet color, which is characteristic of potassium and its compounds.

As will be seen subsequently, the examination of flame colorations, aided by the use of colored glasses, and of a hollow glass prism filled with indigo solution, is of great use in ordinary analysis.

EXP. 49 *a.*—Look at the yellow flame coloration produced by sodium chloride, through a piece of cobalt glass, or through the indigo prism, no yellow color will be visible. Now dip the wire-loop into a mixture of a few drops of sodium chloride and potassium chloride solutions; when held in the flame only the sodium coloration is seen, but through the blue glass or prism the potassium is seen to produce a red flame coloration. By means of the blue glass or prism then potassium is discoverable in the presence of sodium, by which its presence is otherwise masked.

By far the most perfect method of examining flame colorations, more especially for the rarer elements or for mere traces of the more common ones, is by the use of the spectroscope; every advanced student should make himself familiar with the use of this instrument, at least in its simplest and smallest form. A paper on the use of the spectroscope for qualitative analysis will be found in the appendix; this is well worthy of the attention of senior students.

It is well to keep two platinum wires, one to be used for borax beads and the other for flame colorations, since a loop which has been used for borax colors the flame yellow.

**33. *Ignition on Charcoal in the Blowpipe Flame.***—Common wood charcoal forms a very useful support for

substances which are to be heated in the blowpipe flame. It cannot itself be melted, it conducts heat very badly, and possesses other properties which render it peculiarly adapted for ignitions. It is, if necessary, sawn into small pieces of the required shape and size, with a fine-toothed saw.

Ignition on charcoal serves chiefly to detect the presence of metals; the substance for this purpose is heated either alone or mixed with certain other substances (usually sodium carbonate and potassium cyanide) in the inner blowpipe flame; globules or scales of the metal are thus frequently produced, also in many cases the oxide of the metal forms a characteristic coating or "incrustation" on the charcoal; an examination of the color and properties of the metal, or incrustation, or of both, will frequently lead to the detection, or confirm the presence, of some metal. Charcoal is apt to leave a white ash when heated alone in the blowpipe flame; this must not be mistaken for an incrustation. A powder frequently requires to be moistened with water before being heated by the blowpipe flame, in order to prevent the flame from blowing it off the charcoal.

EXP. 50.—Choose a piece of wood-charcoal free from large cracks, scoop out a small hollow near one end with a knife, and in the hollow place a small quantity of a mixture of powdered lead acetate and sodium carbonate (or potassium-oxalate). Heat this mixture in the inner blowpipe flame, holding the charcoal in such a way that the rest of the flame plays over its surface (Fig. 31). After continuing the heating for some time, bright globules of lead will be seen in the cavity, and on removing the charcoal from the flame, the surface round the cavity will be found to be colored with a yellow "incrustation." If one of the little globules of metal





be then taken off the charcoal with the point of a penknife, and be struck smartly with a pestle on the bottom of a mortar turned upside down, it will flatten out into a cake, showing that the metal is "malleable," not "brittle." The metal lead is the only metal which gives a yellow incrustation and malleable globules; this test therefore serves to detect its presence.

**33 a.** The melted mass left on the charcoal after fusing a metallic salt with sodium carbonate is best examined

FIG. 32.



for metallic powder or scales by detaching it from the charcoal with the point of a penknife, then crushing it in an agate mortar or in a watch-glass with a little water; on stirring and quickly decanting the water down a wet glass rod or pestle held against the edge of the vessel (Fig. 32), the metallic powder, if present, will be left; by repeating this process of "levigation," the metal is obtained free from soluble substances and charcoal; it may then be further examined by a lens or magnet. A mixture of copper sulphate with sodium carbonate, when heated on charcoal and "levigated" as directed above, will furnish little red scales of copper; a mixture of ferrous-sulphate and sodium carbonate will yield a gray powder consisting of metallic iron, which, when touched with the pole of a magnet, will become attached to it.

**34. Use of Test-papers.**—Certain vegetable colors undergo remarkable changes when acted upon by chemical substances. Some of these are used for analytical purposes; the following experiment will explain how they are made use of.

EXP. 51.—Boil some pieces of a common red pickling cabbage-leaf in water; a blue liquid will be produced. Pour some of this liquid into a test-tube and add several drops of hydrochloric acid, the color will change to bright red; to another portion of the blue liquid add several drops of ammonia, the color changes to green; to a third portion of the blue liquid add a little ammonium chloride solution, the color remains unchanged.

Mix in a test-tube a little ammonia solution with three times as much distilled water, and add this weak ammonia gradually to the liquid reddened by several drops of hydrochloric acid; by careful addition the blue color may be exactly restored, the acid previously added being “neutralized” by the ammonia.

EXP. 52.—Pour some blue litmus solution (made by boiling solid litmus in water and filtering) into three test-tubes; to the first add hydrochloric acid drop by drop until it turns distinctly red, preserve this liquid; to the second add ammonia solution, the liquid remains blue; to the third add ammonium chloride solution, the color will not be affected.

Divide the reddened liquid into three parts; to one part add ammonia solution, the original blue color is restored; to another part add hydrochloric acid, the red color is unchanged; to the other part add ammonium chloride, the red color remains unaltered.

These experiments may also be made with litmus test-papers, which are prepared by soaking paper in the blue and red liquids respectively and allowing it to dry.

EXP. 53.—Repeat the preceding experiment, using the litmus-papers instead of the solution, and moistening them with the liquid by dipping a glass rod into it, and then touching the paper with the moistened end.

It will be seen then that the three substances employed differ in their action on litmus test-papers.

Hydrochloric acid . .	reddens	blue litmus	does not affect red litmus.
Ammonia . . . . .	does not affect	“ “	blues “ “
Ammonium chloride .	“ “ “ “	“ “	does not affect “ “

Each of these substances is a sample of a whole class, the members of which resemble it in their action on litmus.

The class of *acid substances*, including sulphuric acid, nitric acid, etc., and many metallic salts, resemble hydrochloric acid in reddening blue litmus and not affecting red.

The class of *alkaline substances*, including potassium hydrate, sodium hydrate, etc., and many salts of the alkalies, resemble ammonia in bluing red litmus, and not affecting blue.

The class of *neutral substances*, including potassium chloride, sodium chloride, etc., resemble ammonium chloride in not affecting either blue or red litmus.

Hence in analysis we use red and blue litmus-papers to test whether a liquid is "acid," "alkaline," or "neutral."

The red cabbage liquid, as will be seen, is better suited for a test-paper than litmus, since one paper shows both alkalies and acids; but the color is very apt to spoil with keeping.

Sometimes a liquid in which a colored precipitate is suspended has to be tested by litmus-paper, the precipitate then hides the color of the paper when a drop of the liquid is placed upon it. In such a case moisten the paper with a drop of liquid, and wash the liquid off after a short time with a stream of distilled water from the wash-bottle; the color of the paper is then distinctly seen.

**35.** An acid substance by being mixed with a proper proportion of an alkaline substance becomes "neutral;" similarly an alkaline substance is rendered "neutral" by mixture with a proper quantity of an acid. Thus by mixing hydrochloric acid and ammonia in the proper proportions, the neutral substance ammonium chloride is formed; the acid is then said to be "neutralized" by the alkali, or the alkali by the acid.

It is often required to "neutralize" an acid with an alkali or *vice versâ*; the following experiment will show how this process of neutralizing an acid is performed.

EXP. 54.—Pour a little hydrochloric acid into a porcelain dish, add to it ammonia, several drops at a time, keeping the liquid constantly stirred with a glass rod, and occasionally touching a slip of blue litmus-paper with the end of the rod moistened by the well-stirred liquid; as soon as the paper is only faintly reddened by the liquid the ammonia is added very carefully, stirring well after the addition of each drop, and trying the action of the liquid by touching the wet glass rod against first blue and then red litmus-paper, waiting a short time to give the liquid time to act on the paper. When the colors of both blue and red paper remain unaffected, the process of “neutralization” is completed.

It is well to add *dilute* ammonia towards the end of the process; this is prepared by mixing a little ammonia with three or four times as much distilled water in a test-tube.

The process of neutralizing may often be simplified when the liquid is clear and colorless, by dropping into it a small piece of blue litmus-paper, and after each addition of ammonia noting its color; the condition of the liquid should, however, even in this case be proved towards the end by moistening both red and blue paper.

The process of neutralizing an alkali with an acid is precisely similar to that of neutralizing an acid with an alkali, save that red litmus is at first used instead of blue.

35 a. It will be readily understood from what has been said above, that if an alkaline or neutral liquid is to be “acidified,” acid is added gradually as was described in the preceding experiment, until a drop of the well-mixed liquid reddens blue litmus. If an acid or neutral liquid has to be made alkaline, the alkali is added until a drop of the liquid turns red litmus blue. These processes are often termed adding acid or alkali “in excess.”

35 b. *Yellow turmeric paper* is occasionally employed instead of red litmus to test for alkalies, which turn it reddish-brown; it is unaffected by acids, with the exception of boracic acid. The most important use for tur-

meric paper is to test for boracic acid, which, if dried on the paper at a gentle heat, turns it orange-red.

EXP. 55.—This may be shown by stirring a little borax upon a watch-glass with dilute hydrochloric acid until it is dissolved, moistening the lower part of a slip of turmeric paper with this liquid, and drying it by holding it at some distance above the flame, or better by placing it in a steam-oven; the portion of the slip which was moistened will now appear reddish-brown, the color being well seen by contrast with the light-yellow upper portion of the paper.

### FRACTIONAL SOLUTION, CRYSTALLIZATION, AND DISTILLATION.

1. *Fractional Solution*.—Since different substances show differences in solubility in water and acids, it is often possible by means of solvents to separate the constituents of a mixture. This process is termed separation by “fractional solution.” Illustrate this by making a mixture of powdered ferrous sulphate ( $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ ) and ferric oxide ( $\text{Fe}_2\text{O}_3$ ); on warming a portion of this with water,  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$  alone dissolves, leaving  $\text{Fe}_2\text{O}_3$ ; if another portion is warmed with  $\text{HCl}$ , both substances dissolve.

2. *Fractional Crystallization*.—If a solution of two different salts be boiled down in an evaporating basin, one salt will usually furnish crystals before the other does, and by this process of “fractional crystallization” a separation of the salts may often be effected. Boil down a solution of sodium chloride ( $\text{NaCl}$ ) and nitre ( $\text{KNO}_3$ ):  $\text{NaCl}$  will crystallize out of the boiling liquid in hollow cubes, and on cooling prisms of nitre will separate.

3. *Fractional Distillation*.—Liquids which boil at different temperatures may often be separated by distillation (p. 36); that part of the mixture which first distils over consisting chiefly of the liquid with the lowest boiling-point, and the successive portions of the distillate be-

coming richer in the liquids of higher boiling-points. The process of "fractional distillation" may be illustrated by distilling a mixture of alcohol and water ; since alcohol boils at a lower temperature than water, the first part of the distillate contains the greater part of the alcohol, as is shown by its burning and by its dissolving camphor or iodine in quantity. The last portion of the distillate does not give these tests for alcohol, and is proved to be water by turning dry  $\text{CuSO}_4$  blue and dissolving it.

## SECTION IV.

## ANALYTICAL REACTIONS.

*Course of Practical Analysis to be pursued by the Student.*—The text is so arranged that two main courses may be framed from it in this and the following sections. The one is suited to senior students and dispenses with the training in the detection of simple substances, which, although useful as a stepping-stone to the full analytical method, is in itself of only artificial importance. The other course is suited to junior students, and forms an easy approach to the more difficult complete methods of analysis: it teaches the plan of analyzing simple substances which may contain one acid and one metal. This course is therefore also suited to those who are learning analytical chemistry with the view of preparing themselves for examinations in which the candidate is not required to detect more than one acid and one metal. These courses are described below as the "Senior Course" and "Junior Course."

*The Senior Course* consists in working through the analytical reaction in Section IV; and as those for each group are completed, several substances whose composition is unknown to the students are tested for one member only by the Table of Differences at the end of the group; as soon as these are detected with readiness, substances which may contain two or more of the members mixed together are analyzed by the group table in Section VI: a reference to the Table of Differences will show at a glance on what principles these group tables are founded, and this is further explained by a statement following the Table of Differences. This plan of working with each group table before going on to the reactions for the next group is to be recommended, since,

whilst the differences upon which the separation and special tests of the elements in the Groups are fresh in the student's memory, they are practically applied. The Analytical Tables are thus gradually rendered intelligible one after another, whilst the student is proceeding through the reactions. The book is so arranged, however, that all the Analytical Tables may be left until the reactions for all the groups have been finished. When all the groups have been worked through in this manner, the analysis of substances is commenced which may contain any members of any of the different analytical groups; the directions for the systematic method of analysis required for this purpose being contained in Section VI.

*The Junior Course.*—Since this is intended to teach only the detection of one metal and one acid, the student whilst working through Section IV omits all those paragraphs marked "s" after the number, relating to the separation and detection of the mixed members of a group; and after finishing the reactions for each group, he proceeds to detect the members occurring singly in several substances of unknown composition by the Table of Differences and by reference to the reactions. After working in this way through Section IV, he at once proceeds to analyze simple substances by Section V.

As stated above, this Junior Course may occasionally be made to precede the Senior with advantage, that is to say, the student after working through the Junior Course, and learning how to detect simple substances, may analyze mixtures containing two or more members of each group by the directions given at the end of each of the groups in Section IV, and may then proceed to the full method of analysis contained in Section VI.



## INTRODUCTORY REMARKS ON ANALYTICAL CHEMISTRY.

The science of Chemistry reveals to us the fact that every substance on this earth, and probably in the universe, consists either of one kind of matter only, or of two or more different kinds, which we cannot further separate. Those substances which contain only one kind of matter, are called "chemical elements;" their number at present amounts to a little over sixty. A list of them is given in paragraph 558 at the end of the book.

It is not often that these elements occur singly: two or more of them are commonly found together either in a state of mere "mixture," or united in a much more intimate manner by the force of "chemical affinity" to form a "chemical compound."

Analytic chemistry or analysis (literally meaning an *unloosing* or *separation*), teaches us how to discover what element, elements, or sometimes what groups of elements, any substance of unknown composition contains: it also enables us to ascertain whether any particular element or group of elements is present in a given substance. In order to become a thorough analyst, it is necessary to be intimately acquainted with the properties of every element and of the compounds it forms with other elements, and therefore efficiency in analysis depends in a large measure upon the knowledge of what is usually termed "Theoretical," or more appropriately perhaps "Descriptive, Chemistry." But although we recognize a substance by its properties, still a knowledge of *all* its properties is not requisite to enable us to detect its presence: for ordinary analytical purposes certain marked properties of each element or group of elements are selected, and by these properties, called "tests" or "reactions," we detect its presence.

The "tests" or "reactions" most frequently selected are the following: 1st. The behavior of a substance when heated under various conditions; and 2d. Its be-

havior when mixed with certain liquid or solid substances. Such substances, added for analytical purposes, are called *reagents*. Some reagents serve for detecting one element or compound only, and are therefore called *special reagents*; others separate or detect a group of elements or compounds, and are called *group reagents* or *general reagents*: these latter reagents serve to separate substances into "Analytical Groups," as will be more fully explained hereafter.

The student must bear in mind that, although an acquaintance with the most important reaction of substances is indispensable to success as an analyst, caution, skill, and neatness in manipulation are not less necessary, and the most scrupulous attention must be paid to all directions bearing on these matters.

**36. Analytical Classification.**—The chemical elements are frequently divided into two classes, viz., *metals* and *non-metals*. These classes form, with a slight modification, convenient divisions for analytical purposes. In the class of "metals" must be included the hypothetical metallic radicle ammonium ( $\text{NH}_4$ ), also the element hydrogen. The class of "non-metals," either singly or in groups formed by their combination *inter se* or less frequently with a metal, unite with hydrogen and with metals forming "salts;" the hydrogen salts of these radicles being often called "acids." Thus  $\text{Cl}^*$ ,  $\text{SO}_4$ ,  $\text{PO}_4$ ,  $\text{MnO}_4$  form the hydrogen salts (acids),  $\text{HCl}$ ,  $\text{H}_2\text{SO}_4$ ,  $\text{H}_3\text{PO}_4$ ,  $\text{HMnO}_4$ , and the metallic salts,  $\text{KCl}$ ,  $\text{MgSO}_4$ ,  $\text{Na}_3\text{PO}_4$ ,  $\text{KMnO}_4$ ; metal hydrogen salts (acid salts), such as  $\text{NaHSO}_4$ ,  $\text{Na}_2\text{HPO}_4$ , are also known. The general name "acid radicle" is conveniently applied to  $\text{Cl}$ ,  $\text{SO}_4$ ,  $\text{PO}_4$ ,  $\text{MnO}_4$ , etc., it being understood that although most of these are unisolated compound radicles, *i. e.*, contain more than one element—some of them are elements; their common property is that of forming in combination with hydrogen and with metals—salts. For analytical

---

\* The meaning and use of chemical symbols will briefly be explained hereafter (40); but the student is presumed to be already acquainted with them. In p. 540 will be found a list of the symbols of the different elements.

purposes, then, we may separate the substances we have to detect into the groups of metals and acid radicles. In trying the reactions for these substances, it is usual to employ, not the substances themselves, but certain compounds—usually salts—which contain them.

For convenience of hasty writing, it is not uncommon to employ abbreviations for the compound radicles, especially for such as are constantly met with or are of unusual complexity. In the text the only such abbreviations employed are:

Am for  $(\text{NH}_4)$ , the metallic radicle ammonium.

T “  $(\text{C}_4\text{H}_4\text{O}_6)$ , the acid radicle of tartaric acid and the tartrates.

A “  $(\text{C}_2\text{H}_3\text{O}_2)$ , the acid radicle of acetic acid and the acetates.

Cfy “  $(\text{FeCy}_6)$ , the acid radicle of ferro- and ferricyanides.

Ho “  $(\text{HO})$ , the radicle Hydroxyl, occurring in hydrates, etc.

**37. Analytical Groups.**—The metals are separated by their behavior with certain general reagents into five principal groups, two of these being further subdivided into two sub-groups. Each of these groups receives a distinctive name—either a number from its place in the systematic course, or a name from some conspicuous or important member of the group, or from the general reagent used to precipitate the group. Thus we speak indifferently of the 4th Group, the Barium Group, or the Ammonium-carbonate Group. In sec. v, par. 330 will be found a table showing the analytical groups with their distinctive numbers, names, and group reagents placed at the head of each column; and the names and symbols of the elements contained in the group arranged vertically beneath.

It is perhaps well to avoid referring to a group by its number, as different analysts number the groups differently, and the name thus given is therefore apt not to denote the group with certainty.

**38. Method of Trying the Analytical Reactions in Section IV.**—In order to become fully acquainted with the behavior of different substances with “reagents,” and thus to be able to detect them by their “reactions,” each of these reactions should be performed with the substance itself or with one of its compounds, and the appearances presented carefully observed, so that the substance would be easily recognized in a body of unknown composition by means of those tests.

The following general rules must be carefully attended to in trying the reactions; they may be read through before commencing the reactions par. 47:

1. A *solution* of the substance which is directed to be used must always be employed, unless it is specially stated that the *solid* is required. Solid substances are required as a rule only for blowpipe reactions. A solution can be readily made from the solid, if necessary, by warming it with water, or in some cases with dilute acid (529). This is often called the “original solution.”

2. Commence by taking in a test-tube or small beaker a small quantity (about 15–20 c.c.) of the solution of the substance. If the solid substance is also required, take some of it upon a watch-glass.

3. Pour a small portion of the solution into a perfectly clean test-tube, and add the first-mentioned reagent to it. The reagent must always be added in small quantity at first; more can be employed afterwards if excess is required. The student must from the first acquire the habit of working with small quantities both of solutions and reagents.

For each of the reactions a separate portion of the “original solution” is thus poured into a clean test-tube and the required reagent added to it.

4. Before trying each reaction the student must carefully read through the description of the results he is to obtain; he must then perform the experiment literally following out the directions given. He must consider it a necessary condition of after success that each result is obtained precisely as stated in the text, and must never on any account pass on until he has conscientiously sat-

isfied himself that the statements of the book are true, and that he could at any time repeat the test successfully.

39. *Entry in Note-book.*—Each reaction, as soon as it has been satisfactorily tried, should be neatly entered in the note-book in a short form. Since nothing so much aids brevity as the symbolic notation commonly used by chemists, the student should invariably adopt this chemical shorthand in entering results. A full account and explanation of chemical notation may be found in any work on theoretical chemistry, and merely a few hints will therefore be here given which will be of special use for the purpose of briefly entering reactions.

40. *Chemical Notation.*—Each element has its corresponding *symbol* consisting of one or two letters; thus S stands for sulphur, Cl for chlorine. These symbols also stand for one *atom* of each element; by being written one after another they give the *formula* of a compound substance, showing what elements, and how many atoms of each of these elements, its molecule contains; thus KCl stands for one molecule of potassium chloride, a compound of one atom of potassium and one atom of chlorine;  $\text{PtCl}_4$  stand for one molecule of platinum chloride, which consists of one atom of platinum combined with four atoms of chlorine, the small figure placed *below* a symbol to the right denoting how many atoms of that element are present. A number placed before a formula multiplies each symbol in that formula; thus  $2\text{PtCl}_4$  signifies two molecules of platinum chloride, containing together two atoms of platinum and eight atoms of chlorine. If two or more symbols are included in brackets, any number placed outside, either *before* or *below* the brackets multiplies each symbol contained in the brackets; thus both  $\text{Sr}(\text{NO}_3)_2$  and  $\text{Sr}_2(\text{NO}_3)_4$  stand for  $\text{Sr}_2\text{N}_2\text{O}_{12}$ , and this may also be written  $\text{Sr}_2\text{NO}_6$ . In paragraph 540 will be found a list of the elements, with their symbols and atomic weights; paragraphs 524, 525, 526, 529, and 530 contain a list of the names and corresponding formulæ of all the most common compound substances mentioned in the course of the book. Symbols and formulæ are always used in the text instead of

names, but the student can readily find for these corresponding names by reference to the above tables; or to the labels on the bottles, which should contain both names and formulæ.

41. On adding a "reagent" to a substance some change usually occurs which gives rise to an alteration of color, or very frequently to the formation of a precipitate having a certain color, appearance, or behavior with other liquids, which are characteristic of that particular substance, or of some element or group of elements contained in it; this change should be briefly described in the note-book in words, and then represented by a *chemical equation*. The rules for drawing out an equation may be found in any treatise on chemistry; a few general directions only are here given, which will be of service for the special object in question.

42. *Rules for writing down an Equation.*—Write down the formulæ of the two substances which are mixed together for the reaction, with the sign of addition (+) between them; then write the sign of equality (=), followed by the formula of the precipitate produced. In a completed equation the formulæ written on the left-hand side of the sign (=) are usually called the "left-hand side" of the equation; those to the right the "right-hand side." Since most of the following reactions are cases of "double decomposition"—i. e., cases where an exchange of certain elements or groups of elements occurs between the two compounds,—a little consideration will usually show how many molecules of the substances on the left-hand side of the equation are required to yield the formula for the precipitate, and also whether any other, and if so what other, substance is at the same time formed.

It is always necessary that the number of atoms of any one element on both sides of the equation should be equal. The following are dissected examples:

The first test for K (47):



This is simply placing together the substances mixed

and the precipitate formed ; the equation thus formed is evidently not correct, since we have two atoms of K on the right-hand side and only one on the left, also 6 atoms of Cl on the right, and only 5 on the left ; but this inequality is at once removed if we place 2 before the KCl, thus :




Evidently if the above equation is correct the precipitate is here the only substance formed, since the number of atoms in the substances on the left-hand side is exactly equal to that in the substance on the right.

Take now the first test for Ba (69) :



This represents the substances mixed and the precipitate formed, but evidently the left-hand side of the equation contains more than the right (viz., Am<sub>2</sub> and Cl<sub>2</sub>) ; hence we write it fully thus :



which states that on mixing together barium chloride (BaCl<sub>2</sub>) and ammonium carbonate (Am<sub>2</sub>CO<sub>3</sub>), we obtain barium carbonate (BaCO<sub>3</sub>) and ammonium chloride (AmCl).

In writing out these equations a systematic method should be observed. Write down on the left-hand side of the equation first the substance whose reaction is being tried, then the reagent added ; and on the right-hand side write first the formula for the precipitate with a line drawn over it by way of distinction, then any substance or substances which may be formed with it.

In the reactions which follow, the formulæ of the substance used, of the reagent added, and of the precipitate produced will always be given, these being sufficient data to enable the student to form the equation ; only where the equation is exceptionally difficult or complicated will it be given in full.

43. The following contractions are recommended :

<i>expt.</i>	for	experiment.	<i>insol.</i>	for	insoluble.
<i>pp.</i>	"	precipitate.	<i>soln.</i>	"	solution.
<i>ppn.</i>	"	precipitation.	<i>crystn.</i>	"	crystalline.
<i>ppd.</i>	"	precipitated.	<i>coln.</i>	"	coloration.
<i>sol.</i>	"	soluble.			

44. Examples are given below showing how these reactions should be entered in the note-book ; they should be referred to as soon as the student commences the reactions for the metals :


1st. [See pars. 47-50.]

POTASSIUM (K).—Used KCl.

$PtCl_4$ : stirred on watch-glass: yellow cryst<sup>a</sup> pp, sol. in much water and in KHO, insol. in alcohol:



$NaHT$ : well shaken in a test-tube: white cryst<sup>a</sup> pp., sol. in water, KHO, and HCl:




*Flame col<sup>n</sup>*: pale violet, violet-red through the indigo-prism or cobalt glass.

*Heated in tube or on platinum foil* (solid KCl): melted, but gave no fumes; hence not volatile.

2d. [See pars. 69-74.]

BARIUM (Ba).—Used  $BaCl_2$ .


$Am_2CO_3$ : white pp., sol. in  $H\bar{A}$ :



$CaSO_4$ : heavy white pp., insol. in boiling HCl:



$K_2CrO_4$ : light yellow pp., sol in HCl, insol. in  $H\bar{A}$ :



*Flame col<sup>n</sup>*: on loop of plat<sup>a</sup> wire, yellowish-green color.

45. *General Rules to be observed whilst working.*—The student should attend carefully to the following precautions, which will most materially aid the progress and accuracy of his work ; if they are observed from the commencement they will not be felt irksome:

1. Before commencing work look through the reagent



bottles belonging to the working bench; replace any which are absent, and arrange the bottles, if necessary, in the order indicated on an accessible list, or shown by the number on the labels of the bottles, or by the labels on the edge of the shelf. Then proceed to fill any which are empty and to relabel any whose labels are imperfect or loose. If any of the liquid reagents are not quite clear when shaken they must be filtered before use.

2. Arrange the apparatus required for use, on the hinder part of the bench, so as to leave the front free to work upon; this will lessen the risk of upsetting anything.

3. Before commencing work, all glass and porcelain which is not in use should be carefully cleansed, if not already clean; but it is best never to put apparatus away dirty.

4. In using a reagent bottle take it in the right hand, remove the stopper by taking hold of it between any two of the fingers, or between the fourth finger and palm, of the left hand; hold the stopper in this way and replace it immediately after the bottle has been used. Pour the liquid out at the side opposite to that bearing the label, and prevent the last drop from running down the outside by touching the lip of the bottle either with the stopper or against the clean edge of the vessel. Solid reagents should be taken out of the bottle on a clean knifeblade or spatula, or with a spoon of wood or horn; or a small quantity should be turned out upon a piece of paper or upon the palm of the hand. *The bottle must be restoppered and replaced on the shelf in its proper place with the label outwards immediately after use, and must never be left standing on the working bench.*

5. If any solution or precipitate has to be put by for some time, always label it with a piece of gummed paper bearing an inscription or a reference number which is explained in the note-book, and is sufficient to recall to mind what the solution or precipitate consists of; never trust to memory in this matter.

6. Brass crucible-tongs must never be employed for holding hot evaporating basins containing acids, as some

of the brass is apt to be dissolved and thus introduced into the solution. A hot dish may be moved either by holding it with a cloth or by taking hold of the stand upon which the dish is placed.

7. *Liquids only* are to be poured down the sinks; all *solid* refuse, such as soiled filters, broken glass, pieces of charcoal, etc., which would tend to stop the holes in the sink, must be thrown into a box placed in any accessible position in the laboratory, or into a small earthenware jar or vessel placed upon each working bench.

8. When a student is examining a substance to detect its composition, he must fully write down in his note-book each test or process, with the result obtained, as soon as it is completed. The analysis is thus written out gradually as it progresses; *on no account must the entry of the results be left until the completion of the analysis.* This rule cannot be too strongly enforced, as neglect of it is always attended with loss to the beginner. The most convenient form of entry is that of the Analytical Tables.

\* \* Cleanliness and neatness in analytical work cannot be too carefully attended to; more confusion and error may be caused by using dirty test-tubes, beakers, dishes, and funnels than would be thought possible by the beginner.

---

### REACTIONS FOR THE METALS.

*Note.*—In trying through the reactions for the metals, the usual analytical order of the groups has been reversed, because the most simple reactions occur in the last groups, and they are therefore better suited for a beginner.

#### GROUP V.—POTASSIUM GROUP.

46. This group includes K,  $\text{NH}_4$ ,\* Na, Mg. † The first three metals (K,  $\text{NH}_4$ , Na) are known as the “alkali metals.”

The members of this group are not precipitated by any of the five group reagents.

---

\* A hypothetical metal radical, “Ammonium.”

POTASSIUM (K).—Use KCl.


Refer if necessary to paragraphs 38 and 44 for directions how to work.

47.  $PtCl_4$  added after a drop of dilute HCl to some of the KCl solution on a watch-glass and stirred with a glass rod, gives a yellow crystalline precipitate ( $K_2PtCl_6$ ), forming first along the lines where the rod has rubbed the glass.

Stir up the precipitate and pour off the liquid with the precipitate into three test-tubes; to one add much distilled water and heat, the precipitate dissolves; to the other part add alcohol, the precipitate does not dissolve. Hence this precipitate is soluble in water, insoluble in alcohol. By warming the other portion with solution of KHO the precipitate is also shown to be soluble in that liquid.

*Remarks.*—The following remarks are of great importance in using the above test for K. Since this precipitate is soluble in alkalis the liquid should always be neutral or only faintly acid before adding  $PtCl_4$ , and since it is more insoluble in alcohol than in water, addition of alcohol causes the precipitate to form more rapidly in dilute solutions; owing to the solubility of the precipitate in water, very diluted solutions must, however, be evaporated nearly to dryness before applying the test. It is always well to add several drops of HCl before the  $PtCl_4$ , and to make sure by test-paper that the liquid is acid in reaction. These remarks apply also to the precipitate produced by  $PtCl_4$  with ammonium salts (52).

48.  $H_2\bar{T}$  (or better  $NaH\bar{T}$ ) mixed with some of the solution in a test-tube gives a white crystalline precipitate ( $KH\bar{T}$ ); usually this precipitate appears only when the mouth of the test-tube is closed with the thumb and the tube well shaken for some time.



Shake up and pour some of the liquid and precipitate

into four test-tubes ; add to these portions water,\*  $\text{KHO}$ , and  $\text{HCl}$  respectively ; on being shaken and warmed the precipitate will dissolve in each of these liquids. Hence this precipitate is soluble in  $\text{H}_2\text{O}$ ,  $\text{KHO}$ , and  $\text{HCl}$ . To the fourth portion add alcohol ; the precipitate does not dissolve, since it is insoluble in alcohol.

*Remarks.*—In using the above test for the detection of  $\text{K}$ , the precipitate is not obtained in dilute solutions at once, but its formation is much hastened by the addition of alcohol ; very dilute solutions must first be concentrated by evaporation. The solution to be tested should be neutral or only feebly alkaline ; if acid in reaction, the acid should be boiled off (if volatile) or neutralized by  $\text{Na}_2\text{CO}_3$  ; feebly acid solutions may, however, be at once mixed with  $\text{NaHT}$  ; to an alkaline solution  $\text{H}_2\text{T}$  should be added till the reaction is strongly acid.

*Note.*—In the following reactions when a precipitate is stated to be soluble or insoluble in certain liquids, it must be proved to be so in the above manner unless different directions are given.

**49. Flame Coloration (32).**—Hold a loop of platinum wire in the flame, and see that it gives no color to the flame ; then dip it into some  $\text{KCl}$  solution, or better still, moisten it and dip it into some powdered  $\text{KCl}$ , and again hold the loop in the upper part of the flame. A *pale violet* color will be imparted to the flame if the  $\text{KCl}$  is pure ;† but whatever color is thus seen, the flame-color will always appear *pale blue* or *violet* through the thinner portions of an indigo-prism, and deep *purple-red* through the thickest parts.

The light of a Bunsen flame itself usually appears pale blue through the indigo-prism ; this color should be noted before using the prism for examining a flame coloration.

The flame coloration due to potassium shows in the spectroscope a red line ( $\alpha$ ) and an indigo-blue line ( $\beta$ ).

*Note.*—A piece of blue cobalt glass may be used instead of the

\* By "water" is always meant "distilled water," unless otherwise stated.

†  $\text{KNO}_3$  usually gives a purer potassium coloration.

indigo-prism, but is by no means to be recommended, as the color of different specimens is somewhat variable in shade and in intensity, whereas the color of the indigo can be obtained of an invariable tint.


50. Heat a little solid KCl in a small, dry test-tube, or better on a piece of platinum foil; the KCl will melt, but will not pass into vapor producing white fumes unless heated very strongly, since K-salts are non-volatile at a moderate heat. The fused residue is often transparent and invisible if in small quantity; its presence is shown by a slight crackling being heard during cooling.

---

AMMONIUM ( $\text{NH}_4$ , or Am).—Use  $\text{NH}_4\text{Cl}$  (AmCl).

*Note.*—The symbol Am is often written for  $\text{NH}_4$ ; it is very convenient, as it does away with the necessity of using brackets; *e. g.*,  $\text{Am}_2\text{S} = (\text{NH}_4)_2\text{S}$ .

51.  $\text{KHO}^*$  (or  $\text{NaHO}$ ) poured into some of the solution, or upon the solid, in a test-tube and heated gives off  $\text{NH}_3$  gas:



This gas is known by the following properties:

- a. Its pungent smell—that of common smelling salts.
- b. By turning moist red litmus-paper blue. A small piece of red litmus-paper is wetted with distilled water and placed upon a clean glass rod, and is then held in the test-tube, taking great care not to touch with it the liquid or the sides of the tube; the paper becomes blue.
- c. A glass rod dipped into strong HCl or strong HA, and held over the mouth of the test-tube produces dense white fumes.

52.  $\text{PtCl}_4$  stirred on a watch-glass with some AmCl solution gives a yellow crystalline precipitate ( $\text{Am}_2\text{PtCl}_6$ ), soluble in water and in KHO, insoluble in alcohol. The

---

\* In all cases where KHO is to be used NaHO may be substituted, and either KHO or NaHO may be used unless it is specially stated that "pure NaHO" is to be employed.

remarks at the end of par. 47 apply here also. This precipitate if boiled with KHO evolves  $\text{NH}_3$  (difference from  $\text{K}_2\text{PtCl}_6$ ).

53.  $\text{H}_2\bar{\text{T}}$  (or better  $\text{NaH}\bar{\text{T}}$ ) shaken for some time in a test-tube with AmCl solution gives a white crystalline precipitate ( $\text{AmH}\bar{\text{T}}$ ), soluble in  $\text{H}_2\text{O}$ , KHO, and HCl, insoluble in alcohol. The remarks in par. 48 apply here also.

*Note.*—This precipitate is more soluble than the  $\text{KHT}$  precipitate, and therefore does not form so readily. Indeed with  $\text{H}_2\bar{\text{T}}$  a precipitate rarely forms unless alcohol be added, in which the precipitate is less soluble than in water. With  $\text{NaH}\bar{\text{T}}$  a precipitate is obtained on shaking the liquid well for some time unless it is very dilute.

54. *Flame Coloration.*—AmCl, if pure, gives on platinum wire no color to the flame.

55. *Heat a little solid AmCl* in a small dry test-tube, or on a piece of platinum foil; it will be converted into vapor producing white fumes, which are seen (if platinum foil is used) by removing the foil for a moment from the flame. Hence AmCl is volatile.

If the experiment is performed in a dry test-tube, a white coating or “sublimate” forms on the upper part of the tube.

---

SODIUM (Na).—Use NaCl.

56. *Flame Coloration.*—NaCl gives an *intense yellow* color to the flame; the color is almost invisible or pale blue when viewed through the indigo-prism, *and never shows the slightest tinge of red and purple*, even when seen through the thickest parts of the prism. A crystal of red  $\text{K}_2\text{Cr}_2\text{O}_7$  appears colorless if illuminated by this yellow flame. Examined by a small spectroscope, the sodium coloration gives a single yellow line (a).

The yellow color of the sodium flame is always readily seen, since it overpowers that of K; its production is the most reliable, and often the only possible, test for sodium.


*Note.*—If both K and Na are present, the K coloration is unseen and the yellow coloration of Na is alone visible; but on examining this flame through the indigo prism or cobalt glass, the purple coloration of the K is at once seen; prove this by mixing together KCl and NaCl and trying the flame coloration without and with the indigo prism. The spectroscope also shows the lines of Na and K quite distinct from one another.

57. *Some solid NaCl heated in a dry test-tube, or better on platinum foil, melts without producing white fumes, unless heated very strongly, since NaCl is not volatile at a moderate heat.*

---

MAGNESIUM (Mg).—Use  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  solution.

58.  $\text{Na}_2\text{HPO}_4^*$  added after AmCl and AmHO gives a white crystalline precipitate ( $\text{MgAmPO}_4$ ); if much water is present this precipitate forms only when the liquid is warmed and well stirred or shaken. This precipitate is soluble in acids (use HCl or  $\text{HNO}_3$ ), insoluble in AmHO.



59. *KHO* gives a white precipitate ( $\text{MgHo}_2$ ),† soluble in acids.

60. *Flame Coloration.*— $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , if pure, gives no color to the flame.

61. *AmHO* gives a white precipitate ( $\text{MgHo}_2$ ); but if AmCl is added to the  $\text{MgSO}_4$  solution before the AmHO, no precipitate is produced.

62.  $\text{Am}_2\text{CO}_3$  gives a white precipitate ( $\text{MgCO}_3$ ), but only in strong solutions, and on standing for some time. AmCl prevents the formation of this precipitate.

---

\*  $\text{NaAmHPO}_4$  is better than  $\text{Na}_2\text{HPO}_4$ , since it produces the precipitate more readily than does  $\text{Na}_2\text{HPO}_4$  in dilute solutions, and always in a crystalline condition

† This convenient symbol (Ho) for the group (Hydroxyl), first proposed by Frankland, will be employed throughout in the formulæ for all hydrates containing more than one atom of Hydroxyl; it prevents the necessity of using brackets, as a number placed below the symbol multiplies both H and O, thus  $\text{Ho}_2$  equals  $(\text{HO})_2$  or  $\text{H}_2\text{O}_2$ ; and  $\text{Ho}_3$  equals  $(\text{HO})_3$  or  $\text{H}_3\text{O}_3$ .

63. *Blowpipe Test.*—A little solid  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , placed in a small cavity on a piece of wood charcoal and then heated in the tip of the outer blowpipe flame, shines brightly and gives a white unmelted mass of  $\text{MgO}$ ; no white fumes are given off, since  $\text{MgO}$  is not volatile. If this white mass after cooling is moistened with several drops of cobalt-nitrate solution and again strongly heated for some time\* in the outer blowpipe flame, it becomes *delicate pink* in color; this color is best seen when the mass is *quite cold*, by comparison with a white piece of paper held near the residue on the charcoal.

---

\* It is necessary to heat strongly after moistening with  $\text{Co}(\text{NO}_3)_2$ , else blue  $\text{Co}(\text{NO}_3)_2$  remains.



## 64. GROUP V.—TABLE OF DIFFERENCES.

*Note.*—A blank (thus —) signifies that no marked or characteristic change occurs. For the method of using the table see par. 65.

Tests.	K—Salts.	NH <sub>4</sub> —Salts.	Na—Salts.	Mg—Salts.
<i>*a. For Solutions.</i>				
1. Boil with KHO solution.	—	NH <sub>3</sub> gas given off: known by its smell and by turning moist red litmus blue; it also gives dense white fumes with strong HCl.	—	White precipitate.
2. Heat in flame on clean platinum wire.	<i>Pale violet flame, which appears pale violet or purple red, through the indigo-prism.</i>	—	<i>Intense yellow flame, which shows nothing of red through the indigo-prism.</i>	—
3. Add AmCl, AmHO, and Na <sub>2</sub> HPO <sub>4</sub> .	—	—	—	White crystalline precipitate.
4. Add PtCl <sub>4</sub> and a drop of dilute HCl, and stir vigorously.	Yellow crystalline precipitate, forms only in strong solutions.	Yellow crystalline precipitate, forms only in strong solutions.	—	—
5. Add H <sub>2</sub> T̄ (or better NaHT̄) and shake well.	White crystalline precipitate in strong solutions.	White crystalline precipitate in strong solutions.	—	—
<i>b. For Solids only.</i>				
6. Heat to redness in a dry test-tube, or on platinum foil.	Non-volatile, unless heated to bright redness.	Volatile; white fumes are given off and the substance passes into vapor (at least partially) forming a sublimate on the upper part of the tube.	Non-volatile, unless heated to bright redness.	Non-volatile.
7. Heat strongly on charcoal in the outer blowpipe flame.	Melt easily and sink into the charcoal, coloring the flame pale violet. (See 2.)	Are volatile, giving off white fumes.	Melt easily and sink into the charcoal, coloring the flame intense yellow. (See 2.)	Leave ultimately a white infusible residue which shines brightly, and which if moistened with Co (NO <sub>2</sub> ) <sub>3</sub> solution and reheated becomes pink.

\* In using these tests for the detection of a single member of the group in solution, only the first four need be employed; the fifth test may be substituted for the fourth. Tests 6 and 7 are used only for the examination of solids. They may be supplemented, however, by tests 1 and 2.

65. The method of using the foregoing table may be explained in a few words. Suppose a *solution* to be given which is known to contain one member of this group, the first five tests in the table (four and five being alternative) will enable us to decide which member is present. It is simply necessary to examine *separate portions* of the solution by these tests, until a result is obtained which conclusively proves the presence of one of the four substances. Tests 1, 2, 6, and 7 are similarly employed for examination of a *solid* substance. The student should never rest content with one test only when a second can be tried in confirmation, and he should always consider the tables of differences as mere summaries, and should refer back to the separate reactions for fuller accounts of the tests when required.

The results of these analyses, and of all analyses subsequently made by using the tables of differences, are best entered in three parallel columns, as shown in the upper part of the table in par. 67. The statement of the experiment or test performed is entered in the first column, the result observed being placed beside it in the second column, whilst beside this in the third column is stated the conclusion inferred from the result obtained.

At the end of each analysis state the result thus:

*Found K.*

---

66. (s.)\* *Detection of Members Mixed.*—We can also devise from the above table a plan for detecting the members of this group when mixed together. It is evident—

1. That  $\text{NH}_4$  can always be detected by boiling with KHO solution.

2. That Na can always be detected by its yellow flame coloration.

3. That Mg can always be detected by giving a

---

\* Paragraphs marked with an (s) are intended for a student using the senior course, and are passed over in the junior course.

white precipitate on addition of  $\text{AmCl}$ ,  $\text{AmHO}$ , and  $\text{Na}_2\text{HPO}_4$ .

4. That K may always be detected by giving a flame coloration which appears pale violet through the indigo-prism.

5. That the confirmatory test for K, by stirring with  $\text{PtCl}_4$ , may be tried if no  $\text{NH}_4$  is present; but it cannot be employed when  $\text{NH}_4$  has been found, since  $\text{NH}_4$  gives a yellow precipitate exactly like that given by K. Hence we first get rid of  $\text{NH}_4$  by evaporating some of the solution to dryness, and strongly heating the solid substance thus obtained on platinum foil as long as white fumes come off; the residue (if any) on the foil will then be free from  $\text{NH}_4$ , and can be dissolved and tested for K by  $\text{PtCl}_4$ .

The above remark concerning the  $\text{PtCl}_4$  test applies also to the  $\text{H}_2\text{T}$  test for K, since  $\text{NH}_4$  gives a precipitate with  $\text{H}_2\text{T}$  precisely similar to that yielded by K.

The following scheme, which embodies the above tests, must be used to test for Mg, K, Na, and  $\text{NH}_4$ , when they may be present together.

The student should examine several substances by this table for the four members of Group V, writing down the results he obtains fully in the form adopted in the table; that is to say, three columns are ruled; in the first is described the test performed, in the second the result observed, in the third what substance is inferred to be present or absent from that result.

**67. (s.) TABLE FOR TESTING FOR Mg, K, Na, NH<sub>4</sub>, IN  
A SOLUTION WHICH MAY CONTAIN ONE OR ALL OF  
THEM.**

(See note at the end of the Table.)

Experiment.	Observation.	Inference.
I. Add to a small part of the solution $\text{AmCl}$ , then $\text{AmHO}$ and $\text{Na}_2\text{HPO}_4$ , shake well and allow to stand if no precipitate appears at once.	1. A white precipitate forms.	Presence of Mg.
	[2. No precipitate appears.]	[Absence of Mg.]
II. Dip a small loop of clean platinum wire into the solution, and hold it near the top of the Bunsen flame.	1. A yellow flame.* Look at the yellow flame through the indigo-prism; if a reddish-purple color is seen.	Presence of Na, possibly also of K.
	2. A pale violet flame appearing reddish-purple when seen through the indigo-prism.	Presence of K and absence of Na.
	[3. No color is imparted to the flame.]	[Absence of K and Na.]
III. Boil some of the solution in a test-tube with $\text{KHO}$ solution, and try whether $\text{NH}_3$ is given off by smelling; or if it cannot be smelt, by testing with moist red litmus paper (51 b), or strong $\text{HCl}$ .	1. $\text{NH}_3$ gas is smelt, or the red litmus becomes blue, and white fumes are obtained with $\text{HCl}$ .	Presence of $\text{NH}_4$ .
	[2. No $\text{NH}_3$ is smelt, and moist red litmus paper is not blued, neither are white fumes obtained with $\text{HCl}$ .]	[Absence of $\text{NH}_4$ .]
IV. A confirmatory test for K may be made by adding $\text{PtCl}_4$ or $\text{NaHT}$ . If Exp. III has shown that $\text{NH}_4$ is present, proceed to test for K as under A; if $\text{NH}_4$ is absent, test for K as under B. (See 66, 5.)		
A. Boil some more of the solution down to dryness in a porcelain dish; scrape out the white solid substance left in the dish, and heat it strongly on a piece of platinum foil with turned-up edges as long as any white fumes are seen to be given off when the foil is taken out of the flame for an instant. Place the foil when cold in a test-tube, and boil it with <i>very little</i> water, to which two or three drops of dilute $\text{HCl}$ have been added; cool the liquid, and pour it out upon a watch-glass, add a little $\text{PtCl}_4$ ,† and stir well. A yellow precipitate forms:— <i>Presence of K</i> .		B. Pour a little of the solution upon a watch-glass, add $\text{PtCl}_4$ , and stir well: A yellow precipitate:— <i>Presence of K</i> . If no precipitate forms, K may nevertheless be present, the solution being too dilute to allow the precipitate to form; therefore pour some of the solution into a porcelain dish, boil down nearly to dryness, cool and test this liquid by stirring it with $\text{PtCl}_4$ on a watch-glass, adding alcohol if the precipitate does not appear at once: A yellow precipitate:— <i>Presence of K</i> . <i>Note</i> .—Addition of alcohol will frequently produce the precipitate without evaporation.

\* Almost all substances contain traces of Na, and therefore when heated in the Bunsen flame tinge it more or less yellow: hence, unless the yellow coloration is very intense, enter in the results "Na trace."

† In this and all subsequent places where  $\text{PtCl}_4$  is directed to be used as a test for K, the  $\text{NaHT}$  test may be substituted for economical reasons.

*Note.*—Separate portions of the solution are to be used for each of these experiments. It must be understood that *only one* of the numbered results in the second column can be obtained in each experiment, but one or other of these must invariably be observed. In subsequent tables, that result only is stated in the second column which indicates the presence of a substance, and the absence of that result proves the absence of the substance; the portions in brackets will therefore be omitted in future.

Exp. IV may be omitted by a beginner, K being tested for by the flame coloration alone (Exp. II).

After writing out the tests tried in the above form, state at the end what has been found in the solution: thus—"Found, K and Na." This is to be done after writing out each analysis.

#### GROUP IV.—BARIUM GROUP.

68. This group includes Ba, Sr, Ca.

The members of this group differ from those of Group V, by being precipitated as carbonates by  $\text{Am}_2\text{CO}_3$ , even in the presence of  $\text{AmCl}$ .\* They are not precipitated by any other of the group reagents.

BARIUM (Ba).—Use  $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ .

69.  $\text{Am}_2\text{CO}_3$  added after some  $\text{AmCl}$ : white precipitate ( $\text{BaCO}_3$ ), which is at first flocculent, but if heated and allowed to stand slowly shrinks in bulk and becomes crystalline.

Pour a little of this into another test-tube and add to it some  $\text{H}\bar{\text{A}}$ , the precipitate dissolves entirely if sufficient  $\text{H}\bar{\text{A}}$  is added.

70.  $\text{CaSO}_4$  (or  $\text{H}_2\text{SO}_4$ ): heavy white precipitate ( $\text{BaSO}_4$ ): insoluble in  $\text{HCl}$  even on boiling.

71.  $\text{K}_2\text{CrO}_4$  added after some  $\text{H}\bar{\text{A}}$ : yellow precipitate ( $\text{BaCrO}_4$ ), soluble in warm  $\text{HCl}$ , insoluble in warm  $\text{H}\bar{\text{A}}$ . (Diff. from Sr.).

72.†  $\text{H}_2\text{SiF}_6$ : semi-transparent precipitate ( $\text{BaSiF}_6$ ): the precipitate often does not appear in dilute solution until after some time, or until the liquid is boiled, or alcohol is added. Shaking or stirring this liquid hastens its formation.

73.  $\text{Am}_2\text{C}_2\text{O}_4$ : white precipitate ( $\text{BaC}_2\text{O}_4$ ).

74. *Flame Coloration*:  $\text{BaCl}_2$  gives a yellowish-green color to the flame, visible through the indigo-prism.

\* On boiling with  $\text{AmCl}$  solution, however, the precipitates are more or less perfectly dissolved.

† A reaction in small type is comparatively unimportant, and may be omitted by beginners.

The spectrum consists of a number of lines, the most characteristic of which are three green lines ( $\alpha$ ,  $\beta$ ,  $\gamma$ ).

STRONTIUM (Sr).—Use  $\text{Sr}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$ .

75.  $\text{Am}_2\text{CO}_3$  added after some  $\text{AmCl}$ : white precipitate ( $\text{SrCO}_3$ ): soluble in  $\text{H}\bar{\text{A}}$ . This precipitate is flocculent at first; but if heated it quickly becomes crystalline and shrinks very considerably.

75 a.  $\text{CaSO}_4$  (or  $\text{H}_2\text{SO}_4$ ): white precipitate ( $\text{SrSO}_4$ ): this precipitate does not usually form at once in a cold solution, but only after standing for some time. The precipitate, however, *appears at once on boiling the liquid*.

76.  $\text{K}_2\text{CrO}_4$  added after some  $\text{H}\bar{\text{A}}$ : no precipitate, since  $\text{SrCrO}_4$  is soluble in  $\text{H}\bar{\text{A}}$ .

77.  $\text{H}_2\text{SiF}_6$ : no precipitate.

78.  $\text{Am}_2\text{C}_2\text{O}_4$ : white precipitate ( $\text{SrCr}_2\text{O}_4$ ).

79. *Flame coloration*: crimson-red; this color appears *intense red* through the indigo-prism unless the flame-coloration is very faint.

The strontium spectrum contains many lines; the most characteristic are the orange line ( $\alpha$ ), the red lines ( $\beta$ ,  $\gamma$ ), and the blue line ( $\delta$ ).

CALCIUM (Ca).—Use  $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$ .

80.  $\text{Am}_2\text{CO}_3$  added after  $\text{AmCl}$ : white precipitate ( $\text{CaCO}_3$ ), soluble in  $\text{H}\bar{\text{A}}$ . This precipitate is flocculent at first, but on standing for some time or on being gently heated, it shrinks considerably and becomes crystalline.

81.  $\text{CaSO}_4$ : no precipitate even on standing or on being boiled.

82.  $\text{H}_2\text{SO}_4$ : white precipitate ( $\text{CaSO}_4$ ) forms at once in strong solutions, and often in weak solutions on being boiled; but since  $\text{CaSO}_4$  is not quite insoluble in water, some of it will always remain dissolved; prove this by boiling the above liquid containing the precipitate and filtering it, then keep adding  $\text{AmHO}$  to the filtrate and stirring it until a drop of the solution turns red litmus blue and the liquid smells of  $\text{NH}_3$ , then add  $\text{Am}_2\text{C}_2\text{O}_4$ , a white precipitate will form, showing the presence of

102 GROUP IV.—TABLE OF DIFFERENCES. [83-88.]

Ca (see 85 below) which must have been dissolved as  $\text{CaSO}_4$ .

83.  $\text{K}_2\text{CrO}_4$ : no precipitate.

84.  $\text{H}_2\text{SiF}_6$ : no precipitate.

85.  $\text{Am}_2\text{C}_2\text{O}_4$ : white precipitate ( $\text{CaC}_2\text{O}_4$ ), soluble in acids except  $\text{H}\bar{\text{A}}$  and  $\text{H}_2\text{C}_2\text{O}_4$ .

86. *Flame Coloration*: yellowish-red; this color viewed through the indigo-prism *does not appear red but dingy green*, thus differing from that given by Sr.

In the calcium-spectrum the most characteristic lines are the green line ( $\beta$ ), and the intense orange line ( $\alpha$ ).

87. GROUP IV.—TABLE OF DIFFERENCES.

The first three tests only need be used for the detection of a member of this group occurring by itself in solution: test 1 serves also for solids.

Tests.	Ba—Salts.	Sr—Salts.	Ca—Salts.
1. Flame coloration.	Greenish yellow.	Intense crimson, appears <i>red</i> through the indigo-prism.	Yellowish red, appears <i>dingy green</i> through the indigo-prism.
2. $\text{CaSO}_4$ added to the cold solution.	An immediate precipitate.	A precipitate appearing only after a short time or immediately on boiling.	—
3. $\text{K}_2\text{CrO}_4$ added after $\text{H}\bar{\text{A}}$ .	Light yellow precipitate.	—	—
4. $\text{H}_2\text{SiF}_6$ added.	White precipitate.	—	— ]
5. $\text{H}_2\text{SO}_4$ added in excess and boiled.	Entirely precipitated as $\text{BaSO}_4$ .	Entirely precipitated as $\text{SrSO}_4$ .	Partially precipitated as $\text{CaSO}_4$ ; hence the filtrate from the precipitate contains Ca, which may be detected by adding $\text{AmHO}$ in excess, and then $\text{Am}_2\text{C}_2\text{O}_4$ ; a white precipitate forms.

When analyzing a solution which is known to contain only one member of this group, its detection by the first three of the above tests will be exceedingly simple.

88. (s.) *The separation and detection of Ba, Sr, and Ca, when mixed*, depends upon the following differences:

1. The flame colorations, examined if necessary through the indigo-prism, or better by means of the spectroscope.

2. The difference in behavior with  $\text{CaSO}_4$ , which at once indicates the presence or absence of Ba.

3. The separation of Ba, if present, by  $\text{K}_2\text{CrO}_4$  in the presence of  $\text{H}\bar{\text{A}}$ .

4. The precipitation of Sr by  $\text{CaSO}_4$  on being boiled, which will indicate the presence of Sr in the absence, or after the separation, of Ba.

5. The precipitation of Sr, if present, by boiling with  $\text{H}_2\text{SO}_4$ ; sufficient  $\text{CaSO}_4$  then remaining in the solution to give, after making the filtrate alkaline with  $\text{AmHO}$ , a precipitate with  $\text{Am}_2\text{C}_2\text{O}_4$ .

89. (s.) A solution which has to be-examined for Ba, Sr, and Ca, and can contain only these substances, is made alkaline, if not already so, by addition of  $\text{AmHO}$ ; then  $\text{Am}_2\text{CO}_3$  is added as long as it is seen to cause any precipitate; the liquid is then filtered (23), and a little more  $\text{Am}_2\text{CO}_3$  added to the clear filtrate; if any further precipitate forms, more  $\text{Am}_2\text{CO}_3$  is added and the liquid is again poured through the same filter, pouring the first portion of the filtrate once more through the filter as it is sure to come through turbid. As soon as the filtrate gives no further precipitate with  $\text{Am}_2\text{CO}_3$ , all the members of this group which were present in the solution have been precipitated as carbonates, and on filtration are obtained on the filter. The precipitate is then examined by Table IV (438).

*Note.*—In using the group tables in Section VI it must be understood that they are drawn out to meet the case of all the members of the group, being present. If therefore in any case no precipitate forms on addition of a reagent, the substance or substances whose presence would have been indicated by the formation of the precipitate must be absent. Hence a filtration which is directed to be made is often unnecessary, and the solution itself is treated as is directed for the filtrate.

### GROUP III A.—IRON GROUP.

90. This group includes Al, Fe, Cr.

The members of this group differ from those of Groups III B, IV, and V, by being precipitated by  $\text{AmHO}$  after addition of  $\text{AmCl}$ ; they are not precipitated however by the group reagents for Groups II and I.



This group is also precipitated by  $\text{Am}_2\text{S}$ , or by  $\text{H}_2\text{S}$  added to the alkaline solution.

Group III A further differs from Group III B in being completely precipitated by adding  $\text{BaCO}_3$  shaken up with water; this reagent affords the most perfect means of separating the two groups.

The members of this group show no characteristic flame colorations; but, with the exception of Al, they give borax beads of characteristic color.


**ALUMINIUM (Al).**—Use  $\text{AlAm}(\text{SO}_4)_3 \cdot 12\text{H}_2\text{O}$  or  $\text{AlK}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ , ammonia- or potash-alum.

91.  $\text{AmHO}$ : white flocculent precipitate ( $\text{Al}_2\text{H}_6\text{O}_6$ ), which is seen best on boiling the liquid: soluble in  $\text{HCl}$  and in  $\text{H}\bar{\text{A}}$ ; slightly soluble in  $\text{AmHO}$ , especially in the absence of  $\text{AmCl}$ .



92.  $\text{KHO}$  added in very small quantity: same precipitate as  $\text{AmHO}$ : soluble in excess of  $\text{KHO}$ . If to a part of the  $\text{KHO}$  solution  $\text{HCl}$  be added gradually until it is neutral, the  $\text{Al}_2\text{H}_6\text{O}_6$  is precipitated again; if more acid is added the precipitate dissolves, but is precipitated again by adding  $\text{AmHO}$  in excess. From another portion of the  $\text{KHO}$  solution the  $\text{Al}_2\text{H}_6\text{O}_6$  may be reprecipitated by addition of sufficient  $\text{AmCl}$ .

93.  $\text{Am}_2\text{S}$ : same precipitate ( $\text{Al}_2\text{H}_6\text{O}_6$ ),  $\text{H}_2\text{S}$  gas coming off or remaining dissolved in the liquid:



94. *Blowpipe Reaction.*—Some solid  $\text{AlAm}(\text{SO}_4)_3 \cdot 12\text{H}_2\text{O}$  heated on charcoal in the outer blowpipe flame, then moistened, after cooling, with  $\text{Co}(\text{NO}_3)_2$  solution and again heated in the outer flame, gives a *fine blue mass*.

95. **IRON (Fe).** *Note.*—Two classes of iron compounds are known, which differ in appearance and properties, and behave differently with reagents; they are distinguished as *ferrous* and *ferric* compounds respectively, the former being supposed to contain the metal *ferrosium*


(Fe''),\* the latter *ferricum* (Fe'''). It is usual in stating analytical results to mention in which state of combination the iron exists, and therefore some distinctive tests are described below (101). The whole of the reactions of ferrous compounds are not given, because Fe, if present, is always separated and detected in the course of analysis as ferricum, and ferrosium is tested for by special reactions in the original substance. Ferrous are readily converted into ferric compounds by boiling for a short time with a little strong  $\text{HNO}_3$ , or with  $\text{HCl}$  and a crystal of  $\text{KClO}_3$ .

**FERRICUM (Fe''').**—Use  $\text{Fe}_2\text{Cl}_6$ .

Ferric salts are usually yellow or reddish-yellow; ferrous salts usually pale green, or white, if perfectly dry.


**96.**  $\text{AmHO}$  (or  $\text{KHO}$ ): reddish-brown, flocculent precipitate  $\text{Fe}_2\text{Ho}_6$ : insoluble in  $\text{KHO}$ , soluble in  $\text{HCl}$ .

**97.**  $\text{Am}_2\text{S}$ : a black precipitate ( $\text{FeS}$ ), mixed with white  $\text{S}$ : soluble in boiling acids; insoluble in  $\text{KHO}$ :



In a very weak solution of Fe a green color is produced at first by  $\text{Am}_2\text{S}$ , and the black precipitate separates only after a time.

**98.**  $\text{H}_2\text{S}$ :† same black precipitate in alkaline solutions: in neutral or acid solutions  $\text{S}$  precipitates, and the ferric is changed to a ferrous salt, the color of the solution changing from yellow to pale-green as is seen after boiling and filtering.



**Blowpipe Reactions.**—These are the same for ferrous as for ferric compounds.

**99.** Fused with  $\text{Na}_2\text{CO}_3$  on charcoal in the inner flame, a gray magnetic powder is left (see 33 a); this is shown to be magnetic by being attracted when touched under

\* This distinction of the two conditions of Fe and other elements by dashes or Roman numbers placed above the symbol, first proposed by Odling, is very convenient for brevity.

†  $\text{H}_2\text{S}$  may be added as a solution of the gas in trying the reactions; in the course of analysis the gas is usually passed into the liquid.

water by the end or pole of a magnet, or by a magnetized knife-blade.

100. *Borax Bead:* { Outer flame: *reddish-brown* while hot: *yellow* when cold.  
Inner “ *greenish* bead (the color of common green bottle glass).

### 101. DISTINCTIVE TESTS FOR FERROUS AND FERRIC SALTS.

*Note.*—For these tests several drops of dilute HCl should be first added to the Fe solution.

Reagents to be added.	Ferric Salts.	Ferrous Salts. (For Ferrous Salt use $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ solution.)
1. $\text{K}_4\text{FeCy}_6$ added in small quantity.	<i>Dark-blue precipitate</i> (“Prussian blue”): insoluble in HCl,* soluble in $\text{H}_2\text{C}_2\text{O}_4$ and slightly soluble in $\text{K}_4\text{FeCy}_6$ added in excess: turned brown by $\text{KHO}$ .	<i>Light-blue precipitate</i> , becoming dark-blue in the air, or on addition of $\text{HNO}_3$ or Cl-water.
2. $\text{K}_2(\text{FeCy}_6)_2$	<i>No precipitate</i> : the solution darkens, but on dilution with water is seen to contain no precipitate.	<i>Dark-blue precipitate</i> (“Turnbull’s Blue”); insoluble in HCl. In very dilute solution only a dark bluish-gray color is produced.
3. $\text{KCys}$ .	<i>Blood-red coloration</i> : no precipitate is produced, the liquid being perfectly clear on dilution; the color is immediately destroyed by adding $\text{HgCl}_2$ solution in sufficient quantity; its production is also hindered by $\text{HNO}_3$ and by $\text{H}_2\text{A}$ .	<i>No red coloration</i> unless small quantities of ferric salts are present.

**CHROMIUM (Cr).** Use  $\text{CrK}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ .

Chromic salts are usually green or violet in color.

102. *AmHO*: pale bluish-green precipitate ( $\text{Cr}_2\text{H}_6\text{O}_6$ ): if *AmHO* is added in large quantity and the liquid is

\* The solubility and insolubility of this precipitate must be shown by warming separate portions of the blue liquid containing the precipitate with HCl,  $\text{H}_2\text{C}_2\text{O}_4$  and  $\text{K}_4\text{FeCy}_6$ , then separately filtering each: if the precipitate has been dissolved, the filtrate will be dark-blue.

heated, some of the precipitate is dissolved, producing a beautiful violet-red solution whose color is best seen after filtering; but from this solution the  $\text{Cr}_2\text{H}_6\text{O}_6$  is precipitated again on boiling the liquid for several minutes in a porcelain dish.

103.  $\text{KHO}$  added in small quantity gives the same precipitate ( $\text{Cr}_2\text{H}_6\text{O}_6$ ): if more cold  $\text{KHO}$  is added the precipitate is entirely dissolved to a green fluid; on diluting this liquid with water and boiling for several minutes, the  $\text{Cr}_2\text{H}_6\text{O}_6$  is precipitated again, and the liquid becomes colorless:  $\text{AmCl}$  added to the  $\text{KHO}$  solution also precipitates again the  $\text{Cr}_2\text{H}_6\text{O}_6$ .

103 a. If to some of the green liquid, obtained by adding  $\text{KHO}$  in excess to the chromium solution, a little red-lead or lead-peroxide ( $\text{PbO}_2$ ) is added, and the liquid is then boiled, a yellow liquid is obtained from which a yellow precipitate ( $\text{PbCrO}_4$ ) is thrown down by addition of  $\text{H}\bar{\text{A}}$  in excess.

104.  $\text{Am}_2\text{S}$  precipitates  $\text{Cr}_2\text{H}_6\text{O}_6$ :  $\text{H}_2\text{S}$  gas being given off or remaining dissolved. The reaction is similar to that with  $\text{AlAm}(\text{SO}_4)_3$ , par. 93.

105. *Blowpipe Reaction*.—If  $\text{Na}_2\text{CO}_3$  and  $\text{KNO}_3$  be mixed with some solid  $\text{CrK}(\text{SO}_4)_2$ , or with the dried precipitate produced by  $\text{AmHO}$  or  $\text{KHO}$ , and the mixture be fused on a piece of platinum foil or in a porcelain crucible or lid, or in a loop of platinum wire as was directed for the borax bead (31), a yellow mass is formed colored by  $\text{Na}_2\text{CrO}_4$ . If this yellow mass is dissolved by boiling it with water, a yellow solution is obtained: this solution, if made acid with  $\text{H}\bar{\text{A}}$  and boiled for a few minutes to drive off  $\text{CO}_2$ , gives with  $\text{Pb}\bar{\text{A}}_2$  solution a yellow precipitate of  $\text{PbCrO}_4$ .

106. *Note*.—Cr forms two classes of compounds: in one class Cr in combination with oxygen functions as an acid radicle: this class includes the chromates such as  $\text{K}_2\text{CrO}_4$  and  $\text{K}_2\text{Cr}_2\text{O}_7$ , they are usually yellow or reddish in color; but Cr also forms a series of salts in which it acts as a base; these are usually green or violet, and give the above reactions for Cr. The latter compounds pass by oxidation into the former, as in reactions (103 a) and (105) where the oxidation is caused by  $\text{PbO}_2$  and  $\text{KNO}_3$  respectively. Chromates pass by reduction into the green compounds; examples of this change will be found under the tests for chromic acid (286).

# 108 GROUP III A.—TABLE OF DIFFERENCES. [107.

This reduction of chromic acid causes it to be detected in the course of analysis as a base, and a special experiment has to be performed to ascertain whether the Cr was present originally as a base or as an acid radicle.

## 107. GROUP III A.—TABLE OF DIFFERENCES.

Tests.	Al—Salts.	Fe'''—Salts.	Fe''—Salts.	Cr—Salts.
<i>a. For liquids.</i>				
1. Add AmHO.	White flocculent precipitate.	Reddish-brown flocculent precipitate.	Dingy - green precipitate, rapidly turning brown when left exposed to the air.	Pale-green flocculent precipitate, color unaltered by exposure to air.
2. Add KHO.	White flocculent precipitate, easily soluble in excess of KHO.	Reddish-brown flocculent precipitate, insoluble in excess of KHO.	The same precipitate as with AmHO, insoluble in excess of KHO.	Pale-green precipitate, soluble in excess of cold KHO, but reprecipitated on diluting and boiling for some time.
3. Add KCyS and a few drops of HCl.	— •	Deep blood-red coloration, destroyed by HgCl <sub>2</sub> solution.	No coloration unless Fe''' is also present.	—
<i>b. Blowpipe tests for solids.</i>				
4. Fuse with Na <sub>2</sub> O <sub>2</sub> and KNO <sub>3</sub> on platinum foil.	—	On cooling, a white mass of Na <sub>2</sub> CO <sub>3</sub> remains with dark-brown particles of Fe <sub>2</sub> O <sub>3</sub> .	Same as Fe''.	On cooling, a yellow mass remains, easily soluble in water; the solution, after having been boiled with excess of HA, gives a yellow precipitate with PbA <sub>2</sub> .
5. Heated on charcoal.	In the outer flame when cool if moistened with Co (NO <sub>2</sub> ) <sub>2</sub> solution and reheated strongly gives a fine blue mass.	In inner flame mixed with Na <sub>2</sub> CO <sub>3</sub> gives a gray magnetic powder. (33a.)	Same as Fe''.	—
6. Fused in borax bead.	—	Reddish-yellow in outer flame. Greenish-yellow in inner flame.	Same as Fe''.	Green both in outer and inner flames.

Members of this group occurring singly are easily distinguished by one or other of the above tests, the first three serving for liquids, the last three for solids; tests 2 and 5 are the best for Al, 2, 4, and 6 for Cr, and 1, 3, or 6 for Fe.

**108.** (s.) *The separation and detection of these three substances when mixed* depend upon the following differences:

1. The solubility of  $\text{Al}_2\text{H}_6$  in boiling  $\text{KHO}$ , in which  $\text{Fe}_2\text{H}_6$  and  $\text{Cr}_2\text{H}_6$  are insoluble.

2. The conversion of  $\text{Cr}_2\text{H}_6$  by fusion with  $\text{Na}_2\text{CO}_3$  into soluble  $\text{Na}_2\text{CrO}_4$ :  $\text{Fe}_2\text{H}_6$  remaining as  $\text{Fe}_2\text{O}_3$  which is insoluble in water.

3. The yellow color of the solution of  $\text{Na}_2\text{CrO}_4$  in water, and production of a yellow precipitate by acidifying it with  $\text{HAc}$  and adding  $\text{PbAc}_2$ .

4. The blood-red coloration obtained by adding  $\text{KCys}$  to the  $\text{Fe}_2\text{O}_3$  dissolved in  $\text{HCl}$ .

**109.** (s.) A solution which may contain Al, Fe, Cr, is examined in the following manner:

To a few drops of the solution acidified with  $\text{HCl}$  add a little  $\text{K}_4\text{FeCy}_6$ ; if a blue precipitate is produced Fe is present; this portion is rejected. Two other small portions of the solution also acidified are then tested; one with  $\text{K}_6(\text{FeCy}_6)_2$ , which by giving a dark-blue precipitate shows the presence of  $\text{Fe}''$ ; the other with  $\text{KCys}$ , which if it gives a blood-red color proves the presence of  $\text{Fe}'''$ .

Boil the rest of the solution for several minutes, after adding a few drops of strong  $\text{HNO}_3$  if  $\text{Fe}''$  is present; add some  $\text{AmCl}$ , then  $\text{AmHO}$  until the liquid after being well stirred is alkaline and smells of  $\text{NH}_3$ ; boil, filter, and examine the precipitate by Table III A (436), disregarding all the appended notes.

---

### GROUP III B.—ZINC GROUP.

**110.** This group includes Zn, Mn, Ni, Co; its members differ from those of Group IV and V, by being precipitated by  $\text{Am}_2\text{S}$  in neutral or alkaline solution, and

by  $\text{H}_2\text{S}$  in alkaline solution; they are not precipitated by the group reagents for Groups III A, II, and I. Mn is however liable to be precipitated by  $\text{AmHO}$ , even in the presence of  $\text{AmCl}$ , if the solution stands exposed to air for some time.


The members of this group show no characteristic flame colorations, but with the exception of Zn give characteristic colors to a borax bead.

$\text{BaCO}_3$  does not precipitate this group as it does Group III A, unless its members are present as sulphates.


ZINC (Zn).—Use  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ .

111.  $\text{Am}_2\text{S}$ : white precipitate ( $\text{ZnS}$ ): the precipitate often appears yellow from the presence of excess of yellow  $\text{Am}_2\text{S}$  in the liquid: its true color is seen in the next reaction, or by letting the precipitate produced by  $\text{Am}_2\text{S}$  settle, or by filtering it. For the solubility of this precipitate see the end of the next reaction.

112.  $\text{H}_2\text{S}$ .\* white precipitate ( $\text{ZnS}$ ): the Zn is only partly precipitated from a neutral solution since  $\text{H}_2\text{SO}_4$  is separated and dissolves the  $\text{ZnS}$ :



but if an alkali ( $\text{KHO}$  or  $\text{AmHO}$ ) be added to neutralize the  $\text{H}_2\text{SO}_4$  when it is set free, the Zn may be entirely precipitated: addition of  $\text{Na}\bar{\text{A}}$  also causes complete precipitation, since it replaces  $\text{H}_2\text{SO}_4$  in the solution by  $\text{H}\bar{\text{A}}$ , and  $\text{ZnS}$  is insoluble in  $\text{H}\bar{\text{A}}$ :



Add  $\text{H}_2\text{S}$  to some  $\text{ZnSO}_4$  solution made alkaline with a few drops of  $\text{AmHO}$ , and show by adding to separate portions of this liquid  $\text{HCl}$ ,  $\text{H}\bar{\text{A}}$  and  $\text{Am}_2\text{S}$ , that  $\text{ZnS}$  is soluble in  $\text{HCl}$ , insoluble in  $\text{H}\bar{\text{A}}$  and in  $\text{Am}_2\text{S}$ .†

\*  $\text{H}_2\text{S}$  may be used in this group as sulphuretted hydrogen water.

† The tests which show the solubility of the sulphides of this group are best tried on the precipitate obtained by  $\text{AmHO}$  and  $\text{H}_2\text{S}$ , unless freshly prepared colorless  $\text{Am}_2\text{S}$  can be obtained, since from yellow  $\text{Am}_2\text{S}$  acids precipitate sulphur.

113. *KHO* added in small quantity gives a white precipitate ( $\text{ZnHo}_2$ ); if more *KHO* is added, this precipitate dissolves; the  $\text{ZnHo}_2$  is however precipitated again as such by adding much water to this solution and boiling it, or as  $\text{ZnS}$  by passing  $\text{H}_2\text{S}$  gas into it.

114. Solid  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$  powdered with  $\text{Na}_2\text{CO}_3$  in a mortar, then heated on charcoal in the inner blowpipe flame, gives a white incrustation on the charcoal, which is *yellow* whilst hot. It cannot be driven away by the outer blowpipe flame, but is easily removed by the inner flame. If this incrustation, on cooling, is moistened with cobalt-nitrate solution and strongly heated in the outer blowpipe flame it becomes *green*.

*Note.*—The above changes of color are more distinctly obtained by igniting a little solid  $\text{ZnSO}_4$  in the outer blowpipe flame on charcoal, it is *yellow* whilst hot, and *white* when cold: if moistened with  $\text{Co}(\text{NO}_3)_2$  solution, and reheated in the outer flame, it becomes *green*.

#### MANGANESE (Mn).—Use $\text{MnCl}_2$ or $\text{MnSO}_4$ .

Manganous salts are usually of a pale pink color. Alkaline manganates are green, permanganates purple.

115.  $\text{Am}_2\text{S}$ : flesh-colored or pinkish precipitate ( $\text{MnS}$ ): the liquid should be filtered, since the precipitate often appears discolored by the yellow  $\text{Am}_2\text{S}$ , the color of the precipitate on the filter is then easily seen; the color of the moist precipitate darkens to brown on standing in the air upon the filter.

For the solubility of this precipitate see Exp. (116).

116.  $\text{H}_2\text{S}$  precipitates  $\text{MnS}$  partly from neutral solutions, entirely from alkaline solutions, but not at all in presence of free  $\text{HCl}$  or  $\text{H}\bar{\text{A}}$ : show with separate portions of the  $\text{MnS}$  precipitate, obtained by adding  $\text{H}_2\text{S}$  solution to the liquid after some  $\text{AmHO}$ , that  $\text{MnS}$  is soluble in  $\text{HCl}$  and in  $\text{H}\bar{\text{A}}$ , but insoluble in  $\text{Am}_2\text{S}$ .

117. *KHO*: white precipitate ( $\text{MnHo}_2$ ), quickly turned brown by pouring the liquid containing the precipitate into a white porcelain dish or upon a filter: the precipitate is insoluble in excess of *KHO*.



**118.** *AmHO* gives the same precipitate, but if sufficient *AmCl* is first added, *AmHO* produces no immediate precipitate, the solution, however, on standing exposed to the air turns brown, and the *Mn* is gradually precipitated as brown *Mn<sub>2</sub>Ho<sub>6</sub>*.

**118 a.** Pour a little *MnSO<sub>4</sub>* (not *MnCl<sub>2</sub>*) solution upon some red or brown lead oxide (*PbO<sub>2</sub>*), add *HNO<sub>3</sub>*, which must be quite free from *Cl*, boil the mixture and allow the powder to settle; the clear liquid is colored deep red by the formation of permanganic acid (*HMnO<sub>4</sub>*). The production of the color is prevented by the presence of chlorine in any soluble form of combination.

**119.** *Blowpipe Tests*.—If some solid *MnCl<sub>2</sub>* be fused in the outer flame with a mixture of *Na<sub>2</sub>CO<sub>3</sub>* and a little *KNO<sub>3</sub>*, a *bluish-green mass* is obtained on cooling. The mixture may be fused on a piece of platinum foil by heating the lower surface of the foil with the blowpipe flame; or a bead of the *Na<sub>2</sub>CO<sub>3</sub>* and *KNO<sub>3</sub>* mixed may be made in a loop of platinum wire, and the experiment tried just as with a borax bead, heating it in the extreme tip of the outer flame.

**119 a.** Fused with *Na<sub>2</sub>CO<sub>3</sub>* on charcoal in the inner flame a gray magnetic powder is obtained (33 a).

**120.** *Borax Bead*.—Use *very little\** solid *MnCl<sub>2</sub>*:

In the outer flame { *Violet-red* bead whilst hot.  
                              *Amethyst-red* on cooling.

In the inner flame.—A colorless bead.

---

NICKEL (*Ni*).—Use *NiSO<sub>4</sub>·7H<sub>2</sub>O*.

Nickel salts are usually pale bright green in color.

**121.** *Am<sub>2</sub>S*: Black precipitate (*NiS*); add more *yel-*

---

\* Borax beads are often opaque from the use of too much substance; in such a case fuse the bead in the blowpipe flame, then by a sudden jerk throw some of the fused mass off, and fuse again the remaining portion with fresh borax; repeat this if necessary. The fused bead may also be flattened by squeezing with small pin-cers.

low  $\text{Am}_2\text{S}_3$ ; \* boil and filter; a brown filtrate runs through, colored by  $\text{NiS}$  dissolved in the excess of  $\text{Am}_2\text{S}_3$ ; pour this filtrate into a porcelain dish and boil for some time, adding distilled water if necessary to prevent the dish becoming dry; the black  $\text{NiS}$  will be precipitated and may be filtered off, giving a colorless filtrate. If  $\text{H}\bar{\text{A}}$  is added to the dark filtrate until it is acid, the  $\text{NiS}$  is also precipitated from it.

122.  $\text{H}_2\text{S}$ : black precipitate ( $\text{NiS}$ ) in neutral solutions or in solutions acid with  $\text{H}\bar{\text{A}}$ , but  $\text{HCl}$  prevents the precipitation; show with portions of the liquid containing  $\text{NiS}$ , obtained by adding  $\text{H}_2\text{S}$  to  $\text{NiSO}_4$  solution to which a little  $\text{AmHO}$  has been added, that  $\text{NiS}$  is insoluble in cold dilute  $\text{HCl}$  and in  $\text{H}\bar{\text{A}}$ , but soluble when heated with  $\text{HCl}$  after adding a crystal of  $\text{KClO}_3$ .

123.  $\text{KHO}$ : light-green precipitate ( $\text{NiHo}_2$ ).

124.  $\text{AmHO}$  added in very small quantity; bluish-green precipitate ( $\text{NiHo}_2$ ); soluble in excess of  $\text{AmHO}$  to a violet-blue liquid; soluble also in  $\text{AmCl}$ . If  $\text{AmCl}$  be first added  $\text{AmHO}$  causes no precipitate.

125.  $\text{KCy}$  added in small quantity gives a yellowish-green precipitate ( $\text{NiCy}_2$ ); by further addition of  $\text{KCy}$  this precipitate is redissolved, but  $\text{HCl}$  again precipitates  $\text{NiCy}_2$  from this solution.

126. If some nickel sulphate solution is made acid, with several drops of  $\text{HCl}$ , then  $\text{KCy}$  gradually added, whilst stirring or shaking the liquid, until the precipitate is redissolved, and the solution thus obtained is boiled for a short time, then cooled and divided into two parts;  $\text{HCl}$  added in excess to one part will produce a precipitate of  $\text{NiCy}_2$ , often only appearing after a short time;  $\text{Na}_2\text{Cl}_2\text{O}$ , or  $\text{Br}$ -water after excess of  $\text{NaHO}$ , added to the other part will give on warming a black precipitate ( $\text{Ni}_2\text{Ho}_3$ ).

127. *Blowpipe Reactions*.—Solid  $\text{NiSO}_4 \cdot 7\text{H}_2\text{O}$  powdered with  $\text{Na}_2\text{CO}_3$  in a mortar and then fused on char-

---

\* The  $\text{Am}_2\text{S}_3$  must be yellow; colorless  $\text{Am}_2\text{S}_3$  does not dissolve  $\text{NiS}$ .

coal in the inner blowpipe flame, leaves a gray powder which is attracted by the magnet. (See 33 a.)

**128. Borax Bead.**

In outer flame	{ Violet or sherry-red bead whilst hot. Pale yellow on cooling.
In inner flame, after being heated for some time.	
	{ Blackish or opaque bead.

COBALT (Co).—Use  $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ .

Cobalt salts are usually reddish-pink in color.

**130.  $\text{Am}_2\text{S}$ :** black precipitate ( $\text{CoS}$ ); on adding much  $\text{Am}_2\text{S}$ , boiling and filtering, the filtrate is colored yellow by  $\text{Am}_2\text{S}$  and is not dark-colored, since  $\text{CoS}$  is insoluble in  $\text{Am}_2\text{S}$  (difference from  $\text{NiS}$ ). For the solubility of  $\text{CoS}$  see Exp. 131.

**131.  $\text{H}_2\text{S}$ :** black precipitate ( $\text{CoS}$ ), only forms in alkaline solutions or in solutions acid with  $\text{H}\bar{\text{A}}$ ;  $\text{HCl}$  prevents the precipitation. Add  $\text{H}_2\text{S}$ -water to some  $\text{Co}(\text{NO}_3)_2$  solution, first made alkaline with a few drops of  $\text{AmHO}$ , and pour into separate portions of this liquid  $\text{HCl}$  and  $\text{H}\bar{\text{A}}$ , the precipitate does not dissolve; to the portion containing  $\text{HCl}$  add a crystal of  $\text{KClO}_3$  and heat, the precipitate dissolves readily.

**132.  $\text{KHO}$ :** blue precipitate ( $\text{CoHo}_2$ ), turning green if poured out upon a watch-glass and allowed to stand in the air, and becoming pale red on being boiled; the red color is, however, frequently more or less masked by a brown cobalt hydrate precipitated at the same time.

**133.  $\text{AmHO}$ :** bluish-green precipitate, having the same properties as that given by  $\text{KHO}$ , soluble in  $\text{AmCl}$ ; hence if  $\text{AmCl}$  is added before  $\text{AmHO}$  no precipitate is produced.

**134.  $\text{KC}_y$**  added in small quantity gives a reddish-brown precipitate ( $\text{CoCy}_2$ ); add more  $\text{KC}_y$  slowly and whilst shaking the solution the precipitate dissolves; now add  $\text{HCl}$ , the  $\text{CoCy}_2$  is precipitated again.

**135.** If some  $\text{Co}(\text{NO}_3)_2$  solution is made acid with a few drops of  $\text{H}\bar{\text{A}}$ , then  $\text{KC}_y$  added slowly until the pre-

precipitate at first formed just redissolves, and the liquid is boiled until no small of  $\text{HCy}$  is evolved, then cooled and divided into three parts, it will be found that neither  $\text{HCl}$ , nor  $\text{Na}_2\text{Cl}_2\text{O}$ , nor Br-water with excess of  $\text{NaHO}$  solution, produces a precipitate on warming the liquid. [Difference from  $\text{Ni}$ , see (126).]

*Note.*—The difference of behavior of the  $\text{Ni}$  and  $\text{Co}$  solutions which have been boiled with excess of  $\text{KC}_y$ , is due to the fact that  $\text{NiC}_y$  forms with  $\text{KC}_y$  a feeble compound ( $\text{NiC}_y \cdot 2\text{KC}_y$ ) which is soluble in water, but is easily decomposed, whereas  $\text{CoC}_y$  forms with  $\text{KC}_y$  in the air the very stable soluble compound  $\text{K}_6(\text{CoC}_y)_2$  "Potassium Cobalt cyanide."

**136. Blowpipe Reactions.**—Fused with  $\text{Na}_2\text{CO}_3$  on charcoal in the inner flame,  $\text{Co}(\text{NO}_3)_2$  gives a gray metallic powder attracted by the magnet (33 a).

**137. Borax Bead.**—Strong solution may be employed, the bead being dipped into it. *Fine blue* bead in both inner and outer flames, opaque if too much  $\text{Co}(\text{NO}_3)_2$  has been used.

138. GROUP III B. — TABLE OF DIFFERENCES.

Tests.	Zn—Salts.	Mn—Salts.	Ni—Salts.	Co—Salts.
<p><i>a. For liquids.</i></p> <p>1. Add <math>\text{AmS}_2</math> or better <math>\text{H}_2\text{S}</math> and a few drops of <math>\text{AmHO}</math>.</p> <p>2. Add <math>\text{KHO}</math>.</p> <p>3. Add <math>\text{KCY}</math>. [<i>Note.</i>—This test being employed to distinguish between Ni and Co need only be tried when a <i>black</i> precipitate has been obtained by Test No. 1.]</p>	<p><i>White precipitate:</i> soluble in cold dilute <math>\text{HCl}</math>; insoluble in <math>\text{H}_2\text{A}</math>.</p> <p><i>White precipitate:</i> soluble in excess of <math>\text{KHO}</math>.</p> <p>—</p>	<p><i>Pink precipitate:</i> soluble in cold dilute <math>\text{HCl}</math>; <i>soluble</i> in <math>\text{H}_2\text{A}</math>.</p> <p><i>White precipitate:</i> turning brown in the air; <i>insoluble</i> in <math>\text{KHO}</math>.</p> <p>—</p>	<p><i>Black precipitate:</i> soluble in boiling yellow <math>\text{AmS}_2</math> to a dark solution; insoluble in cold dilute <math>\text{HCl}</math> and in <math>\text{H}_2\text{A}</math>.</p> <p><i>Green precipitate:</i> insoluble in <math>\text{KHO}</math>.</p> <p>Added in excess to the slightly acid solution and boiled for some time, <math>\text{Na}_2\text{C}_2\text{O}_4</math> (or <math>\text{Br}</math>-water, and excess of <math>\text{NaHO}</math>) added, gives a black precipitate on heating.</p>	<p><i>Black precipitate:</i> insoluble in boiling yellow <math>\text{AmS}_2</math>* in cold dilute <math>\text{HCl}</math>, and in <math>\text{H}_2\text{A}</math>.</p> <p><i>Blue precipitate:</i> insoluble in <math>\text{KHO}</math>.</p> <p>Added in excess to slightly acid solution and boiled for some time, <math>\text{Na}_2\text{C}_2\text{O}_4</math> (or <math>\text{Br}</math>-water and excess of <math>\text{NaHO}</math>) gives no precipitate on heating.</p>
<p><i>b. Blowpipe tests for solids.</i></p> <p>4. Fused with <math>\text{Na}_2\text{CO}_3</math>.</p>	<p>On charcoal in the inner flame: gives a white incrustation, which, if moistened with <math>\text{Co}(\text{NO}_3)_2</math> and heated in the outer flame, turns <i>green</i>. The substance itself when strongly heated after having been moistened with <math>\text{Co}(\text{NO}_3)_2</math> also becomes <i>green</i>.</p>	<p>On charcoal in the inner flame: a gray magnetic powder (33 a). On platinum foil in outer flame: a <i>bluish-green</i> mass, showing more rapidly if a little <math>\text{KNO}_3</math> be mixed with the <math>\text{Na}_2\text{CO}_3</math> before fusing.</p>	<p>On charcoal in the inner flame, a gray magnetic powder (33 a.)</p>	<p>Same as Ni.</p>
<p>5. Borax bead.</p>	<p>—</p>	<p>Outer flame—<i>Amethyst-red</i>. Inner flame—<i>Colorless</i>.</p>	<p>Outer. { <i>Violet-red</i> or <i>yellow</i> while hot.           <i>Yellow</i> when cold. Inner.—<i>Gray</i> or <i>black opaque</i> bead.</p>	<p>Outer and inner flames—<i>fine blue</i> bead.</p>

\* In order to ascertain whether the precipitate has been dissolved, filter and observe whether the filtrate is dark in color.

**139.** When a solution is to be examined for only one member of this group, its detection by one or more tests in the above table will be very easy. The precipitate with  $\text{Am}_2\text{S}$  is distinctive;  $\text{ZnS}$  and  $\text{MnS}$  being easily recognized by their different colors;  $\text{NiS}$  and  $\text{CoS}$ , which differ from  $\text{ZnS}$  and  $\text{MnS}$  by being black, are distinguished from one another by the solubility of  $\text{NiS}$  in excess of  $\text{Am}_2\text{S}$ , the liquid therefore gives a brown filtrate.  $\text{KHO}$  also gives characteristic precipitates with each of these substances. The most distinctive tests for  $\text{Ni}$  and  $\text{Co}$  are 3 and 5 for  $\text{Zn}$  and  $\text{Mn}$ , 2, 4, and 5.

**140. (s.)** *The method for separating and detecting Zn, Mn, Ni, Co, when mixed, depends on:*

1. The solubility of  $\text{NiS}$  in yellow  $\text{Am}_2\text{S}$ ;  $\text{ZnS}$ ,  $\text{MnS}$ , and  $\text{CoS}$  being insoluble.\*  $\text{NiS}$  is precipitated from this solution by boiling or by addition of  $\text{H}\bar{\text{A}}$ , and its presence confirmed by fusion in a borax bead.

2. The solubility of  $\text{ZnS}$  and  $\text{MnS}$  in cold dilute  $\text{HCl}$ ,  $\text{NiS}$  and  $\text{CoS}$  being almost insoluble. (Note 1.)

3. The solubility of  $\text{ZnHo}_2$  in cold  $\text{KHO}$ , in which  $\text{MnHo}_2$  is insoluble; white  $\text{ZnS}$  is then precipitated from this solution by  $\text{H}_2\text{S}$ .

4. The production of a bluish-green mass by fusing  $\text{MnHo}_2$  with  $\text{Na}_2\text{CO}_3$  and  $\text{KNO}_3$ .

5. The difference of behavior of the  $\text{KCy}$  solution of  $\text{Ni}$  and  $\text{Co}$  with  $\text{Na}_2\text{Cl}_2\text{O}$  (or with  $\text{Br}$ -water and excess of  $\text{NaHO}$ ); and the marked difference in their borax beads. (Note 2.)

*Note 1.*— $\text{Mn}$  may also be separated from  $\text{Ni}$  and  $\text{Co}$  by passing  $\text{H}_2\text{S}$  into the solution containing free  $\text{H}\bar{\text{A}}$ ;  $\text{Mn}$  remains in solution.  $\text{NiS}$  and  $\text{CoS}$  are precipitated.

*Note 2.*—A modification by Henry of Rose's method for separating  $\text{Ni}$  and  $\text{Co}$  is also to be recommended; it depends upon the fact that, whilst  $\text{Co}$  is precipitated as  $\text{Co}_2\text{Ho}_2$  by  $\text{BaCO}_3$  in presence of  $\text{Br}$ ,  $\text{Ni}$  remains in solution. The sulphides are dried and then strongly ignited in an open porcelain crucible; the metals remaining as oxides are dissolved by boiling with a little strong  $\text{HCl}$ ; carbon is filtered off if necessary, and to the clear solution, after being cooled and mixed with  $\text{Br}$ -water, excess of  $\text{BaCO}_3$  is added

---

\* In order to ascertain whether the precipitate has been dissolved, filter and observe whether the filtrate is dark in color.

in a small corked flask; the flask is then well shaken and allowed to stand for about half an hour; on filtering, Co if present is detected in the precipitate by the borax bead; Ni is precipitated from the filtrate by heating it with KHO, filtering and confirming the presence of Ni in the precipitate by fusing a portion of it in the borax bead.

**141.** (s.) A solution is examined for Zn, Mn, Co, Ni, by adding AmCl to it in a boiling tube, then AmHO, if necessary, until it is alkaline; then Am<sub>2</sub>S until, after being well stirred or shaken, the liquid smells of it. The liquid is then boiled, a few drops are poured upon a filter and the color of the filtrate noted; if it is colorless sufficient Am<sub>2</sub>S has not been added, more Am<sub>2</sub>S is then to be poured in, and the liquid again boiled. When a few drops run through the filter brown or yellow in color, Am<sub>2</sub>S has been added in excess, the whole is boiled and filtered, and the precipitate examined by Table III B (437); the filtrate if yellow is rejected, if brown it is examined for Ni according to (431).

---

#### GROUP II A.—COPPER GROUP.

**142.** This group includes Hg'', Pb, Bi, Cu, Cd; its members differ from those of Groups III A, III B, IV, and V, in being precipitated by H<sub>2</sub>S in acid solutions. With the exception of Pb, which in a strong solution is partially precipitated by HCl, the members of this group are not precipitated by HCl.

The sulphides differ from those of Group II B, by being insoluble in Am<sub>2</sub>S or KHO. If very much acid is present, H<sub>2</sub>S does not precipitate the members of this group readily or completely, they are precipitated however on dilution.



---

MERCURICUM (—''Hg'').—Use HgCl<sub>2</sub>.


**143.** Hg forms two series of compounds distinguished as *Mercuric* and *Mercurous*; the mercury in these two states behaves differently with reagents, and may be distinctively called *Mercuricum* and *Mercuriosum*; in the latter state it is classed under Group I.

**144.**  $H_2S$ :\* black precipitate ( $HgS$ ); if the  $H_2S$  solution is added slowly, the precipitate is first *white*, then *brown* or *orange*, and ultimately becomes *black*; these changes of color during the addition of  $H_2S$  are characteristic of mercuric salts. Filter and wash the precipitate; place portions of it in three test-tubes; boil one portion with strong  $HNO_3$ , and another with strong  $HCl$ ; the precipitate is not dissolved by either acid; mix the contents of the two tubes and warm again, the precipitate will now dissolve; boil the third portion with  $Am_2S$ ; the precipitate does not dissolve. Hence  $HgS$  is insoluble in hot strong  $HNO_3$ , in hot strong  $HCl$ , and in  $Am_2S$ , but is easily soluble in a mixture of  $HNO_3$  and  $HCl$  ("aqua regia").

**145.**  $SnCl_2$ : white precipitate ( $Hg_2Cl_2$ ); if more  $SnCl_2$  is added and the liquid boiled, the white precipitate becomes gray and consists of fine particles of  $Hg$ ; if the liquid be decanted and the gray precipitate be then boiled with strong  $HCl$ , the  $Hg$  particles unite to globules visible by a lens or frequently to the naked eye.



**146.** *Cu*: If a small strip of copper sheet or a copper coin, whose surface has been polished and cleansed by rubbing it with sand-paper, be immersed in some  $HgCl_2$  solution made acid with a drop or two of  $HCl$ , it will soon become coated with a gray film of  $Hg$ :



If the surface of the *Cu*, after having been immersed for several minutes, is dried and rubbed hard with a cloth or piece of wash-leather, it will appear more or less whitened like silver, the liquid metal  $Hg$  having "amalgamated" the *Cu*. The  $Hg$  may be readily driven off by heating the *Cu* strongly; if this heating is per-

---

\* A test-tube half full of the solution must be used in order to get sufficient precipitate for the experiments below. This remark applies also to the liquid used for obtaining the  $H_2S$  precipitate with the other members of this group.



formed in a test-tube, or better in a small hard-glass tube sealed at one end (10), globules of Hg are sublimed upon the cool sides of the tube.

147. Mix *a little* solid  $\text{HgCl}_2$  intimately with about three times as much  $\text{Na}_2\text{CO}_3$ , by powdering them together in a mortar, or with a pestle on a watch-glass. Pour some of this dried mixture into the bottom of a piece of perfectly dry hard-glass tubing closed by being drawn out (10) at one end (see note below), and cover the mixture with a small layer of  $\text{Na}_2\text{CO}_3$ ; now heat strongly first the layer of  $\text{Na}_2\text{CO}_3$ , then gradually the mixture; Hg will sublime, condensing in globules on the sides of the tube. Sometimes the globules are so small that the layer appears as a gray film; they may be united into visible globules by rubbing the film with a splinter of wood or with a thin glass rod (11 *a*), or the separate globules may be seen by the use of a pocket lens.

*Note.*—The powdered mixture is best dried by spreading it out upon a watch-glass and placing it in a steam or water-oven; it may be dried also by supporting it on the filter dryer (25) at some height above a *small* flame. A powder such as the above is most easily placed in a narrow closed tube by alternately scooping the powder up with the mouth of the tube, and tapping the tube so as to shake the powder to the bottom; or by turning the powder out upon a piece of note-paper folded over into a trough and pouring it thence into the tube, holding both over the watch-glass to catch any substance which falls. Both the inside of the tube and the powder must be perfectly dry, else some powder obstinately adheres to the sides of the tube and obscures the Hg sublimate. The heating must never be commenced until the sides of the tube are perfectly cleansed if necessary with a twisted slip of filter-paper or a wooden match; also if any drops of water condense on the inside of the tube during the first stage of the heating they must be removed by a twisted piece of filter-paper, or by a small piece of filter-paper rolled round a wooden match or a thin piece of wire.

148. Heat a little solid  $\text{HgCl}_2$  in a tube closed at one end; the substance sublimes in white fumes, since Hg compounds are volatile.

LEAD (Pb).—Use  $\text{PbA}_2 \cdot 3\text{H}_2\text{O}$  dissolved in water to which a little HA has been added.

149.  $\text{H}_2\text{S}$ : black precipitate ( $\text{PbS}$ ); this precipitate is

red if much hydrochloric acid is present in the solution, but becomes black on diluting with water and passing  $H_2S$ , or on adding more  $H_2S$  solution. Filter or decant, and show with separate portions of the precipitate that  $PbS$  is insoluble in  $KHO$  or  $Am_2S$ , soluble in boiling dilute  $HNO_3$ , but converted by boiling strong  $HNO_3$  into insoluble  $PbSO_4$ .

150.  $HCl$ : white precipitate ( $PbCl_2$ ), forms only in cold and strong solutions; on boiling, the precipitate dissolves,\* but the  $PbCl_2$  separates again in beautiful crystals on cooling.

151.  $H_2SO_4$ : white precipitate ( $PbSO_4$ ); this precipitate is much less soluble in dilute  $H_2SO_4$  than in water, hence  $H_2SO_4$  should be added in excess to a pretty strong solution of  $Pb$ ; pour off into two test-tubes and let the liquid stand; decant the liquid from the precipitate, and show that the precipitate may be dissolved by pouring upon it  $H\bar{A}$  or  $H_2\bar{T}$ , then  $AmHO$  in excess, and boiling; show also that it is dissolves in boiling strong  $HCl$ .

152.  $K_2CrO_4$  (or  $K_2Cr_2O_7$ ); yellow precipitate ( $PbCrO_4$ ); pour off into two tubes and show that the precipitate is soluble in  $KHO$ , but insoluble in  $H\bar{A}$ .


153. *Blowpipe Reaction*.—Mix well some powdered  $Pb\bar{A}_2 \cdot 3H_2O$  with about twice as much  $Na_2CO_3$ ; heat the mixture in a cavity on a piece of charcoal in the inner flame; a yellow incrustation will form around the cavity and small bright white globules of  $Pb$  will be seen within it. The incrustation is readily removed when heated in the inner blowpipe flame, coloring the flame azure-blue. Detach one of the globules with the point of a knife, place it on the bottom of a mortar turned upside down and give it a smart blow with the pestle; it flattens out without breaking into powder, because  $Pb$  is *malleable* not *brittle*. If one of the globules be fixed upon the point of a penknife, it will be found by gentle friction on paper to mark it as a black lead-pencil does.

---

\* If the precipitate does not entirely dissolve, add a little water and boil again.

**BISMUTH (Bi).**—Use  $\text{Bi}(\text{NO}_3)_3 \cdot 5\text{H}_2\text{O}$  dissolved in dilute  $\text{HCl}$ .

**154.**  $\text{H}_2\text{S}$ : brownish-black precipitate ( $\text{Bi}_2\text{S}_3$ ); let stand, and decant most of the water; then pour some of the precipitate into three test-tubes, and show that it is insoluble in  $\text{Am}_2\text{S}$  and in  $\text{KHO}$ , but soluble in strong  $\text{HNO}_3$ .



**155.**  $\text{AmHO}$  (or  $\text{KHO}$ ): white precipitate ( $\text{BiHO}_3$ ); pour off a small quantity of the liquid and precipitate into a tube, add much  $\text{AmHO}$  and warm, the precipitate is undissolved; filter off the rest of the precipitate and dissolve it by pouring upon the filter a few drops of hot dilute  $\text{HCl}$ ; keep this solution.

**156.**  $\text{H}_2\text{O}$ : pour the  $\text{HCl}$  solution obtained from the last reaction, or some of the original solution, into a large quantity of distilled water contained in a beaker; on stirring and letting stand a milkiness appears, due to the formation of  $\text{BiOCl}$ :



Pour a little of the milky liquid into two test-tubes; warm one portion after adding to it a little strong  $\text{HCl}$ , the precipitate dissolves: to the other add  $\text{H}_2\text{T}$  and warm, the milkiness remains (difference from  $\text{SbOCl}$ ). If but little  $\text{Bi}$  is present this milkiness often only appears after stirring and letting the liquid stand for five or ten minutes.

**157.**  $\text{K}_2\text{CrO}_4$  or  $\text{K}_2\text{Cr}_2\text{O}_7$  (see note below): yellow flocculent precipitate [ $\text{Bi}_2(\text{CrO}_4)_3$ ]: add  $\text{KHO}$ , the precipitate does not dissolve (difference from  $\text{PbCrO}_4$ ); it dissolves completely in  $\text{HCl}$  or  $\text{HNO}_3$  added in excess.

*Note.*—Since this precipitate is soluble in  $\text{HCl}$ , and  $\text{HCl}$  is present in the  $\text{Bi}$  solution used, the free  $\text{HCl}$  must first be removed by dissolving in a little of the  $\text{Bi}$  solution a sufficient quantity of solid  $\text{Na}\bar{\text{A}}$ ; the free  $\text{HCl}$  is thus replaced by  $\text{H}\bar{\text{A}}$ , which does not dissolve the precipitate:



**158.**  $\text{H}_2\text{SO}_4$ : no precipitate.

**159.** *Blowpipe Reaction.*—A mixture of powdered  $\text{Bi}$

( $\text{NO}_3$ )<sub>3</sub> and  $\text{Na}_2\text{CO}_3$  heated on charcoal in the inner flame, gives an incrustation *orange-red* whilst hot, *yellow* when cold; also white globules of Bi, which are very brittle, being crushed to powder by a sharp blow with a pestle (153).

---

COPPER (Cu).—Use  $\text{CuSO}_4 \cdot 6\text{H}_2\text{O}$ .

Copper salts are usually either blue or green in color; their color commonly becomes very pale or disappears when the salts are thoroughly freed from water of crystallization.

160.  $\text{H}_2\text{S}$ : brownish-black precipitate ( $\text{CuS}$ ): filter, keeping the funnel carefully covered with a glass plate, as air quickly oxidizes  $\text{CuS}$  to  $\text{CuSO}_4$  which dissolves and runs through with the filtrate. Place some of the precipitate in five test-tubes and show that it is insoluble in  $\text{KHO}$ , in  $\text{Am}_2\text{S}$ , and in boiling dilute  $\text{H}_2\text{SO}_4$ , but dissolves in boiling  $\text{HNO}_3$  (dilute or strong), and in  $\text{KCy}$  solution.

161.  $\text{AmHO}$  added in very small quantity: a *greenish-blue* precipitate; if more  $\text{AmHO}$  is added this precipitate dissolves, yielding an *intensely blue* liquid containing  $(\text{N}_2\text{CuAm}_4\text{H}_2)\text{SO}_4$ , which becomes again light-blue on adding an acid in excess.

162.  $\text{H}_2\text{SO}_4$ : no precipitate.

163.  $\text{K}_4\text{FeCy}_6$ : reddish-brown precipitate ( $\text{Cu}_2\text{FeCy}_6$ ), insoluble in  $\text{HA}$ ; the color is best seen by doing the reaction in a white porcelain dish. In *very* dilute solution only a reddish color is produced.

164.  $\text{Fe}$ : a bright strip of steel or iron, such as the blade of a penknife, freed from grease by rubbing with sand-paper or by boiling in a little dilute  $\text{KHO}$  and washing, when dipped into  $\text{CuSO}_4$  solution made acid with a few drops of  $\text{H}_2\text{SO}_4$ , is covered with a red film of Cu after a time.

165.  $\text{Zn}$  and  $\text{Pt}$ : a strip of bright Zn if placed upon a piece of platinum foil or wire in some  $\text{CuSO}_4$  solution contained in a porcelain dish or watch-glass, causes a red film of Cu to deposit on the platinum: the  $\text{CuSO}_4$  solution should be first made acid with a few drops of  $\text{H}_2\text{SO}_4$ .

**166. Blowpipe Reactions.**—A mixture of powdered  $\text{CuSO}_4 \cdot 6\text{H}_2\text{O}$  and  $\text{Na}_2\text{CO}_3$  if heated on charcoal in the inner blowpipe flame, gives red scales of Cu, which are best seen by separating them as directed in par. 33 a.

**167. Borax bead.**  $\left\{ \begin{array}{l} \text{In outer flame:} \\ \text{In inner flame:} \end{array} \right\} \begin{array}{l} \text{Green whilst hot, blue when cold.} \\ \text{Red or colorless bead when cold, obtained only when very little Cu is present, and the bead is long heated in a good reducing flame.*} \end{array}$

**168. Flame Coloration:** a loop of platinum wire dipped into  $\text{CuSO}_4$  solution, and held in the inner blowpipe flame, or in the Bunsen flame gives a *green* coloration; the flame shows a *blue centre*, after having been moistened with HCl.

CADMIUM (Cd).—Use  $\text{CdSO}_4 \cdot 4\text{H}_2\text{O}$ .

**169.  $\text{H}_2\text{S}$ :** bright yellow precipitate (CdS); boil off the  $\text{H}_2\text{S}$ , and pour into several tubes; show with these separate portions that the precipitate is insoluble in  $\text{Am}_2\text{S}$ , in KHO, and in KCy solution, but soluble in boiling dilute  $\text{HNO}_3$ , and in boiling dilute  $\text{H}_2\text{SO}_4$ : dilute the  $\text{H}_2\text{SO}_4$  solution with much water and pass  $\text{H}_2\text{S}$  for a short time, yellow CdS is again precipitated.

**170.  $\text{AmHO}$**  added in small quantity, best by using  $\text{AmHO}$  much diluted: white precipitate ( $\text{CdHo}_2$ ); on adding more  $\text{AmHO}$  the precipitate readily dissolves.

**171.  $\text{H}_2\text{SO}_4$ :** no precipitate.

**172. Blowpipe Reaction.**—Powdered  $\text{CdSO}_4$ , mixed with  $\text{Na}_2\text{CO}_3$  and heated on charcoal in the inner blowpipe flame, gives a brown incrustation.

**173. GROUP II A.—TABLE OF DIFFERENCES.**—(See opposite page.)

\* The red bead is much more easily and rapidly obtained by fusing a fragment of Sn or Zn into the bead; the former gives a clear, the latter a turbid bead.

GROUP II A.—TABLE OF DIFFERENCES.

Tests.	Hg—Salts.	Pb—Salts.	Bi—Salts.	Cu—Salts.	Cd—Salts.
<i>a. For liquids.</i> 1. Pass $H_2S$ or add $H_2S$ water.	<i>Black</i> precipitate, when filtered and well washed, insoluble in boiling strong $HNO_3$ , and unchanged by it.	<i>Black</i> precipitate, almost insoluble in boiling strong $HNO_3$ , changed by it into white $PbSO_4$ .	<i>Black</i> precipitate; soluble in boiling strong $HNO_3$ , to form a colorless solution.	<i>Black</i> precipitate; soluble in boiling concentrated $HNO_3$ , to form a blue solution; insoluble in boiling dilute $H_2SO_4$ .	<i>Yellow</i> precipitate; soluble in boiling $HNO_3$ , and in boiling dilute $H_2SO_4$ .
2. Add $H_2SO_4$ .	—	White precipitate.	—	—	—
3. Add $AmHO$ .	White precipitate, insoluble in excess.	White precipitate, insoluble in excess.	White precipitate, insoluble in excess.	Blue precipitate easily soluble in excess to a deep blue liquid.	White precipitate, easily soluble in excess.
<i>b. Blowpipe reactions for solids.</i> 4. Borax bead.	—	—	—	Outer flame { <i>Green</i> , hot <i>Blue</i> , cold. Inner flame { <i>Colorless</i> <i>Colorless</i> bead	—
5. Flame coloration.	—	—	—	<i>Green</i> when moistened with $HCl$ shows <i>blue</i> centre.	—
6. Heated with $Na_2CO_3$ . <i>a.</i> In a tube, closed at one end. <i>b.</i> On charcoal in the inner blow-pipe flame.	Sublimate of $Hg$ .  —	Yellow incrustation; white malleable globules.	<i>Orange-red</i> incrustation; white <i>brittle</i> globules.	No incrustation; red metallic scales.	<i>Brown</i> incrustation; no globules.

174. The color of the  $H_2S$  precipitate, and its behavior with strong  $HNO_3$ , enable the analyst to detect one member of this group when present alone: the most characteristic special tests for each member of this group will be found under its reactions.

175. (s.) The separation of Hg, Pb, Bi, Cu, Cd, depends upon the following differences:

1. The solubility of  $Bi_2S_3$ , CuS, CdS in strong boiling  $HNO_3$ ; HgS and PbS being insoluble.

2. The solubility of  $PbSO_4$  in  $H\bar{A}$  and excess of AmHO; HgS being insoluble.

3. The solubility of  $CuHO_2$  and  $CdHO_2$ , and insolubility of  $BiHO_3$ , in excess of AmHO.

4. The solubility of CdS and insolubility of CuS in boiling dilute  $H_2SO_4$ . The insolubility of CdS in KCy solution, in which CuS easily dissolves, furnishes another means of separation.

After having been separated, the presence of each metal is confirmed by some special test.

176. (s.) *A solution which may contain Pb, Hg, Bi, Cu, Cd, mixed*, is first made acid, if not already acid, with a little HCl;\*  $H_2S$ -water is then added, or the gas is passed in a rapid stream through the solution for about five minutes; the precipitate is filtered off, the filtrate mixed with more  $H_2S$ -water, or diluted with a little water and  $H_2S$  again passed for a short time, to ascertain whether all the above metals have been completely precipitated: if no further precipitate is produced, the filtrate may be rejected; but in case  $H_2S$  causes further precipitation more  $H_2S$  solution must be added, or the gas must be passed for some time longer, and the liquid poured again through the filter, the filtrate only being rejected when  $H_2S$  no longer produces any precipitate in it; the precipitate is then examined by Table II (435), commencing at Group II A (Copper Group), and using only the left-hand side of the table.

---

\* A white precipitate of  $PbCl_2$ , which may form if the solution is very strong, being neglected, since it is readily converted into PbS by  $H_2S$ .

## GROUP II B.—ARSENIC GROUP.

**177.** This group includes As, Sb, Sn; its members differ from those of Groups III A, III B, IV, and V by being precipitated by  $H_2S$  in a solution made acid with HCl; they differ from those of Group I, in not being precipitated by HCl.

The sulphides of this group, which are precipitated by  $H_2S$ , differ from those of Group II A, by being soluble in  $Am_2S$  or  $KHO$ .

**178.** Each member of this group forms two series of compounds which resemble each other in many reactions, but also present several differences. The two classes are distinguished conveniently by the terminations *-ic* and *-ous*; thus we speak of *arsenic* and *arsenious* acid.

---

ARSENIC (As).	{	For <i>arsenious</i> compound use HCl solution of $As_2O_3$ .
	{	For <i>Arsenic</i> compound use water solution of $Na_2HAsO_4 \cdot 12H_2O$ .

A. TESTS WHICH GIVE ULTIMATELY THE SAME RESULT WITH BOTH SETS OF COMPOUNDS.

The two solutions should be taken in separate test-tubes, and a portion of each tried in succession with each reagent; the difference of behavior is thus more easily appreciated and remembered.

*Arsenic* compounds are converted into *arsenious* by boiling with HCl; this explains reactions 179 and 180 with *arsenic* solutions.



Reagent.	Arsenious solution.	Arsenic solution ( $\text{Na}_2\text{HAsO}_4 \cdot 12\text{H}_2\text{O}$ ).
<b>179.</b> $\text{H}_2\text{S}$ .	In neutral solutions only a yellow color is produced, but if the solution is acid with $\text{HCl}$ a light yellow flocculent precipitate ( $\text{As}_2\text{S}_3$ ) forms immediately even in the cold. Shake up and pour some of the liquid into another test-tube and show that $\text{KHO}$ and $\text{Am}_2\text{S}$ , added to the separate portions drop by drop, dissolve the precipitate, which, however, is thrown down again on addition of $\text{HCl}$ in excess.*	No precipitate is produced in the cold solution acidified with $\text{HCl}$ ; but if the acid solution is boiled, $\text{H}_2\text{S}$ causes in the hot solution, first a milky precipitate of $\text{S}$ , then yellow $\text{As}_2\text{S}_3$ .
<b>180.</b> $\text{Cu}$ : three or four narrow slips brightly polished with sand-paper.	<i>Reinsch's Test.</i> —If boiled in the solution to which some dilute $\text{HCl}$ has been added, $\text{Cu}$ becomes coated with a black film ( $\text{As}_2\text{Cu}_3$ ); if the $\text{Cu}$ slips are carefully dried by being pressed between filter-paper, and are heated near the lower end of a piece of hard-glass tube open at both ends and 4 and 5 inches in length, which is held somewhat obliquely in the flame or bent as seen in Fig. 85 (p. 132), a white sublimate of $\text{As}_2\text{O}_3$ forms in the upper part of the tube; on being examined with a lens this is seen to consist of white octohedral crystals. This sublimate may be dissolved when cold in a little hot water, and the presence of $\text{As}$ confirmed by tests 179 and 187. The presence of $\text{As}$ in the deposit should always be confirmed in this way, since other metals besides $\text{As}$ are deposited on $\text{Cu}$ as a black film.	The same deposit is obtained as with $\text{As}_2\text{O}_3$ but only when strong $\text{HCl}$ is added and the liquid boiled.


\* The  $\text{KHO}$  solution of the precipitate and the  $\text{As}_2\text{S}_3$  precipitated from it by  $\text{HCl}$  are often discolored by black sulphides formed by the action of  $\text{H}_2\text{S}$  upon  $\text{Pb}$  or  $\text{Fe}$  present as an impurity in the  $\text{KHO}$ : this discoloration may be almost entirely prevented by boiling the liquid before adding  $\text{KHO}$ , the free  $\text{H}_2\text{S}$  is thus expelled.

TESTS DEPENDING ON THE FORMATION OF  $\text{AsH}_3$ .

*Note 1.*—As  $\text{AsH}_3$  is a very poisonous gas, the following experiments with it should be performed in a cupboard provided with a good draught, or in the open air.

*Note 2.*—Arsenic compounds give these reactions less readily than arsenious, but by being boiled with  $\text{HCl}$  for a short time they are converted into arsenious compounds, which then readily answer to the following reactions.

**181. Hofmann's Method.**—If some scraps of  $\text{Zn}$  and some dilute  $\text{HCl}$ , both free from  $\text{As}$ , be placed together in a small flask fitted with a funnel-tube dipping below the liquid and a delivery-tube, as shown in Fig. 33,\*  $\text{H}$  gas will be given off either at once or on adding a little strong  $\text{HCl}$ ;




The hydrogen can be made to bubble through some  $\text{PbA}_2$  solution contained in a second small flask

connected with the first by a piece of india-rubber tubing; the gas, thus freed from any  $\text{H}_2\text{S}$  with which it might possibly have been mixed, is then allowed to bubble through some  $\text{AgNO}_3$  solution contained in a test-tube which is supported in a small beaker; if the  $\text{Zn}$  and acid were free from  $\text{As}$ , no precipitate or color is produced in the  $\text{AgNO}_3$  solution.

If a little  $\text{As}_2\text{O}_3$  solution be now poured into the larger flask through the funnel, a black precipitate ( $\text{Ag}$ ) will


FIG. 33.




\* In order to be sure that the corks and tubes fit air-tight, moisten the outside of the cork after it has been placed in the neck of the flask, and having closed one tube, blow down the other. No air-bubbles must be seen or heard to escape.

Several of these apparatus may be kept ready fitted in the laboratory for general use.

begin to form in the  $\text{AgNO}_3$  solution; this is produced by the  $\text{AsH}_3$  evolved by the action of the H upon the  $\text{As}_2\text{O}_3$ :




The  $\text{AsH}_3$  on passing into the  $\text{AgNO}_3$  solution, reacts thus:



Hence Ag is precipitated, and  $\text{H}_3\text{AsO}_3$  remains in solution.

As soon as a copious precipitate has been obtained the test-tube is removed. It will be best to put by the test-tube with its contents to be examined afterwards as directed below, and whilst the gas is coming off to proceed at once to use it for reactions 182 and 183.

The liquid in the test-tube is filtered, the precipitate rejected, and several drops of  $\text{AgNO}_3$  solution are added to clear the filtrate. On mixing with this liquid drop by drop some very dilute  $\text{AmHO}$  (made by pouring several drops of  $\text{AmHO}$  solution into a test-tube, then nearly filling up with distilled water and shaking the tube) and stirring or shaking after each addition, a light yellow precipitate ( $\text{Ag}_3\text{AsO}_3$  or  $\text{Ag}_2\text{HAsO}_3$ ) will be obtained. The formation of this precipitate on addition of  $\text{AmHO}$ , is explained by the fact that  $\text{H}_3\text{AsO}_3$  is in the solution in the presence of  $\text{AgNO}_3$  and also of  $\text{HNO}_3$  (see last equation); now  $\text{AgNO}_3$  forms with  $\text{H}_3\text{AsO}_3$  a yellow precipitate ( $\text{Ag}_3\text{AsO}_3$ ) if no free acid is present in the liquid (see reaction 187); hence on neutralizing with  $\text{AmHO}$  the free  $\text{HNO}_3$  already present, and that which is formed by the action of  $\text{AgNO}_3$  on  $\text{H}_3\text{AsO}_3$ , the precipitate appears:



This precipitate is also readily soluble in  $\text{AmHO}$ ;

hence great caution is requisite to use very dilute  $\text{AmHO}$ , and to add it gradually.


*Caution.*—Both flasks should be rinsed out each time after use, else some of the  $\text{AsH}_3$  may remain in it and be driven out in a future operation.

182. Disconnect the larger flask from the smaller, and slip into the india-rubber joint a tube of hard glass about four inches in length whose end is drawn out to a fine jet and supported on the ring of the retort-stand, as shown in Fig. 34; if the gas is not coming off briskly pour in through the funnel-tube a little strong  $\text{HCl}$ , and light the gas as it issues from the jet.\* The flame will differ from that of pure  $\text{H}$  by burning with a bluish tinge, and by giving off white fumes of  $\text{As}_2\text{O}_3$ .†

Press down upon the flame the inside of a porcelain crucible lid or of a porcelain crucible or dish, a dusky black film of  $\text{As}$  will be deposited upon the cool surface; pour upon this some  $\text{Na}_2\text{Cl}_2\text{O}$  solution, the stain is rapidly dissolved. Warm also a stain of  $\text{As}$ , obtained inside a porcelain dish with  $\text{Am}_2\text{S}_3$ , it will dissolve and on gentle evaporation will leave *yellow*  $\text{As}_2\text{S}_3$ .

183. Add a little more strong  $\text{HCl}$  if necessary to

Fig. 34.




\* If the experiment is performed as here directed, the  $\text{H}$  will have been evolved for some time before being lighted, and there is no fear of an explosion occurring; but if the  $\text{H}$  is being produced for the first time, allow it to escape briskly for four or five minutes, and ascertain that a test-tube full of the gas burns quietly (see Exp. 8, p. 22) before adding the liquid to be tested for  $\text{As}$  and lighting the gas at the jet.

† If these appearances are not noticeable, pour a little more  $\text{As}_2\text{O}_3$  solution into the flask through the funnel and mix by gentle shaking.

cause the gas to come off rapidly, light it at the jet and heat a point near the middle of the tube strongly (see Fig. 34); a black mirror of As will form just beyond the part heated by the flame, and the flame burning at the jet will meanwhile become colorless, showing that the arsenic has been thus removed from the hydrogen. Heat another point in the tube, so as to produce a second mirror; then divide the tube between the mirrors, and show by warming one portion when cold with some  $\text{Na}_2\text{Cl}_2\text{O}$ ; solution in a test-tube, that the mirror is soluble in  $\text{Na}_2\text{Cl}_2\text{O}$ ; show that the other, when heated in the tube held obliquely in the flame, or bent as shown in Fig. 35, gives a sublimate of  $\text{As}_2\text{O}_3$ . Dissolve the sublimate when cool in a little hot water, and confirm the presence of As by reactions 179 and 187.


FIG. 35.



*Note.*—Tests 182 and 183 were first suggested by *Marsh*; the former is usually known as *Marsh's test*.

**183 a.**  $\text{AsH}_3$  produces a characteristic color with  $\text{HgCl}_2$  solution. Remove the cork from the larger flask and place a stopper of cotton-wool in its neck to stop any spirting from the liquid in the flask, then cover the mouth of the flask with a small piece of filter-paper moistened with  $\text{HgCl}_2$  solution. The paper becomes colored first yellow then brown.

**184.** If some pieces of Zn or Al be boiled in  $\text{KHO}$  solution,  $\text{H}$ -gas is evolved which is free from odor, but if a little  $\text{As}_2\text{O}_3$  solution be now added and the liquid be again boiled, a garlic odor is evolved owing to the formation of  $\text{AsH}_3$ :



If a small slip of filter-paper moistened with  $\text{AgNO}_3$  solution and placed on the end of a glass rod,\* be held in the gas in the mouth of the tube whilst the liquid is being boiled, the paper will be stained black by the separation of Ag (see reaction 181).

This reaction does not succeed with arsenic compounds unless they are first reduced by boiling with  $\text{HCl}$ . It is not given by any Sb compounds (difference between As and Sb).

---

\* A simple method of getting a moistened piece of paper on the rod is to dip the rod into  $\text{AgNO}_3$  solution; then gently press the paper slip upon it whilst the rod is wet.

**185.** If some solid  $\text{As}_2\text{O}_3$  or any As compound, be mixed with powdered  $\text{Na}_2\text{CO}_3$  and  $\text{KCy}$  (or with

FIG. 36.



$\text{K}_2\text{C}_2\text{O}_4$ ) in a mortar, and a small quantity of the mixture, after having been dried for some time at a gentle heat on a watch-glass, be introduced into a little bulb-tube of hard glass (Fig. 36) at least three inches in length; and if the substance be then heated, observing the precautions stated in the note under (147), a black mirror of As forms in the cooler part of the tube and frequently a smell of garlic may be detected at the mouth of the tube. If the bulb be cut off and the mirror be heated by holding the tube obliquely in the flame (Fig. 35), the mirror is converted into a sublimate of  $\text{As}_2\text{O}_3$  crystals.

**186. Blowpipe Reaction.**—A mixture of any solid As compound with  $\text{Na}_2\text{CO}_3$  if heated on charcoal in the inner blowpipe flame emits a smell of garlic, which is noticed if the charcoal is removed from the flame and smelt. The flame is colored *livid blue*.

A little solid  $\text{As}_2\text{O}_3$  heated in a small hard-glass tube closed at one end is readily “sublimed,” and the sublimate is seen under a microscope or a powerful lens to consist of octohedral crystals.

## B. DISTINCTIVE TESTS FOR ARSENIOUS AND ARSENIC COMPOUNDS.

Reagent.	Arsenious compound. Use $As_2O_3$ dissolved in HCl except for (187).	Arsenic compound. Use $Na_2HAsO_4 \cdot 12H_2O$ dissolved in $H_2O$ .
<b>187. <math>AgNO_3</math>,</b> several drops:  <i>Note.</i> —For this reaction $As_2O_3$ dissolved in boiling water is to be employed, since HCl would yield a precipitate of AgCl.	<i>Light-yellow precipitate.</i> $(Ag_2HASO_3 \text{ or } Ag_3AsO_3)$ This precipitate forms only on adding carefully, drop by drop, a little dilute $AmHO$ . <sup>*</sup> Show that it is dissolved by $AmHO$ and by $HNO_3$ . It is also appreciably soluble in $AmNO_3$ ; hence, if too much $AmHO$ has been added, the precipitate does not always form on neutralizing with $HNO_3$ .	<i>Brown precipitate.</i> $(Ag_3AsO_4)$ : show that it is soluble in $AmHO$ and in $HNO_3$ .
<b>188. <math>CuSO_4</math>,</b> several drops:	<i>Yellowish-green precipitate</i> $(Cu''HASO_3)$ . This precipitate forms only on adding dilute $AmHO$ carefully, drop by drop:† show that it is soluble in $AmHO$ and in $HNO_3$ .	<i>Pale-green precipitate,</i> $(CuHASO_4)$ : soluble in $AmHO$ and in $HNO_3$ .
<b>189. <math>CuSO_4</math>,</b> two drops: then $KHO$ added.	The above yellowish-green precipitate is first produced, but on addition of more $KHO$ this dissolves to a clear blue liquid, which on being boiled deposits red $Cu_2O$ .	No clear blue solution is obtained, and no $Cu_2O$ produced on boiling the liquid.
<b>190. <math>AmCl</math>,</b> <i><math>AmHO</math>, and</i> <i><math>MgSO_4</math>.</i>	No precipitate.	White crystalline precipitate $(MgAmAsO_4)$ : if filtered off and moistened with $AgNO_3$ solution on the filter becomes brown.

\*  $As_2O_3$  when dissolved in water yields  $H_3AsO_3$ , which cannot give with  $AgNO_3$  the  $Ag_3AsO_3$  precipitate, because this decomposition would liberate  $HNO_3$  in which  $Ag_3AsO_3$  is soluble; hence the addition of  $AmHO$  is necessary to neutralize the free acid; but since the precipitate is also soluble in  $AmHO$ , the latter must be added carefully in only just sufficient quantity to neutralize the acid. This is best effected by using some very dilute  $AmHO$ , made by pouring a few drops into a test-tube, filling the tube two-thirds with distilled water, then closing it with the thumb and inverting it for a moment. This dilute  $AmHO$  is to be added drop by drop, shaking or stirring the test solution after each addition until the precipitate is obtained.

† The precipitate produced by  $CuSO_4$  is soluble in acids; hence the HCl present in the solution must be neutralized by careful addition of  $AmHO$  before the precipitate will appear.

ANTIMONY (Sb).—Use  $\text{SbCl}_3$ , or  $2(\text{KSbOT})\cdot\text{H}_2\text{O}$  ("Tartar Emetic"), dissolved in dilute  $\text{HCl}$ .

*Note.*—Antimonious compounds give results ultimately identical with antimonious in the following eight reactions, except in (192). After these reactions, which are so common to Sb in both its sets of compounds, are given two which are distinctive: they are, however, rarely used and may be neglected by the student.

191.  $\text{H}_2\text{S}$ : orange-red precipitate ( $\text{Sb}_2\text{S}_3$ ): pour into two tubes and show that the precipitate is dissolved when warmed with pure  $\text{NaHO}$  or with  $\text{Am}_2\text{S}$ , but is reprecipitated from these solutions on addition of  $\text{HCl}$  in excess.


192. If poured into much water,  $\text{SbCl}_3$  solution yields a white precipitate or turbidity ( $\text{SbOCl}$ ): this precipitate redissolves on adding  $\text{HCl}$  and warming, and is also readily soluble in  $\text{H}_2\text{T}$ ; in this latter respect it differs from the similarly formed  $\text{BiOCl}$  precipitate (156).

193. If a few drops of acid  $\text{SbCl}_3$  solution be poured upon a piece of platinum foil\* and a piece of  $\text{Zn}$  be dropped into it, a black deposit of  $\text{Sb}$  will rapidly form on the foil. If the foil be rinsed with water and then boiled with  $\text{HCl}$ , the stain remains undissolved, but it is rapidly removed by boiling  $\text{HNO}_3$ .

#### TESTS DEPENDING ON THE FORMATION OF $\text{SbH}_3$ .

$\text{SbH}_3$  differs from  $\text{AsH}_3$  in not being poisonous, and in being free from smell.

194. If  $\text{SbH}_3$  gas formed by the action of acid  $\text{SbCl}_3$  solution on  $\text{Zn}$ , is passed into  $\text{AgNO}_3$  solution in a precisely similar way to that described for  $\text{AsH}_3$  (181), a black precipitate forms ( $\text{SbAg}_3$ ):



This precipitate is to be filtered off, and the filtrate, which will contain no  $\text{Sb}$ , rejected. Wash the precipitate four or five times with boiling water upon the filter:

---

\* A slip of platinum foil laid in a watch-glass or porcelain dish, or with turned-up edges, may be employed: the inverted lid of a platinum crucible is very convenient for this purpose.



then pour upon it boiling dilute  $H_2T$  solution, which will dissolve the Sb, and receive the liquid in a test-tube; boil it, and pour it once more upon the filter; add HCl to the liquid, filter if necessary, and pass  $H_2S$  into it, orange-red  $Sb_2S_3$  will be precipitated.

195. A stain produced by Sb on porcelain, in the manner described under (182), differs from the As stain in presenting a *dead* surface and in being insoluble in  $Na_2Cl_2O$ ; also when the stain is dissolved by warming with  $Am_2S$ , the solution if gently evaporated leaves *orange-red*  $Sb_2S_3$ .

196. The Sb mirror, obtained as described under (183), differs from the As mirror in being formed much nearer to the flame, and on both sides of the heated part of the tube; it may further be distinguished by the tests given under (195). This mirror, after oxidation, is also insoluble in boiling water: and if dissolved in a little boiling HCl the solution gives an orange-red precipitate ( $Sb_2S_3$ ) when  $H_2S$  is passed into it, whereas the  $As_2O_3$  solution gives yellow  $As_2S_3$ .

196a.  $SbH_3$ , if allowed to act on filter-paper moistened with  $HgCl_2$  solution, as described under 183a for  $AsH_3$ , yields a *grayish-brown* stain.

197. Boiled with Zn or Al and KHO, no  $SbH_3$  is evolved. [Difference from As; see 184.]

198. If a little solid Sb compound, either  $KSbOT$  or  $Sb_2O_3$ , be mixed in a cavity on wood charcoal with  $Na_2CO_3$  and KCy, and the mixture be heated in the inner blowpipe flame, a white incrustation forms on the charcoal, and white globules of metallic Sb are obtained, which are extremely brittle. The flame is colored pale green. If the melted Sb be allowed to stand aside on the charcoal, the globule emits a white smoke, and coats itself with sharp white crystals of the oxide.

The two distinctive tests (199, 200) for antimonious and antimonic compounds given in the following table are seldom used; they may be read through without trying the reactions, and may be referred to hereafter if required for analytical purposes.

## DISTINCTIVE TESTS FOR ANTIMONIOUS AND ANTIMONIC COMPOUNDS.

Reagent.	Antimonious. (Use the above solution of $\text{SbCl}_3$ in dilute $\text{HCl}$ .)	Antimonic. (Use $\text{K}_3\text{SbO}_7$ solution obtained by boiling some "Potassium Metantimonate" with $\text{H}_2\text{O}$ .)
199. Add excess of $\text{KHO}$ , then $\text{AgNO}_3$ solution.	A dark-colored precipitate, which when shaken after addition of $\text{AmHO}$ leaves black $\text{Ag}_2\text{O}$ undissolved.	A brown precipitate, which dissolves entirely on addition of $\text{AmHO}$ .
200. Add excess of $\text{HCl}$ and warm; then pour in a little $\text{KI}$ solution.	No iodine is set free; proved by the liquid not turning brown and giving no color after being cooled and mixed with freshly-prepared starch solution.	Iodine is liberated, giving a brown color to the liquid and yielding a deep blue color when starch solution is added to the cold liquid.

TIN ( $\text{Sn}$ ).—Stannosum and Stannicum.

## DISTINCTIVE REACTIONS FOR STANNOUS AND STANNIC COMPOUNDS.

Test.	Stannous Compounds. Use $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ dissolved in dilute $\text{HCl}$ .	Stannic Compounds. Use $\text{SnCl}_4$ in dilute $\text{HCl}$ .
201. Pass $\text{H}_2\text{S}$ .	<i>Dark-brown precipitate</i> ( $\text{SnS}$ ): pour off two portions and show that the precipitate is soluble in $\text{KHO}$ and in <i>yellow</i> $\text{Am}_2\text{S}$ on heating; and is precipitated again by $\text{HCl}$ , from the $\text{KHO}$ solution as brown $\text{SnS}$ , from the $\text{Am}_2\text{S}$ solution as yellow $\text{SnS}_2$ . $\text{SnS}$ differs from $\text{SnS}_2$ in being insoluble in <i>colorless</i> $\text{Am}_2\text{S}$ , which readily dissolves $\text{SnS}_2$ .	<i>Yellow precipitate</i> ( $\text{SnS}_2$ ), often forming only when the liquid is heated: pour off portions and show that the precipitate is soluble in $\text{Am}_2\text{S}$ (both <i>yellow</i> and <i>colorless</i> ) and in $\text{KHO}$ on heating, and is precipitated again by $\text{HCl}$ as yellow $\text{SnS}_2$ from both solutions.
202. $\text{HgCl}_2$ .	<i>A white precipitate</i> ( $\text{Hg}_2\text{Cl}_2$ ): turns gray on being boiled if the $\text{SnCl}_2$ is in excess, see (145).	—
203. $\text{AuCl}_3$ : added after a few drops of $\text{SnCl}_2$ solution or of $\text{Cl}$ water.	<i>Purple or dark-brown precipitate</i> ("Purple of Cassius.")	—

*Reactions Common to Stannous and Stannic Compounds.*

**204.** Zn in presence of HCl precipitates from Sn solutions a spongy mass of Sn: the Zn and solution should be allowed to stand for some time in a small porcelain dish; the action is much hastened by gentle heat. If this test is done on platinum foil, the Sn is deposited in a spongy state *on the Zn, and does not stain the Pt* (difference from Sb).  $\text{SnCl}_4$  is not so readily acted upon by Zn as  $\text{SnCl}_2$  is; from  $\text{SnCl}_2$  the Sn is often precipitated in beautiful crystals.

**205.** A solid Sn compound,  $\text{SnCl}_2$  or  $\text{SnO}_2$ , mixed with powdered  $\text{Na}_2\text{CO}_3$  and KCy and heated on charcoal in the inner blowpipe flame gives a slight white incrustation and white particles of metallic Sn which are with difficulty fused into globules. The globules of Sn differ from those of Pb by not marking paper, see (153).

## 206. GROUP II B.—TABLE OF DIFFERENCES.

The following tests answer for these elements in both sets of compounds: for distinctive tests see the preceding reactions.

Tests.	As.	Sb.	Sn.
<i>a. For liquids.</i>			
1. Pass $H_2S$ into the solution acidified with HCl and heat: if no precipitate forms heat to boiling, and again pass $H_2S$ .	Yellow precipitate, insoluble in boiling strong HCl.	Orange-red precipitate, soluble in boiling strong HCl.	$\left\{ \begin{array}{l} SnS - \text{Brown precipitate.} \\ SnS_2 - \text{Yellow precipitate.} \end{array} \right.$ Both soluble in boiling strong HCl.
2. Zn and HCl.	$AsH_3$ gas is evolved, which if passed into $AgNO_3$ solution gives a black precipitate of $Ag_3AsO_3$ , being left in solution; on adding dilute $AmHO$ to this solution, yellow $As_2AsO_3$ precipitates.	$SbH_3$ gas is evolved, which if passed into $AgNO_3$ solution gives a precipitate of $Ag_3Sb$ , from which hot $H_2T$ solution dissolves Sb; $H_2S$ gives in this solution, after adding HCl, orange-red $Sb_2S_3$ .	Sn is deposited on the Zn strips.
3. Zn and HCl on platinum.	$AsH_3$ evolved, no stain on the platinum.	Black stain of Sb on the platinum.	Sn deposited on the zinc, no stain on the platinum.
<i>b. For solids.</i>			
4. Heated with $Na_2CO_3$ and KCy on charcoal in the inner blowpipe flame.	No metallic globules; smell of garlic.	Brittle metallic globules, white incrustation.	Malleable metallic globules, white incrustation.
5. Heated with solid $Na_2CO_3$ and KCy in a bulb tube.	Black mirror and smell of garlic.	—	—

The detection of any one member of this group is simple; the color of the precipitate with  $H_2S$ , confirmed by special tests, is quite sufficient to identify it with ease.

207. (s.) *Many methods have been proposed for the separation and detection of As, Sb, and Sn, when mixed; two of the most trustworthy are those introduced by*

*Hofmann* (208) and by *Fresenius* (210); a method more simple, though less trustworthy, is given in (209).

208. (s.) *Hofmann's* method consists in pouring the solution into a flask, in which H is being generated by the action of HCl upon Zn; Sn remains as a spongy mass on the Zn, As and Sb are evolved as  $\text{AsH}_3$  and  $\text{SbH}_3$ ; these gases, if passed into  $\text{AgNO}_3$  solution, yield soluble  $\text{H}_3\text{AsO}_3$  and insoluble  $\text{Ag}_3\text{Sb}$ , which are separated by filtration; the three members of this group thus separated are then detected by special tests.

As, Sb, Sn, occurring together in a solution, are tested for by precipitating them with  $\text{H}_2\text{S}$ , filtering, and testing the filtrate by passing  $\text{H}_2\text{S}$  through it as was directed for precipitating Group II A. The precipitate is then examined by Table II (435 B, Arsenic Group), using only the right-hand side of the table.

The  $\text{H}_2\text{S}$  precipitate may also be examined by the simpler method given in (209), or by the very accurate, but less simple, method in (210).

209. (s.) The following method of examining a precipitate, produced by  $\text{H}_2\text{S}$ , for As, Sb, Sn, is recommended by its comparative simplicity; it is sufficiently accurate for ordinary analyses where traces of these metals have not to be looked for. It depends upon the insolubility of  $\text{As}_2\text{S}_3$  in hot strong HCl, in which  $\text{SnS}$ ,  $\text{SnS}_2$ , and  $\text{Sb}_2\text{S}_3$  are soluble; and the further separation of Sb and Sn is effected either by adding Zn alone to the acid solution, when Sb is evolved as  $\text{SbH}_3$ , and Sn is deposited upon the Zn, or by adding Zn and Pt, when Sn is deposited on the Zn and Sb on the Pt.

The precipitate produced by  $\text{H}_2\text{S}$  is filtered and allowed to drain for some time upon the filter in the funnel; it is best to drain it still further by carefully taking the filter out of the funnel, opening it out, and spreading it upon a piece of filter-paper folded several times upon itself; the precipitate is then removed to a small porcelain dish and heated for some time nearly to boiling with strong fuming HCl; the liquid is cooled and filtered:

The *Residue* will consist chiefly of  $As_2S_3$ , which is almost insoluble in strong  $HCl$ .

Dry the washed residue on the filter at a gentle heat, then mix it with three or four times as much powdered  $KCy$  and  $Na_2CO_3$ , and heat the mixture in a small bulb-tube, removing any drops of water inside the tube by a twisted piece of filter-paper; a black mirror (185):

*Presence of As.*

*Note.*—The  $As_2S_3$  may also be dissolved by heating with a little fuming  $HNO_3$ , boiling off excess of  $HNO_3$  and detecting the  $As$  as  $H_3AsO_4$  by adding  $AmCl$  excess of  $AmHO$  and  $MgSO_4$ ; or by addition of  $AgNO_3$ , and then cautiously neutralizing with  $AmHO$ .

The *Filtrate* may be examined for  $Sb$  and  $Sn$  by either I or II below:

I. Place a piece of platinum foil in a porcelain dish and pour the acid filtrate upon it, then touch the foil with a piece of  $Zn$ ;  $H$  will come off with effervescence, and if either at once or after a few minutes a black stain appears upon the platinum the *Presence of Sb* is indicated.\*

The platinum foil is removed and pieces of  $Zn$  placed into the liquid: as soon as the bubbles of  $H$  cease to be given off,  $Zn$  still remaining undissolved, remove the pieces of  $Zn$ , rubbing and rinsing any dark deposit back into the dish; let this deposit settle, decant the liquid and heat the solid deposit with strong  $HCl$  for several minutes in a test-tube, dilute with a little water, filter if necessary, and add to the solution a few drops of  $HgCl_2$  solution. A white or gray precipitate (204, 202) indicates the *Presence of Sn*.

II. The acid filtrate is poured into a little hydrogen flask (Fig. 84, p. 121), in which  $H$  has been coming off briskly for about five minutes, being produced by the action of a little strong  $HCl$  upon some pieces of  $Zn$ . The  $H$  is lighted at the jet, and the inside of a small porcelain dish or crucible lid pressed down upon the flame: a black stain not dissolved by hot  $Na_2Cl_2O$  solution shows the *Presence of Sb*.

The residue in the flask is tested for  $Sn$ , as directed in the latter part of I (above).

210. (s.) The following method of detecting  $As$ ,  $Sb$ ,  $Sn$ , described by Fresenius, is very delicate and trustworthy.

It depends upon the fact that fusion with  $Na_2CO_3$  and  $NaNO_3$  converts  $As_2S_3$  into soluble  $Na_2HAsO_4$ ,  $Sb_2S_3$  into

\* The  $Sb$  thus deposited on the  $Pt$  may, after rinsing the foil, be dissolved by heating the  $Pt$  in a test-tube with a very dilute  $HNO_3$ ; on cooling, diluting, and passing  $H_2S$ , an orange-red precipitate will form, confirming the *Presence of Sb*.

insoluble  $\text{Na}_2\text{H}_2\text{Sb}_2\text{O}_7$ , and  $\text{SnS}_2$  into insoluble Sn or  $\text{SnO}_2$ .

The precipitated sulphides are dried on the filter, and the precipitate\* mixed well on a watch-glass or in an agate mortar with equal weights of powdered  $\text{Na}_2\text{CO}_3$  and  $\text{NaNO}_3$ ; this powder is dropped in small quantities at a time into some  $\text{NaNO}_3$  kept melted in a small porcelain crucible over a Bunsen flame. When the powder has all been added, the crucible is heated sufficiently to keep the substance melted for a few minutes, and the melted mass is then poured into a small dry porcelain dish; as soon as the substance has cooled, cold water is poured upon it in the dish and also upon the residue adhering to the inside of the crucible, and the solid substance is allowed to soak for some time with occasional stirring, crushing it by pressure with a pestle if it does not easily fall to pieces; it is then filtered:

The *Residue* is washed on the filter with a mixture of equal measures of water and alcohol; the precipitate is then rinsed into a small porcelain dish, using as little water as possible; a few drops of strong  $\text{HCl}$  are added, and the dish is heated. Whether the residue has dissolved or not, place in the liquid a piece of clean platinum foil, and upon the foil a piece of pure Zn. If a black stain forms after a time upon the platinum, it proves the *Presence of Sb*.† When the Zn ceases to be acted upon, Sn, if present, will remain as a spongy deposit. This is to be washed by decantation, dissolved by boiling with  $\text{HCl}$  in a test-tube, and  $\text{HgCl}_2$  added immediately to the solution; a white precipitate (202) shows the *Presence of Sn*. Test 203 may be substituted for 202.

*Filtrate*:  $\text{HNO}_3$  is added until the solution is just acid after being boiled; then  $\text{AgNO}_3$  solution is poured in as long as it causes any precipitate, and very dilute  $\text{AmHO}$  is added little by little;‡ a brown precipitate readily dissolving in excess of  $\text{AmHO}$  shows the *Presence of As*.

\* If the quantity of the precipitate is very small, the part of the paper containing the precipitate is cut up into small pieces and treated as the precipitate.

† See foot-note on p. 141.

‡ Instead of mixing the dilute  $\text{AmHO}$  with the liquid, it may be carefully poured upon the top of it; a brown color is then seen at the surface of contact.

## GROUP I.—SILVER GROUP.

**211.** This group includes Pb, Ag, Hg' (Mercurorum); its members differ from those of all the other analytical groups by being precipitated as chlorides by HCl: since, however,  $\text{PbCl}_2$  is rather soluble in water, Pb is not entirely precipitated by HCl, and is usually found also in Group II A as  $\text{PbS}$ .

---

**LEAD (Pb).**—Reactions already given in Group II A (149-153).

---

**SILVER (Ag).**—Use  $\text{AgNO}_3$  solution.

**212.** *HCl*: white precipitate ( $\text{AgCl}$ ), becomes curdy on being shaken or heated: pour off into four tubes; filter one, wash the precipitate and let it stand for some time exposed to sunlight or common daylight, it will become blackish-purple; show with the other portions that  $\text{AgCl}$  dissolves readily when heated with  $\text{AmHO}$  or with  $\text{KCy}$  solution, and is precipitated again from these solutions on adding  $\text{HNO}_3$  in excess; also that it is insoluble in  $\text{HNO}_3$  even on boiling.

**213.**  $\text{H}_2\text{S}$  (or  $\text{Am}_2\text{S}$ ): black precipitate ( $\text{Ag}_2\text{S}$ ): insoluble in  $\text{Am}_2\text{S}$  or  $\text{KHO}$ ; soluble in boiling dilute  $\text{HNO}_3$ .

**214.** *KHO*: brown precipitate ( $\text{AgHO}$ ): insoluble in excess.

**215.** *AmHO*: light-colored precipitate, produced only when *very dilute*  $\text{AmHO}$  is added drop by drop; easily soluble in excess.

**216.** A mixture of a solid Ag compound and powdered  $\text{Na}_2\text{CO}_3$  heated on charcoal, gives white malleable globules or scales of Ag and no incrustation.

---

**MERCUROSUM (Hg').**—Use  $\text{Hg}'_2(\text{NO}_3)_2 \cdot \text{H}_2\text{O}$  dissolved in dilute  $\text{HNO}_3$ .

*Note.*—Reactions 145, 146, 147, and 148 given under mercuricum ( $\text{Hg}''$ ) yield precisely similar results with Mercurorum ( $\text{Hg}'$ ) and may be repeated with the  $\text{Hg}'_2(\text{NO}_3)_2$  solution.

In reaction 146 however it is unnecessary to add acid since the  $\text{Hg}'_2(\text{NO}_3)_2$  solution is already acid.

**217.** *HCl*: a white precipitate ( $\text{Hg}'_2\text{Cl}_2$ ): insoluble in dilute acids unless warmed with both  $\text{HCl}$  and  $\text{HNO}_3$ ,



144 GROUP I.—TABLE OF DIFFERENCES. [218–221.

which convert it into soluble  $\text{Hg}''\text{Cl}_2$ ; *becomes black when AmHO is poured upon it, but does not dissolve.*

218.  $\text{AmHO}$  (or  $\text{KHO}$ ); black precipitate, insoluble in excess.

219.  $\text{H}_2\text{S}$ ; black precipitate ( $\text{Hg}'_2\text{S}$ ); this precipitate, like  $\text{Hg}'/\text{S}$  (144), is insoluble in  $\text{Am}_2\text{S}$ , in hot strong  $\text{HCl}$ , and in hot strong  $\text{HNO}_3$ , but is easily dissolved when heated with a mixture of  $\text{HNO}_3$  and  $\text{HCl}$ .

220. GROUP I.—TABLE OF DIFFERENCES.

Test.	Pb.	Ag.	Hg'.
<i>a. For liquids.</i>			
1. Add $\text{HCl}$ .	White precipitate: <i>soluble in boiling water; AmHO neither dissolves the precipitate nor changes its color.</i>	White curdy precipitate: <i>insoluble in boiling water, easily soluble in warm AmHO, reprecipitated from this solution by adding <math>\text{HNO}_3</math> in excess.</i>	White precipitate: <i>insoluble in boiling water, and insoluble in AmHO, but blackened by it.</i>
2. Add $\text{AmHO}$ .	White precipitate: <i>insoluble in excess.</i>	Brown precipitate: <i>readily soluble in excess.</i>	Black precipitate: <i>insoluble in excess.</i>
3. Add $\text{K}_2\text{CrO}_4$ .	Bright yellow precipitate.	Chocolate-red precipitate.	Scarlet-red precipitate.
<i>b. For solids.</i>			
4. Fused with $\text{Na}_2\text{CO}_3$ on charcoal in the inner blowpipe flame.	White malleable globules of metal which mark paper; yellow incrustation on the charcoal.	White malleable globules or scales which do not mark paper; no incrustation.	No metallic globules.
5. Heated in a bulb-tube, mixed with $\text{Na}_2\text{CO}_3$ .	—	—	Gray sublimate, consisting of globules of metallic mercury.

By the above differences any one member of this group is readily distinguished.

221. (s.) *The separation and detection of Pb, Ag, and Hg', when mixed, depend upon:*

1st. The solubility of  $\text{PbCl}_2$  in boiling water, in which  $\text{AgCl}$  and  $\text{Hg}'_2\text{Cl}_2$  are insoluble.

2d. The solubility of  $\text{AgCl}$  in  $\text{AmHO}$ , in which  $\text{Hg}'_2\text{Cl}_2$  is insoluble.

The presence of each member, when thus separated, is then confirmed by one of its special tests.

**222. (s.)** A solution which may contain Pb, Ag, Hg' is examined by first precipitating by HCl added in excess, then filtering and adding more HCl to the clear filtrate to ascertain that no further precipitate is produced. The precipitate is then examined by Table I (433).


The foregoing reactions only include the metals of common occurrence. For information concerning the detection and separation of the rarer metals, which will only be required by a somewhat advanced student, the appendix or larger analytical treatises must be consulted. The metals gold and platinum, however, are briefly noticed here on account of their frequent employment for chemical and other purposes. These metals are dissolved only by a mixture of HCl and HNO<sub>3</sub>; their most important reactions are given below.

**GOLD (Au).—Use AuCl<sub>3</sub> solution.**

Gold solutions are usually bright yellow in color.

**223.** SnCl<sub>2</sub> containing a little SnCl<sub>4</sub>; purple or dark-brown precipitate ("Purple of Cassius"), best seen by performing the experiment in a white porcelain dish.

**224.** FeSO<sub>4</sub> produces either at once, or on heating the solution, a very finely divided precipitate of Au; the liquid usually appears bluish by transmitted light; always copper-red by reflected light:




**224 a.** H<sub>2</sub>SO<sub>4</sub> produces on boiling a similar precipitate of Au; by boiling the liquid for some time in a porcelain dish, the Au settles in small black masses and the solution loses its color.

**PLATINUM (Pt).—Use PtCl<sub>4</sub> solution.**

**225.** AmCl added to a strong solution produces on standing for some time, or more quickly on being stirred or evaporated, a yellow precipitate (Am<sub>2</sub>PtCl<sub>6</sub>); rather soluble in hot water.

**226.** SnCl<sub>2</sub> gives a reddish-brown color in the solution acidified with HCl, owing to the formation of PtCl<sub>2</sub>:



The method of separating and detecting Au and Pt is fully stated in paragraphs 472-474.

## 146 REACTIONS FOR ACID RADICLES.—GROUP I.

**\*\* Note.**—Before commencing the reactions for acids the student may with advantage analyze several substances which are liable to contain any one or more of the members of the metallic groups. If only *one* metal has to be looked for (see column 1, paragraph 539), the directions given in paragraphs 330–369 may be followed, omitting those portions which relate to the detection of acid radicles. If two or more metals may be present (see columns 2, 3, paragraph 539), full directions for ascertaining to which analytical group or groups they belong will be found in the General Table (419); and the group precipitate or precipitates thus obtained may be then examined by the Group Tables (433–439), with which the student will probably already be familiar.

---

### REACTIONS FOR ACID RADICLES.

In the course of analysis acid radicles are usually detected by special tests; they cannot advantageously be precipitated in groups and the members of each group then separated and identified, as is done in the case of metals. Accordingly the arrangement here adopted consists in simply placing together in a group such acid radicles as in some respects resemble one another in their reactions, and at the end of each group stating upon what differences the detection of its members when occurring together depends.

The reactions for acid radicles are worked through in a manner similar to that employed for the reactions of the metals,—a compound containing the radicle (usually a salt) being employed. At the end of each group the student may with advantage detect one or more of its members as was done with the metals.

---

### GROUP I.—SULPHATE GROUP.

The sulphates are the only commonly occurring salts which give with  $\text{BaCl}_2$  a precipitate insoluble in boiling  $\text{HCl}$ . Hydrofluosilicates resemble sulphates in this respect, but differ in so many other reactions that they are considered hereafter (299–302) in connection with fluorides and silicates, to which they are more closely related.

SULPHATES ( $-\text{SO}_4$ ).—Use  $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ .

**227.**  $\text{BaCl}_2$  [or  $\text{Ba}(\text{NO}_3)_2$ ]: white precipitate ( $\text{BaSO}_4$ ), insoluble when boiled with  $\text{HCl}$  or  $\text{HNO}_3$ .

*Note.*—If  $\text{BaCl}_2$  or  $\text{Ba}(\text{NO}_3)_2$  be added to a solution containing much strong  $\text{HCl}$  or  $\text{HNO}_3$ , a white precipitate may be produced consisting of  $\text{BaCl}_2$  or  $\text{Ba}(\text{NO}_3)_2$ , which, however, is readily distinguished from the precipitate of  $\text{BaSO}_4$  by being dissolved when boiled with water.

$\text{Sr}(\text{NO}_3)_2$  and  $\text{Pb}(\text{NO}_3)_2$  give also white precipitates ( $\text{SrSO}_4$  and  $\text{PbSO}_4$ ).

**228.** *Blowpipe Test.*—Solid  $\text{Na}_2\text{SO}_4$  or any solid sulphate, if mixed with solid  $\text{Na}_2\text{CO}_3$ \* and fused on charcoal in the inner blowpipe flame produced from a spirit-flame, yields  $\text{Na}_2\text{S}$ :



The  $\text{Na}_2\text{S}$  is readily detected by detaching the solid mass, after it has cooled, from the charcoal with the point of a knife; placing a portion of it upon a bright silver coin, then moistening it with a drop of water and crushing it by pressure with a knife-blade. After the moistened mass has remained upon the coin for a short time rinse it off, a black stain will then be seen upon the silver where the substance had rested.

Another portion of the cool mass may be moistened with a drop of  $\text{HCl}$ , when the fetid smell of  $\text{H}_2\text{S}$  will be observed, and a piece of paper moistened with  $\text{PbA}_2$  solution and held above it, will be blackened (230).

*Note.*—This test for a sulphate is reliable only when it is known that sulphur in no other form is present in the substance to be tested; it is also necessary that no sulphur should be introduced in the  $\text{Na}_2\text{CO}_3$ , or by the flame. Now coal-gas sometimes contains sufficient sulphur to give the above reaction with pure  $\text{Na}_2\text{CO}_3$ , hence it is always advisable to employ a spirit rather than a gas flame.

*Hydrogen sulphate, or sulphuric acid* ( $\text{H}_2\text{SO}_4$ ), may be detected by producing black charred stains on paper which has been dipped into it and dried by heat; it also has a strong acid reaction to litmus, and evolves hydro-

---

\* Addition of  $\text{Na}_2\text{B}_4\text{O}_7$  is useful in preventing the fused mixture from sinking into the charcoal.

gen with Zn, properties shown by some metallic and acid sulphates.

## GROUP II.—THE CARBONATE GROUP.

The five following classes of salts give off characteristic gases when acted upon by HCl.

\* \* The systematic Tables of Differences, such as are given for each group of the metals, are not drawn out for the acid radicle groups; the student should have no difficulty in constructing such tables for himself, either mentally or on paper, by looking through the reactions.

CARBONATES ( $-''\text{CO}_3$ ).—Use  $\text{Na}_2\text{CO}_3$ , or white marble ( $\text{CaCO}_3$ ).

229.  $\text{HCl}$  or  $\text{H}\bar{\text{A}}$  or almost any other acid, if poured upon some  $\text{Na}_2\text{CO}_3$  in a test-tube, causes carbon-dioxide gas ( $\text{CO}_2$ ) to come off with effervescence. This gas is recognized by its property of turning lime or baryta-water milky by the production in them of insoluble  $\text{CaCO}_3$  or  $\text{BaCO}_3$ .

The test may be tried in several ways:

1. The acid is poured upon the carbonate in a test-tube. A glass rod which has been dipped into some perfectly clear lime-water\* and has a small drop adhering to its end is then introduced into the test-tube, carefully avoiding touching the liquid or the sides of the tube; the drop will quickly become milky.

2. The acid is poured upon the  $\text{Na}_2\text{CO}_3$  in a small beaker, and this is immediately covered with a watch-glass placed with its convex face downwards, having a drop of clear lime-water adhering to the middle of the face; the drop becomes milky.

3. Since  $\text{CO}_2$  gas is much heavier than air, on adding acid the gas evolved from the carbonate will remain in the test-tube if the tube is held erect and its mouth loosely closed with the thumb; by gradually sloping

\* Baryta-water gives a much more decided result than lime-water.


the tube the heavy gas is poured into another test-tube containing lime-water without allowing any liquid to flow out. On closing this tube with the thumb and well shaking it the lime-water will become milky. Or if the inside of the tube has been rinsed with lime-water the sides become white.

4. The  $\text{CO}_2$ , if it is given off in considerable quantity, may also be made to pass into some lime-water through a doubly-bent tube fitted air-tight by a perforated cork or india-rubber stopper into the tube in which the  $\text{CO}_2$  is generated, as shown in Fig. 37; or the apparatus shown in Fig. 38 may be employed.

Of the above methods 1 and 3 are the most simple; by method 4 the  $\text{CO}_2$  is not found unless given off in somewhat large quantity.

*Note.*—The following precautions must be attended to. Since the milkiness at first caused by  $\text{CO}_2$  disappears gradually when more  $\text{CO}_2$  is absorbed, if the lime-water does not at once become milky, it must be constantly watched to avoid the risk of its becoming milky and again clear before being seen.


Also if the addition of the first few drops of acid does not cause an effervescence of  $\text{CO}_2$  more acid should be added, as many carbonates retain the first portions of  $\text{CO}_2$  by forming acid carbonates:



A solid substance which is being tested by  $\text{HCl}$  for  $\text{CO}_2$  should be first moistened so as to drive out the air-bubbles entangled in it, which, in coming off, might make one suspect effervescence of  $\text{CO}_2$ .

*Hydrogen carbonate* or *carbonic acid* ( $\text{H}_2\text{CO}_3$ ), can only exist in dilute aqueous solution; addition of lime-water to its solution causes milkiness; but, since other substances in solution behave in a similar way, it is best found by boiling the liquid and passing the steam with the  $\text{CO}_2$  which accompanies it into lime-water; the lime-water becomes milky.

FIG. 37. FIG. 38.



**SULPHIDES** (—"S).—Use  $\text{FeS}$  for solid, and  $\text{Am}_2\text{S}$  for liquid.

**230.**  $\text{HCl}$  (or  $\text{H}_2\text{SO}_4$ ) poured upon a small fragment

of  $\text{FeS}$  causes  $\text{H}_2\text{S}$  gas to come off with effervescence; this gas is detected by its fetid smell, resembling that of rotten eggs, and by its property of blackening  $\text{Pb}\bar{\text{A}}_2$  solution by the formation of  $\text{PbS}$ . The gas may be made to act upon the  $\text{Pb}\bar{\text{A}}_2$  solution by the methods given for  $\text{CO}_2$  (229), or better by placing a piece of filter-paper moistened with  $\text{Pb}\bar{\text{A}}_2$  solution on the end of a glass rod so that one-half of the slip adheres to the glass rod and the other hangs free, and holding the rod in the gas as in the test for  $\text{CO}_2$  (229, 1).

*Note.*—This test is made much more delicate if the paper is moistened with a solution prepared by adding  $\text{KHO}$  to boiling  $\text{Pb}\bar{\text{A}}_2$  solution until the precipitate first formed dissolves; a test-paper thus prepared becomes intensely blackened by  $\text{H}_2\text{S}$ .

**231.** Solutions of soluble sulphides if dropped upon a bright silver coin produce a black stain ( $\text{Ag}_2\text{S}$ ); this may be removed by rubbing the coin with moist lime.

**232.** Insoluble sulphides which give off  $\text{H}_2\text{S}$  with  $\text{HCl}$  (*e. g.*,  $\text{FeS}$ ) produce a black stain when placed on a silver coin and moistened with a drop of  $\text{HCl}$ .

**233.**  $\text{Pb}\bar{\text{A}}_2$  solution gives with soluble sulphides a black precipitate ( $\text{PbS}$ ). The solution produced by adding  $\text{KHO}$  in excess to  $\text{Pb}\bar{\text{A}}_2$  solution is much more delicate than  $\text{Pb}\bar{\text{A}}_2$  alone.

**234.** Many sulphides (*e. g.*,  $\text{FeS}_2$ , “Iron Pyrites”) and also free  $\text{S}$ , if strongly heated in a tube open at both ends and held obliquely in the flame, or in the tube shown in Fig. 35 (p. 132), give off  $\text{SO}_2$  gas, recognized by its pungent smell and by turning paper moistened with  $\text{K}_2\text{Cr}_2\text{O}_7$  green.



**235.** Reaction (306) given under  $\text{HCy}$  may be employed also as a very delicate test for soluble sulphides. Test (228) for a sulphate, also answers for a sulphide.

*Hydrogen sulphide*, or *hydrosulphuric acid* ( $\text{H}_2\text{S}$ ), in aqueous solution, is easily recognized by (231) or (233), and, unless the solution is very dilute, also by its smell, and, by suspending lead-paper over the liquid;  $\text{H}_2\text{S}$  gas is detected by (230).

SULPHITES ( $-''\text{SO}_3$ ).—Use  $\text{Na}_2\text{SO}_3$ .

236.  $\text{HCl}$  (or  $\text{H}_2\text{SO}_4$ ) poured upon  $\text{Na}_2\text{SO}_3$  and warmed gives off  $\text{SO}_2$  gas, known by its pungent smell and by turning  $\text{K}_2\text{Cr}_2\text{O}_7$  solution green; the  $\text{K}_2\text{Cr}_2\text{O}_7$  solution may be exposed to the gas by the methods described under (229), preferably by introducing a glass rod with a drop hanging upon its end into the gas; or by moistening a small strip of filter-paper with the  $\text{K}_2\text{Cr}_2\text{O}_7$  solution and making it adhere by one-half of its length to the rod (230), the other half hanging freely, and introducing it into the gas as in (229, 1).

237.  $\text{BaCl}_2$ : white precipitate ( $\text{BaSO}_3$ ), entirely soluble in  $\text{HCl}$ , unless some  $\text{Na}_2\text{SO}_4$  is present, when  $\text{BaSO}_4$  remains undissolved; on adding  $\text{Cl}$  or  $\text{Br}$  water to the  $\text{HCl}$  solution,  $\text{BaSO}_4$  is formed and precipitated.



238.  $\text{Na}_2\text{SO}_3$  added to a mixture of  $\text{HCl}$  and  $\text{Zn}$  which is giving off pure  $\text{H}$ , immediately causes an evolution of  $\text{H}_2\text{S}$  with the  $\text{H}$ ; the  $\text{H}_2\text{S}$  is detected by its smell or by blackening a piece of filter-paper moistened with solution of  $\text{PbO}$  in excess of  $\text{KHO}$ , see note (230).

*Hydrogen sulphite*, or *sulphurous acid* ( $\text{H}_2\text{SO}_3$ ), can be found in aqueous solution by adding a drop of  $\text{K}_2\text{Cr}_2\text{O}_7$  solution, which will at once become green;  $\text{SO}_2$  gas is found as directed under (236).

THEIOSULPHATES\* ( $=''\text{S}_2\text{O}_3$ ).—Use  $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$ .


239.  $\text{HCl}$  (or  $\text{H}_2\text{SO}_4$ ) causes no immediate change in the cold, but the liquid on standing or on being warmed deposits a precipitate of yellow  $\text{S}$ , and  $\text{SO}_2$  is given off; the  $\text{SO}_2$  is recognized by its pungent smell and by changing the color of  $\text{K}_2\text{Cr}_2\text{O}_7$  solution (236).

240.  $\text{Fe}_2\text{Cl}_6$ : gives a reddish-violet color which vanishes after a short time or immediately when heated, the  $\text{Fe}_2\text{Cl}_6$  solution at the same time loses its yellow color owing to the change of  $\text{Fe}_2\text{Cl}_6$  into  $\text{FeCl}_3$ .

\* Formerly known as Hyposulphites.



240 a.  $AgNO_3$ ; a white precipitate ( $Ag_2S_2O_3$ ), becoming black after standing for a short time, or immediately upon being heated, owing to the formation of  $Ag_2S$ :



The above precipitate ( $Ag_2S_2O_3$ ) dissolves very readily in excess of the  $Na_2S_2O_3$  solution, hence it is most easily obtained by dropping the latter into some  $AgNO_3$  solution.


*Hydrogen theiosulphate*, or *theiosulphuric acid* ( $H_2S_2O_3$ ), is extremely unstable, rapidly separating into  $S + H_2SO_4$ .

**HYPPOCHLORITES** ( $-ClO$ ).—Use  $NaClO \cdot NaCl$  ( $Na_2Cl_2O$ ), solution.

Hypochlorites give off a faint smell of  $Cl$  in the air if solid or in strong solution.

241.  $HCl$  (or  $H_2SO_4$  or even  $H_2\bar{O}$ ) in the cold, or when gently warmed if much water is present, sets free  $Cl$  gas, which may be recognized by its smell, its yellowish color, and by bleaching a piece of moistened litmus-paper, which is held for a short time in the tube without touching the liquid or the sides of the tube.

A piece of litmus shaken up with the solution, without addition of acid, is also bleached, the  $CO_2$  in the air probably liberating the  $Cl$ ; on addition of a drop of any acid the bleaching is very rapid:



242.  $Pb(NO_3)_2$  (or  $Pb\bar{A}_2$ ) solution added in large quantity gives a white precipitate, becoming reddish, and then dark-brown ( $PbO_2$ ) when boiled for a short time.

$MnCl_2$  likewise gives a dark-brown precipitate on heating.

*Hydrogen hypochlorite*, or *hypochlorous acid* ( $HClO$ ), is a yellow liquid with sweetish smell, which if strong is very unstable; it bleaches litmus, and evolves  $Cl$  when warmed with  $HCl$ .  $Cl_2O$  gas is of a deep yellowish-green color, and has an irritating smell, it dissolves easily in water to form the above acid.

NITRITES ( $= \text{'NO}_2$ ).—Use  $\text{KNO}_2$ .

243. Warmed with dilute  $\text{H}_2\text{SO}_4$ , reddish-brown "nitrous" fumes, with a characteristic smell, are given off; they are best seen by looking down the tube at some white object.

244. If cold  $\text{FeSO}_4$  solution be added to  $\text{KNO}_2$  solution, it becomes brown; on adding cold dilute  $\text{H}_2\text{SO}_4$  the color becomes much more intense; it is produced by the combination of  $\text{NO}$  with  $\text{FeSO}_4$ . This color is destroyed by boiling, red fumes being given off (243).

245. If solution of  $\text{KI}$  and several drops of starch solution (made by boiling starch in water, and cooling it) be added to  $\text{KNO}_2$  solution and the liquid be then made acid with  $\text{H}^+\text{A}$ , an intense blue color will be produced, owing to the combination of iodine (set free by the  $\text{HNO}_2$ ) with the starch; the experiment is best done in a white porcelain dish, the liquid, which often appears black at first, being diluted if necessary till its blue color becomes visible. This is a most delicate test, and is rather characteristic if the color is not produced till the  $\text{H}^+\text{A}$  is added.

*Hydrogen nitrite, or nitrous acid* ( $\text{HNO}_2$ ), is extremely unstable, decomposing into  $\text{HNO}_3$  and  $\text{NO}$ .

\*.\* The student will have no difficulty in detecting any one of the above acid radicles occurring singly in a liquid or solid; warming with  $\text{HCl}$  will usually indicate which of them is present, and its presence may then be confirmed by a second test chosen from amongst its reactions as stated above.

246. (s.) Most of the acids formed by the salts of Group II on addition of  $\text{HCl}$ , react upon one another; e.g.,  $\text{HClO}$  destroys  $\text{HNO}_2$ ,  $\text{H}_2\text{S}$ , and  $\text{H}_2\text{SO}_3$ , and  $\text{H}_2\text{S}$  destroys  $\text{H}_2\text{SO}_3$ . The analyst may often by skilful use of the above reactions detect these acids when mixed, but their detection in certain mixtures is a problem only soluble by careful consideration, and not unfrequently insoluble.

$\text{SO}_2$ , if evolved, may be detected by  $\text{K}_2\text{Cr}_2\text{O}_7$  paper;

H<sub>2</sub>S by PbA<sub>2</sub> paper; Cl by bleaching moist litmus-paper; N<sub>2</sub>O<sub>3</sub> by its red color. But CO<sub>2</sub> can only be detected in presence of much SO<sub>2</sub> by passing the gases through K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> solution which absorbs SO<sub>2</sub>, CO<sub>2</sub>, passing on and being detected by lime-water.

### GROUP III.—NITRATE GROUP.

The two classes of salts which follow resemble one another in many respects, more particularly in no member of them being insoluble in water; hence no method of precipitating them is known.

NITRATES (—'NO<sub>3</sub>).—Use KNO<sub>3</sub>.

247. Add to some KNO<sub>3</sub> solution contained in a rather broad test-tube about an equal bulk of strong H<sub>2</sub>SO<sub>4</sub>; cool the liquid by letting a stream of cold water run upon the tube from the tap; then hold the tube in a slanting position, and pour some strong cold FeSO<sub>4</sub> solution in a gentle stream down the inside of the tube. If this solution is poured in with proper care, it will form a distinct layer resting on the surface of the acid liquid, and at the surface of contact of the two layers a *dark-brown* film or ring will appear, sometimes extending upwards towards the surface of the liquid. The color is most easily seen by holding a piece of filter-paper between the tube and a bright flame or window, and looking through the tube at the light. A mere *yellow* color is often produced in the absence of a nitrate, and may be disregarded.

The test is made more delicate by letting the tube stand in a small beaker of cold water for a few minutes, since heat prevents or destroys the brown color. If much nitric acid is present, on mixing the acid and FeSO<sub>4</sub> solution by shaking, the brown color extends upwards, until by the heat evolved by dilution of the acid the NO is expelled, forming brown hues in the test-tube, and the liquid loses its dark-brown color (244).

248. Place some small scraps of Cu in some KNO<sub>3</sub>,

solution, and add strong  $\text{H}_2\text{SO}_4$ ; reddish nitrous fumes appear at once or on warming the tube; they are best seen, if small in quantity, by looking down the tube at a white surface. The liquid at the same time becomes blue from the presence in it of  $\text{Cu}(\text{NO}_3)_2$ :



$\text{NO}$  is itself a colorless gas, but yields on mixture with air  $\text{N}_2\text{O}_3$  and  $\text{N}_2\text{O}_4$ , which are reddish-brown.

**249.** If some cold dilute indigo solution be poured into cold  $\text{KNO}_3$  solution until it is decidedly blue, and then  $\text{H}_2\text{SO}_4$  be added, the blue color remains unchanged. (Difference from chlorates.)

**250.** If any solid nitrate be fused for some time at a red heat with fusion mixture on platinum foil, oxygen gas is given off slowly with effervescence,\* and  $\text{KNO}_3$  is formed; if the foil after cooling be boiled in water, a portion of the solution may be proved to contain  $\text{KNO}_3$  by the test in paragraph (245); another portion, if acidified with  $\text{HNO}_3$ , will be found to give no precipitate with  $\text{AgNO}_3$ . (Difference from chlorates.)

A solution of a nitrate may also be reduced to nitrite by nascent  $\text{H}$ . Add to a little  $\text{KNO}_3$  solution some dilute  $\text{H}_2\text{SO}_4$  and starch paste with  $\text{KI}$ , no color will be produced, but on dropping in a fragment of  $\text{Zn}$  a deep blue color is seen (245).

**251.** Solid  $\text{KNO}_3$  fused on charcoal in the blowpipe flame "deflagrates," that is to say, the surface of the charcoal burns rapidly like gunpowder.

*Note.*—Many metallic nitrates [*e.g.*,  $\text{Pb}(\text{NO}_3)_2$ ] when heated in a glass tube closed at one end, evolve reddish-brown fumes with a characteristic smell ( $\text{N}_2\text{O}_4$ ) and oxygen— $\text{PbN}_2\text{O}_6 = \text{N}_2\text{O}_4 + \text{O} + \text{PbO}$ . The fumes are recognized by their color and smell, the  $\text{O}$  by its inflaming a glowing splinter of wood.

*Hydrogen nitrate*, or *nitric acid* ( $\text{HNO}_3$ ), has a strong acid reaction, it gives with  $\text{FeSO}_4$  solution a brown ring

---


\* If this experiment is performed in a small hard-glass tube as described in 254, the oxygen can rarely be detected, since it is evolved very slowly; hence the student may perform the ignition on platinum foil and neglect to test for  $\text{O}$ .

(247), and with Cu brown fumes (248) without addition of  $\text{H}_2\text{SO}_4$ . Evaporated with quill-clippings, it stains them bright yellow.

CHLORATES ( $-\text{'ClO}_3$ ).—Use  $\text{KClO}_3$ .

252. If to a *cold* solution of  $\text{KClO}_3$  some cold dilute indigo solution be added drop by drop till the liquid is colored faintly but distinctly blue, then some  $\text{H}_2\text{SO}_4$  or  $\text{NaHSO}_3$  be poured in and the mixture shaken, the blue color of the indigo is at once destroyed. (Difference from nitrates.)

253. *Strong*  $\text{H}_2\text{SO}_4$  poured upon a little solid  $\text{KClO}_3$  becomes orange-red in color, and evolves when shaken a bright yellow gas ( $\text{Cl}_2\text{O}_4$ ):




This gas has a smell somewhat resembling that of chlorine; it explodes readily when gently heated, hence on warming the mixture in the tube crackling sounds or small explosions will be produced. This experiment may be dangerous, unless the  $\text{KClO}_3$  is used in very small quantity, and heated gently with the mouth of the test-tube turned away from the operator.

254.  $\text{AgNO}_3$  produces no precipitate with  $\text{KClO}_3$  solution if it is free from KCl. But if some solid  $\text{KClO}_3$ \* be heated in a test-tube or better in a small ignition-tube (10), after melting it begins to give off oxygen with effervescence; this gas may be recognized by placing in the tube a slip of wood with a spark at its end, which is easily obtained by kindling a match and suddenly blowing out the flame, the spark is caused to burst into a flame. If the tube is then allowed to cool, and the residue dissolved by boiling with some water, a portion of the solution will be found to give no reaction for  $\text{HNO}_3$  (245); but another portion will give a white precipitate ( $\text{AgCl}$ ) with  $\text{AgNO}_3$ , which does not dissolve in  $\text{HNO}_3$ .

\* If the chlorate used for this test is not an alkaline chlorate, fusion mixture must be mixed with it before heating, in order to obtain a soluble chloride in the residue.

even on boiling, but is easily soluble in AmHO (difference from a nitrate): this precipitate is caused by KCl:



**255. Blowpipe Test.**— $\text{KClO}_3$ , if heated on charcoal in the blowpipe flame, causes the charcoal to “deflagrate.”

*Hydrogen chlorate*, or *chloric acid* ( $\text{HClO}_3$ ), is a colorless, odorless liquid, which first reddens and then bleaches litmus paper; on being kept for some time it changes into  $\text{O}_2$ ,  $\text{Cl}_2$ ,  $\text{HClO}_4$ , and  $\text{H}_2\text{O}$ ; the same change occurs rapidly when it is heated.

**256.** As will be seen, there is little difficulty in distinguishing a nitrate from a chlorate. When present together they more or less interfere with one another's reactions; but they may be detected, in the absence of chlorides and nitrites, by heating strongly for a short time, and testing the residue for nitrite and chloride (250 and 254).

#### GROUP IV.—CHLORIDE GROUP.


Chlorides, bromides, and iodides closely resemble one another; they are all precipitated by  $\text{AgNO}_3$  from solutions to which  $\text{HNO}_3$  has been added in excess; in this respect they differ from all salts except cyanides, ferrocyanides, and ferricyanides, and these are easily distinguished by other means.

CHLORIDES (—'Cl).—Use NaCl.

**257.**  $\text{AgNO}_3$  added to a little NaCl dissolved in water, gives a *pure white* precipitate ( $\text{AgCl}$ ), which collects into curdy masses when heated or shaken, and quickly turns violet in sunlight or more slowly in daylight. Decant the water; to one portion of the precipitate add  $\text{HNO}_3$  and warm, it does not dissolve; to another portion add AmHO, it readily dissolves. Decant the liquid from a portion of the precipitate after shaking and letting the precipitate settle, and warm the precipitate with a little strong  $\text{H}_2\text{SO}_4$ , the acid remains colorless, and no colored vapor is given off.

**258.** Solid NaCl warmed with strong  $\text{H}_2\text{SO}_4$  gives off colorless HCl gas, which is recognized by fuming in the air, reddening moistened blue litmus-paper, and making a drop of  $\text{AgNO}_3$  solution on the end of a glass rod milky.


**259.** If solid NaCl be powdered and mixed with finely powdered  $\text{MnO}_2$ , then strong  $\text{H}_2\text{SO}_4$  added and the mixture warmed, chlorine gas comes off:



This gas is recognized by its property of bleaching a piece of moist blue litmus-paper, introduced into the tube on a glass rod. The most delicate way of testing is to warm the mixture in a small beaker covered with a watch-glass which bears on its under surface a piece of moist litmus-paper, and to let the apparatus stand for some time. Moist starch is not colored when held in the air in the tube.

*Note.*—Many samples of commercial  $\text{MnO}_2$  evolve Cl when heated with  $\text{H}_2\text{SO}_4$ ; hence the  $\text{MnO}_2$ , unless specially prepared by precipitation, must be carefully tested before using it for this reaction; or the  $\text{MnO}_2$  and  $\text{H}_2\text{SO}_4$  may be first boiled together as long as any bleaching action is produced when moist litmus is held in the tube, then the substance to be tested is added, and Cl again tested for whilst heating the mixture.

**260.** If an intimate mixture of solid NaCl with three or four times as much  $\text{K}_2\text{Cr}_2\text{O}_7$  in fine powder be made by rubbing the two substances together in a mortar, and this mixture be then poured into the tubulated flask shown in Fig. 38\* (229) (see note below), and mixed with strong  $\text{H}_2\text{SO}_4$  by pouring in the acid and shaking it round in the flask, on warming the flask reddish-brown vapor ( $\text{CrO}_2\text{Cl}_2$ ) will be evolved:



If the neck of the flask be closed by an india-rubber or a glass stopper, and the delivery-tube be made to dip

---

\* The test-tube fitted as shown in Fig. 37 is much less suitable

into water in a test-tube, the vapor, on passing into the water, will impart to it a reddish-yellow color, owing to the formation of  $\text{H}_2\text{CrO}_4$ :



On adding to this yellow liquid excess of  $\text{AmHO}$ , the color becomes pale yellow; on now adding excess of  $\text{H}\bar{\text{A}}$ , the original reddish-yellow color is reproduced, and in this liquid the presence of  $\text{H}_2\text{CrO}_4$ , and therefore indirectly that of  $\text{HCl}$ , may be shown by the formation of a yellow precipitate on addition of  $\text{Pb}\bar{\text{A}}$  solution.

*Note.*—Since by this method the detection of a chloride depends on the formation of  $\text{H}_2\text{CrO}_4$  in the liquid in the test-tube from the vapor or  $\text{CrO}_2\text{Cl}_2$ , great care must be taken that no  $\text{H}_2\text{CrO}_4$  is introduced into that liquid from any other source. Now, since the mixture introduced into the flask contains a chromate, the greatest precaution must be taken that none of it is allowed to get into the bent delivery-tube, and thence into the test-tube. In introducing the powder into the flask, it must be poured down the side of the *perfectly dry* neck opposite to that at which the delivery-tube enters; also, whilst heating the mixture it must not be allowed to rise or splash into the neck of the flask. If these precautions are not taken, and  $\text{H}_2\text{CrO}_4$  as such is introduced into the water in the test-tube, the test is worthless as an indication of the presence of a chloride.

Care must also be taken that the liquid is not sucked back from the test-tube into the flask: this is prevented by raising the delivery-tube out of the water in the test-tube as soon as the heating is stopped.

*Hydrogen chloride* or *hydrochloric acid* ( $\text{HCl}$ ) is a colorless gas, which fumes in the air and dissolves very readily in water; both the gas and its solution render  $\text{AgNO}_3$  solution, which has been acidified with  $\text{HNO}_3$ , milky. Heated with  $\text{MnO}_2$ , the liquid acid evolves chlorine gas.

BROMIDES (—'Br).—Use  $\text{KBr}$ .

261.  $\text{AgNO}_3$  added to a little  $\text{KBr}$  solution gives a *yellowish-white* precipitate ( $\text{AgBr}$ ), which is easily coagulated by heating or shaking the liquid; it is insoluble in  $\text{HNO}_3$ , but rather soluble in  $\text{AmHO}$ , thus somewhat resembling the  $\text{AgCl}$  precipitate, from which it is



distinguished by its color. By shaking well, allowing to settle, decanting the liquid, and heating the precipitate with strong  $\text{H}_2\text{SO}_4$ , no violet vapor is evolved, the precipitate thus differing from  $\text{AgI}$ .

**262.** Solid  $\text{KBr}$  when heated with strong  $\text{H}_2\text{SO}_4$  gives off *reddish-brown* vapor of  $\text{Br}$ ,\* which somewhat resembles  $\text{Cl}$  in smell, but differs by its color and by turning moist starch *orange-red*, the color disappearing by heat. This last test is best performed by dipping the wetted end of a glass rod into some powdered starch, and then moistening the adhering starch by breathing upon it several times. If  $\text{MnO}_2$  be mixed with the  $\text{KBr}$  powder before adding  $\text{H}_2\text{SO}_4$ , the  $\text{Br}$  is evolved in greater quantity and more readily.

**263.** *Cl-water*, added carefully drop by drop to a little  $\text{KBr}$  solution whilst shaking it, liberates  $\text{Br}$ , which gives the solution an orange-red color; excess of *Cl-water* must be carefully avoided, as it destroys the color:



On warming a part of this solution in a dish, brown fumes are given off which color *orange-red* some starch held in the vapor as described in the preceding reaction.

If  $\text{CS}_2$  be added to another portion and the liquids are then well shaken together, the  $\text{Br}$  is dissolved by the  $\text{CS}_2$ , and on standing the reddish-brown  $\text{CS}_2$  solution sinks to the bottom; on adding a little  $\text{KHO}$  solution and shaking well, the color of the  $\text{Br}$  again disappears from the  $\text{CS}_2$ , owing to the formation of the colorless salts  $\text{KBr}$  and  $\text{KBrO}_3$ :



**264.** A mixture of solid  $\text{KBr}$  and  $\text{K}_2\text{Cr}_2\text{O}_7$  heated with strong  $\text{H}_2\text{SO}_4$  as in (260), gives off reddish-brown vapor of  $\text{Br}$ :




---

\*  $\text{KBr}$  solution does not so easily give off  $\text{Br}$  vapor, but the liquid becomes colored reddish-brown.

This vapor, if received in a little water, imparts its color to the water; if the liquid is shaken up with  $\text{CS}_2$ , however, the Br is dissolved away from the water by  $\text{CS}_2$ , which forms a brown layer at the bottom of the water: by shaking well after addition of AmHO or KHO the color of the solution is destroyed. (Difference from HCl.)


*Hydrogen bromide*, or *hydrobromic acid* (HBr), is a gas resembling HCl; its solution differs by evolving Br when heated with  $\text{MnO}_2$ .

IODIDES (-'I).—Use KI.

**265.**  $\text{AgNO}_3$  added to a little KI solution, gives a *yellow* precipitate (AgI), which is easily coagulated by heating or shaking the liquid: it is insoluble in  $\text{HNO}_3$ , and very slightly soluble in AmHO.

This precipitate differs from AgCl and AgBr by its color, and its slight solubility in AmHO, but more particularly by its behavior when heated with strong  $\text{H}_2\text{SO}_4$ . If the precipitate, coagulated by heating or shaking the liquid, is allowed to settle and the liquor decanted, then strong  $\text{H}_2\text{SO}_4$  poured in and heated with the precipitate, the acid becomes colored red, and violet vapors of iodine are seen especially on looking down the tube; the color is particularly distinct after the tube has been allowed to cool for a short time. This vapor of iodine may be detected when almost invisible by holding in the air of the tube a glass rod previously dipped into freshly made starch solution, which will become colored blue.

**266.**  $\text{CuSO}_4$  mixed with  $\text{H}_2\text{SO}_3$  or  $\text{FeSO}_4$  solution, which reduces it to  $\text{Cu}_2\text{SO}_4$ , gives a white precipitate ( $\text{Cu}_2\text{I}_2$ ); warming, or the addition of a drop of AmHO, causes this precipitate to form more quickly:



If this precipitate is allowed to settle and the water poured off, or if it be filtered, and the precipitate is then

warmed with strong  $\text{H}_2\text{SO}_4$ , the liquid becomes violet and gives off violet fumes of I.

Solutions of chlorides and bromides are not precipitated thus by  $\text{Cu}_2\text{SO}_4$ .

**267.** Solid KI if warmed with strong  $\text{H}_2\text{SO}_4$  gives off violet vapor of I,\* which, if in large quantity, cools on the sides of the tube to black solid I: this vapor colors starch solution blue, as may be proved by dipping a rod moistened with freshly made starch solution† into it, or by pouring the heavy fumes out into a white porcelain dish whose inside has been wetted with starch solution, or by dipping into the vapor a slip of filter-paper moistened with starch solution. The I is more readily and plentifully evolved if some  $\text{MnO}_2$  be mixed with the KI before adding the  $\text{H}_2\text{SO}_4$ .

**268.** *Cl-water* added drop by drop (carefully avoiding excess, which would form colorless  $\text{ICl}_3$ ), sets free I, which dissolves in the water, giving a brown liquid; divide this into three parts:

1. Heat one part; violet vapor of I is given off and is best seen by looking down the mouth of the tube at a white surface. The color is most distinctly seen if, before heating, some strong  $\text{H}_2\text{SO}_4$  is added to the liquid. If a glass rod is dipped into starch solution and held in the vapor, the starch becomes blue.

2. To another portion starch solution is added, which gives an intense blue color, often appearing black unless much starch is added and the liquid largely diluted; on warming the solution the color disappears, but often reappears when the liquid is cooled. To insure the disappearance of the color on being heated, the solution must be diluted and the starch added in excess.

3. To the third portion add a drop of  $\text{CS}_2$  and shake well; the  $\text{CS}_2$  dissolves the I, and gradually settles to the bottom having a beautiful violet color. Add a little

---

\* KI solution does not so easily give off purple vapor, but the liquid becomes reddish-brown or red.

† The starch solution must always be freshly made, as after being kept, it loses the property of yielding a blue color with iodine.

KHO solution and shake well, the violet color of the  $\text{CS}_2$  will disappear owing to the formation of the colorless salts KI and  $\text{KIO}_3$ .

*Note.*—The last two methods of testing for the free I are by far the most delicate known; but Cl-water is by no means the best reagent for setting I free from its compounds, because if added in excess it combines with the I and prevents its detection. The reagents described in (269) are much better suited for liberating I from its compounds.

269. If a little  $\text{KNO}_3$  is added to the KI solution, and then any dilute acid ( $\text{H}\bar{\text{A}}$ , HCl or  $\text{H}_2\text{SO}_4$ ) poured in,  $\text{HNO}_3$  is produced which liberates I. A solution of  $\text{N}_2\text{O}_4$  in dilute  $\text{H}_2\text{SO}_4$ , made by heating  $\text{Pb}(\text{NO}_3)_2$  strongly and passing the red fumes into dilute  $\text{H}_2\text{SO}_4$ , also sets free I from KI. The I thus liberated in solution may be identified by the three methods described in (268), of which the third is the most delicate.

These two reagents for liberating I, do not hinder its detection if they are added in excess, and they are also useful because they do not liberate Br from its compounds.

270.  $\text{HgCl}_2$  gives with KI solution a scarlet precipitate, with KBr or NaCl it gives no precipitate;  $\text{PbA}_2$  gives with KI solution a *bright yellow* precipitate, with KBr or NaCl a *white* precipitate.

*Hydrogen iodide*, or *hydriodic acid* (HI), is a gas resembling HCl and HBr; its solution differs by evolving I when heated with  $\text{MnO}_2$ .

271. A *chloride*, *bromide*, or *iodide*, is easily detected by the foregoing tests. The reactions given in paragraphs (259) and (260) are perhaps the most characteristic tests for a chloride, those described in (263) and (264) for a bromide, whilst an iodide is detected with the greatest ease and certainty by reaction (269).

The color of the precipitate produced by  $\text{AgNO}_3$  solution in the solution made acid with  $\text{HNO}_3$ , and its behavior with  $\text{AmHO}$ , though less reliable than the above tests, serve to indicate which of these acid radicles is present,  $\text{AgCl}$  being *pure white* and very easily soluble

in AmHO, AgBr being *pale yellow* and not readily soluble in AmHO, whilst AgI is *primrose-yellow* and almost insoluble in AmHO. See also the distinctive action of hot strong  $\text{H}_2\text{SO}_4$  on AgI, described in paragraph (265).

272. (s.) *The detection of a chloride, bromide, and iodide when occurring together.*

The method of examination varies according to whether an iodide is present or absent; since a chloride cannot be detected in the presence of an iodide, the latter, if present, must be separated before proceeding to examine for the chloride (see 273). In paragraph (274) is given a process for detecting an iodide and bromide; it is of value only when a chloride has not to be tested for.

273. (s.) To a small portion of the solution, made just acid if necessary by addition of dilute  $\text{H}_2\text{SO}_4$ , add a little cold freshly prepared starch solution and then strong  $\text{HNO}_3$ , or either of the reagents mentioned in (269), drop by drop; a dark-blue coloration\* shows the *presence of an iodide*. The remainder of the solution is examined for a chloride and bromide by the directions given below under A or B, according as an iodide is present or absent:

A. *An iodide is present.*—Add to the solution a reagent consisting of a mixture of  $\text{CuSO}_4$  solution with half its measure of strong  $\text{H}_2\text{SO}_4$  solution, and warm gently for a short time;  $\text{Cu}_2\text{I}_2$  will be precipitated (266). In order to be sure that the iodide is entirely precipitated, a small quantity of the liquid must be filtered and warmed with a little more of the above reagent; if any further precipitate is caused, the filtered portion is returned to the unfiltered liquid, more of the reagent is added to the whole, which is then warmed for a time and again tested as above, to see if the precipitation is complete; these trials are repeated until a small portion, on filtration, gives no further precipitate when warmed with more of the reagent. The whole is then filtered till clear, pure NaHO is added in excess to the filtrate

---

\* Black, unless the starch solution has been added in sufficient quantity to the dilute solution.

and the liquid boiled. The precipitate thus produced is filtered off, and the filtrate, now free from iodide, is tested for bromide and chloride by the directions given in column *B*.

*B. No iodide is present.*—Evaporate the solution (made alkaline, if not already so, by the addition of pure  $\text{NaHO}$ ) to dryness in a porcelain dish, scrape the residue off the sides of the dish and mix it with three or four times as much powdered  $\text{K}_2\text{Cr}_2\text{O}_7$  by rubbing the two substances together in the dish with a pestle (260). Transfer this mixture into the small tubulated flask (Fig. 38, p. 149) by pouring it down the perfectly dry neck on the side opposite to that at which the tube enters, and taking the greatest care to let none of the powder enter the delivery-tube. Then pour into the flask strong  $\text{H}_2\text{SO}_4$  sufficient to cover the powder to the depth of about a quarter of an inch, at once close the neck with an india-rubber or glass stopper, and dip the end of the delivery-tube an inch into water contained in a test-tube. Mix the contents of the flask by cautious shaking, and warm the mixture gently, carefully guarding against allowing any of the mixture to get into the delivery-tube, and thence into the water in the test-tube.\* As soon as no more reddish-brown vapor is visible in the flask or its delivery-tube, discontinue the heating, and immediately withdraw the test-tube. If no colored vapor has been evolved and the water in the test-tube is colorless, chloride and bromide are absent. If the water is colored pour into it  $\text{CS}_2$  sufficient to form a layer at the bottom of the tube about as large as a small nut; close the mouth of the tube with the thumb and shake vigorously; the  $\text{CS}_2$  on settling shows a brown color, indicating the *presence of Br from a bromide*.

Separate the water from the  $\text{CS}_2$  by pouring it through a wetted filter, add to the filtrate  $\text{AmHO}$  in excess, then  $\text{HA}$  in excess, and  $\text{PbA}_2$  solution. A yellow precipitate indicates the presence of a chromate, which proves indirectly that a *chloride was present*.

---

\* See the precautions in the note at the end of par. 260.

**273 a.** (s.) The above method will also serve for the examination for chloride, bromide, and iodide in the  $\text{AgNO}_3$  precipitate. The precipitate is dried, and fused for about ten minutes in a porcelain crucible with three or four times as much fusion mixture; the cool mass is boiled with  $\text{H}_2\text{O}$  for some time and filtered, the filtrate being examined as directed above (273).

**274.** (s.) If only an iodide and a bromide have to be tested for, the following method may be employed. Pour into the solution  $\text{CS}_2$  sufficient to form a large drop at the bottom; add Cl-water drop by drop, shaking thoroughly after each addition: an iodide, if present, shows first by its liberated iodine coloring the  $\text{CS}_2$  purple, but on continuing the addition of Cl-water this color is destroyed; a bromide is then detected by a brown coloration of the  $\text{CS}_2$ , which, however, is also destroyed if too much Cl-water is added.

#### GROUP V.—PHOSPHATE GROUP.

Phosphates and arsenates resemble one another closely in many reactions; the points of difference they present from one another and from other salts will, however, if carefully attended to, render their detection a simple matter.

PHOSPHATES ( $-\text{PO}_4$ ).—Use  $\text{NH}_4\text{PO}_4 \cdot 12\text{H}_2\text{O}$ .

**275.**  $\text{MgSO}_4$  solution, to which some  $\text{AmCl}$  has been added, and then a little  $\text{AmHO}$ , gives with  $\text{Na}_2\text{HPO}_4$  a white crystalline precipitate ( $\text{MgAmPO}_4$ ): this precipitate is almost insoluble in  $\text{AmHO}$ , but easily soluble in acids. If very little  $\text{Na}_2\text{HPO}_4$  is present, the precipitate appears only on warming, stirring, or shaking well, and letting stand.

**276.**  $\text{Fe}_2\text{Cl}_6$  (a few drops only) added after a little  $\text{HA}$  and some  $\text{NaA}$  solution gives, especially on being heated, a yellowish-white precipitate of  $\text{Fe}'''\text{PO}_4$ .

**277.**  $\text{AmHMoO}_4$ , dissolved in  $\text{HNO}_3$ , gives a yellow precipitate if a drop of  $\text{Na}_2\text{HPO}_4$  solution is added to it; the formation of the precipitate is hastened by *gentle*

warming, stirring, or shaking, but it often appears only after a time. Pour off a part of the liquid and show that the precipitate is soluble in alkalies ( $\text{KHO}$ ,  $\text{NaHO}$ ,  $\text{AmHO}$ ), and insoluble in acids, especially in  $\text{HNO}_3$ .

*Note.*—This reaction, if properly performed, is the most delicate test known for a phosphate, but careful attention must be paid to the following precautions: The  $\text{AmHMoO}_4$  solution must be prepared according to the directions given in par. 526, remark 40; the solution to be tested, must not be alkaline to test-paper; it is best made distinctly acid with  $\text{HNO}_3$ ; it is then to be added *in small quantity only* to some of the  $\text{AmHMoO}_4$  solution in a test-tube, more of the liquid being added only if after *gently* warming and stirring or shaking well no yellow precipitate forms after a few minutes. This last precaution must be particularly observed, as an excess of phosphate altogether prevents the formation of the precipitate: show that this is the case by pouring into some  $\text{Na}_2\text{HPO}_4$  solution, contained in a test-tube and made acid with  $\text{HNO}_3$ , a few drops of  $\text{AmHMoO}_4$  solution; it will be found that no precipitate will form even on heating and shaking the liquid, since the phosphate is present in large quantity compared with the  $\text{AmHMoO}_4$ ; but on adding a few drops of this liquid to some fresh  $\text{AmHMoO}_4$  solution the precipitate appears. It must also be remembered that  $\text{HCl}$  retards or prevents the formation of this precipitate, and therefore an  $\text{HNO}_3$  solution should always be used when possible.

**278.**  $\text{AgNO}_3$ ; yellow precipitate ( $\text{Ag}_3\text{PO}_4$ ): pour off a portion and show the precipitate to be soluble in  $\text{AmHO}$  and in  $\text{HNO}_3$ .

**278 a.** A very delicate test for the presence of P in the form of a phosphate or otherwise, consists in strongly heating the dry solid substance with a small piece of Na in a little tube of hard glass closed at one end. On breaking the tube and breathing upon the powdered substance a smell of onions, due to  $\text{H}_2\text{P}$ , is perceived.

A phosphate made into a paste with strong  $\text{H}_2\text{SO}_4$  and heated strongly at the tip of the inner blowpipe flame, gives a bluish-green color to the flame.

*Hydrogen phosphate, or phosphoric acid* ( $\text{H}_3\text{PO}_4$ ), is a colorless crystalline substance; its solution is strongly acid, but does not char paper and organic substances. (Diff. from  $\text{H}_2\text{SO}_4$ .) It yields reactions (275 and 277).

---

ARSENATES ( $-\text{AsO}_4$ ). — Use  $\text{Na}_2\text{HASO}_4 \cdot 12\text{H}_2\text{O}$  solution.

**279.** The precipitates formed in reactions (275, 276, 277) for phosphates are precisely similar in appearance and general properties to those formed with the same reagents by arsenates. The



precipitate, however, which is formed by an arsenate with  $\text{AgNO}_3$ , differs from that given by a phosphate by being *brown* in color; also  $\text{AmHMoO}_4$  gives with an arsenate a precipitate *only on boiling*.

**280.** The following differences serve to detect and separate a phosphate and an arsenate:

1. The fact that an arsenate only yields a precipitate with  $\text{AmHMoO}_4$  when the liquid is *boiled*, whereas the corresponding precipitate with a phosphate is produced by a *gentle heat*.

2. An arsenate solution, if boiled with strong  $\text{HCl}$ , gives when  $\text{H}_2\text{S}$  is passed into the hot liquid, first a white precipitate of  $\text{S}$ , then yellow  $\text{As}_2\text{S}_3$ ; a phosphate solution, under the same circumstances, gives no precipitate.

3. In a *perfectly neutral* solution,  $\text{AgNO}_3$  gives with an arsenate solution a *brown* precipitate, with a phosphate a *canary-yellow* precipitate.

**281.** Hence, if occurring separately, a phosphate and arsenate may be distinguished by one or more of these differences.

**281 a.** The simplest method of ascertaining which of these two acids is present is to throw the precipitate yielded by  $\text{AmCl}$ ,  $\text{AmHO}$ , and  $\text{MgSO}_4$  upon a filter, wash it with a little cold water, and drop  $\text{AgNO}_3$  upon it; if the precipitate consisted of  $\text{MgAmAsO}_4$  it will become *brown*, if of  $\text{MgAmPO}_4$  *canary-yellow* (difference 3). Or the precipitate produced by  $\text{AmCl}$ ,  $\text{AmHO}$ , and  $\text{MgSO}_4$  may be examined for arsenate by dissolving a part of it in dilute  $\text{HNO}_3$ , adding  $\text{AgNO}_3$  as long as it causes any white precipitate, and then very dilute  $\text{AmHO}$  gradually drop by drop; if arsenate is present a *brown* precipitate will form just before the liquid becomes alkaline to test-paper; phosphate gives a *yellow* precipitate. Instead of mixing the dilute  $\text{AmHO}$  it may be poured upon the top of the acid liquid when the color appears at the surface of contact.

**282.** If both arsenate and phosphate have to be tested for,  $\text{AmCl}$ ,  $\text{AmHO}$ , and  $\text{MgSO}_4$  are added as long as any further precipitate is caused; the precipitate is

filtered off\* and dissolved in a little boiling strong HCl,  $H_2S$  is passed for several minutes into this hot HCl solution, when the presence of arsenate is shown as above (par. 280, 2); the precipitate is filtered off if any has formed, and  $H_2S$  passed into the boiling filtrate; if any further precipitate is produced, the gas must be passed again until it no longer causes a precipitate; the clear filtrate from this precipitate is then evaporated just to dryness, dissolved in a little dilute  $HNO_3$ , and tested for phosphate by adding a few drops of it to  $AmHMoO_4$  solution and gently warming. Arsenite, if present, may be detected by  $H_2S$  in the filtrate from the Mg precipitate, acidified with HCl; yellow  $As_2S_3$  is precipitated.

#### UNGROUPED ACID RADICLES (INORGANIC).

The salts of the following acid radicles present no marked resemblances in their reactions to one another or to other salts, and are usually detected by special tests.

BORATES ( $-BO_2$ , or  $-BO_3$ ).—Use  $Na_2B_4O_7 \cdot 10H_2O$

**283.** If some  $Na_2B_4O_7$  be stirred with a little dilute HCl on a watch-glass and a strip of turmeric paper be half immersed in the liquid, then dried on a watch-glass in the steam-oven or on a water-bath,† the part which was moistened appears *reddish-brown*, and becomes *bluish-black* when moistened with  $AmHO$ . This is a most useful test for a borate.

**284.** If some methylated spirit be poured upon a little solid  $Na_2B_4O_7$  in a test-tube or porcelain dish, some strong  $H_2SO_4$  added, and the mixture be heated and set

\* A small portion of the precipitate may be filtered off upon another filter and tested at once for arsenate by dropping upon it  $AgNO_3$  solution. If the precipitate turns brown, arsenate is present and must be separated in the other part of the Mg precipitate before testing for phosphate; if the precipitate turns yellow, phosphate alone is present.

† The paper may also be dried over a small flame, if it is not heated above steam heat ( $100^\circ C.$ ).

fire to, the spirit burns with a green-edged flame, seen best by blowing the flame out after it has burnt for a time, and then relighting it. This test is not quite characteristic, since some chlorides yield a similar green flame.

**285.** A mixture of  $\text{CaF}_2$ ,  $\text{Na}_2\text{B}_4\text{O}_7$ ,\* and  $\text{KHSO}_4$ , if finely powdered, moistened, and heated on a loop of platinum wire in the inner blowpipe flame, colors the outer flame momentarily *green* after being heated for a short time.

This test is not always decisive, since many phosphates and Cu salts give the same coloration, and it is besides liable to be masked by the flame colorations of other substances.

*Hydrogen borate*, or *boric acid* ( $\text{HBO}_3$ ), is a white porous mass; with  $2\text{H}_2\text{O}$  it forms a crystalline scaly hydrate; when strongly heated these yield a fusible glass consisting of  $\text{B}_2\text{O}_3$ . The solution of the acid reddens litmus-paper and *colors turmeric reddish-brown, especially on drying the paper*. The acid is soluble both in  $\text{H}_2\text{O}$  and in alcohol; on evaporating these solutions, the acid passes off with the vapor of the liquid. Reactions 283, 284, and 285 will serve to detect boric acid.

---

**CHROMATES** ( $-\text{CrO}_4$ ).—Use  $\text{K}_2\text{Cr}_2\text{O}_7$ .

All chromates are more or less yellow or red in color. The yellow solution of an alkaline chromate becomes orange-red on addition of an acid, owing to formation of a dichromate

**286.** Various substances, which readily combine with oxygen, deoxidize acid  $\text{K}_2\text{Cr}_2\text{O}_7$  solution to a solution of a chromic salt, the color of the solution at the same time changing from *orange-red* to a *bright green*. To show this add  $\text{HCl}$  or  $\text{H}_2\text{SO}_4$  to some  $\text{K}_2\text{Cr}_2\text{O}_7$  solution, divide the orange-red solution into five parts, and deoxidize each of these portions by one of the following methods:

- a. Pass  $\text{H}_2\text{S}$  into the hot solution; white S separates.
- b. Add some  $\text{NaHSO}_3$  or  $\text{H}_2\text{SO}_3$  and warm.

---



\* Boric acid is better, since the Na flame coloration tends to conceal the green coloration of  $\text{BF}_3$ .

c. Add alcohol and boil.

d. Boil for some time, after adding much strong HCl.

e. Add some Zn to the acidified liquid and warm. The change of color is in this case slow.

In each case the change in composition is shown by the liquid becoming green: the following equations show the decomposition occurring with  $H_2S$  and with HCl, the others may be drawn out by the student:



This green solution gives all the reactions for Cr (102-105).

*Note.*—Since chromic acid is thus reduced by boiling with HCl and by  $H_2S$  to a chromic salt, a chromate leads to precipitation of  $Cr_2H_2O_6$  in Group III in the ordinary course of analysis.

*Hydrogen peroxide* ( $H_2O_2$ ), if added after some ether to an acid solution of a chromate, on well shaking the liquids causes the ether to assume a blue color; addition of a few drops of strong  $HNO_3$  is useful. The  $H_2O_2$  may be produced by adding  $BaO_2$  to the *cold* acid liquid. This reaction is extremely delicate and characteristic.

**287.**  $AgNO_3$ : *purple-red* precipitate ( $Ag_2CrO_4$ ): soluble in  $AmHO$  and in  $HNO_3$ .

**288.**  $Pb\bar{A}_2$ : *yellow* precipitate ( $PbCrO_4$ ): soluble in  $KHO$ , insoluble in  $H\bar{A}$ .

**289.**  $BaCl_2$ : yellowish-white precipitate ( $BaCrO_4$ ): insoluble in  $H\bar{A}$ , soluble in HCl.

*Hydrogen chromate or chromic acid* ( $H_2CrO_4$ ?).  $CrO_3$  dissolves in water to a reddish-yellow liquid, with a strongly acid reaction. The liquid evolves  $Cl$ , and becomes green if it is boiled with HCl; when neutralized by an alkali it becomes yellow, the solution yielding the reactions in pars. 287, 288, 289.

---

SILICATES. { Use  $Na_2SiO_3$  solution for liquid;  
for solid, finely ground sand  
or solid  $Na_2SiO_3$ .

**290.** *HCl* added slowly drop by drop to a solution of  $\text{Na}_2\text{SiO}_3$  whilst stirring or shaking it, will usually, unless the solution is very dilute [see note 2 below], give a gelatinous precipitate of  $\text{H}_2\text{SiO}_4$ ; but this precipitate is very apt to remain dissolved, hence its non-appearance is no proof of the absence of  $\text{H}_2\text{SiO}_4$ .


**290 a.** Small quantities of  $\text{H}_2\text{SiO}_4$ , or  $\text{H}_4\text{SiO}_4$  in dilute solution, may be detected by evaporating the solution, made acid with *HCl*, to dryness in a porcelain dish, and heating the residue to  $100^\circ$  as long as acid fumes come off; on stirring and warming the dry residue with strong *HCl*,  $\text{SiO}_2$  remains undissolved as a white powder.

*Note 1.*—During the last part of the evaporation the silica often separates as a gelatinous mass, and unless the heat is lessened and the residue stirred with a glass rod, it will spirt; spirting is best avoided by finishing the evaporation, as soon as the liquid becomes thick, upon a water-bath. The undissolved  $\text{SiO}_2$  left on treating with *HCl* is best seen by pouring out the liquid into a test-tube; it is soluble in hot *KHO* solution.

*Note 2.*—The solution of  $\text{Na}_2\text{SiO}_3$  used for these reactions is made of such a strength that acids do not precipitate  $\text{H}_2\text{SiO}_4$  from it, but by boiling some of it down to a quarter its volume before adding the acid the precipitate is usually obtained.

**291.** *AmCl* (or  $\text{Am}_2\text{CO}_3$ ) causes a gelatinous precipitate of  $\text{H}_2\text{SiO}_4$ , which usually forms only after a time.

**292.** *Blowpipe Tests.*—A little solid  $\text{SiO}_2$  (or most silicates), if fused into a bead of  $\text{Na}_2\text{CO}_3$ , causes frothing from *CO* escaping:



**293.** Fused in a bead of  $\text{NaAmHPO}_4$ , the  $\text{SiO}_2$  is not dissolved, but floats about in transparent particles and on cooling remains as little opaque masses in the bead. The particles of  $\text{SiO}_2$  are usually best seen in the fused bead.

*Hydrogen silicate*, or *silicic acid* ( $\text{H}_4\text{SiO}_4$ ), is a white flocculent substance with no reaction to litmus; it is insoluble in water, and but very slightly soluble in acids, but dissolves as an alkaline silicate in solutions of the alkalis or alkaline carbonates. On ignition it forms a white insoluble powder ( $\text{SiO}_2$ ) which yields reactions 292 and 293.

FLUORIDES (-'F).—Use  $\text{CaF}_2$ , finely powdered.

*Note.*—The most useful tests for a fluoride depend upon setting free the acid and allowing it to act upon a surface of glass, which it 'etches' or corrodes, causing it to appear dim when perfectly dry.

294. Strong  $\text{H}_2\text{SO}_4$  if warmed, with a little  $\text{CaF}_2$  in a test-tube, liberates HF, which causes the glass to behave towards the liquid as if it were greasy. The HF etches the glass, and, on washing away the cooled mixture and perfectly drying the inside of the tube, it will be found to be dimmed and roughened.

This test is often performed by gently warming the substance, upon which several drops of strong  $\text{H}_2\text{SO}_4$  have been poured, upon a watch-glass, for some time; on washing off the substance and drying the glass thoroughly, it will be found to be dimmed and roughened where the substance and acid rested; this is best seen, if only very slight, by holding the glass so as to see the reflection from its surface.

295. A less simple, but more delicate, plan consists in placing the  $\text{CaF}_2$  in a platinum crucible, or in a small leaden cup made by bending a piece of sheet-lead, and pouring upon it some strong  $\text{H}_2\text{SO}_4$ . The crucible is then covered with a watch-glass with its hollow upwards and filled with cold water to keep the glass cool; the crucible is placed in a porcelain dish containing sufficient hot water to reach half way up the sides of the crucible, the water being occasionally heated, if necessary, by a small flame;\* on removing the watch-glass after ten minutes or more, washing and then thoroughly drying it, the part of the surface covering the crucible will be found to be dimmed by the action of the HF vapors. The delicacy of this method is much increased by the modification described in par. 296.

296. The convex surface of a watch-glass is covered with a thin film of beeswax or solid paraffin by carefully heating the glass held by crucible tongs at some

---


\* Or the crucible may be heated *gently* over a flame or on a sand-bath.

distance above a flame, then rubbing its convex side with a piece of wax or paraffin; it is thus covered with a melted layer, which may be made uniform by once more heating the glass, allowing any excess of wax to drop off, and keeping the glass moving whilst cooling; the coating is finally hardened by immersion for a short time in cold water. Lines or characters are then traced on the glass towards its centre by gentle pressure with the point of a penknife, the object being simply to remove the wax from these parts and not to scratch the glass. The glass thus prepared is exposed (as in par. 295) to the HF for about twenty minutes or half an hour, the wax being prevented from melting by cold water kept in the hollow of the glass. The watch-glass is then removed and cleansed from wax by holding it with crucible tongs above the flame and quickly wiping off the wax when melted; a clean bright surface will be thus obtained on which the traced lines will usually be visible at once on holding the glass up to the light, but if mere traces of a fluoride were present the etched lines will be made visible by breathing upon the cold surface.

297. The presence of  $\text{SiO}_2$ , or a silicate, prevents the liberation of HF by  $\text{H}_2\text{SO}_4$ ,  $\text{SiF}_4$  being given off instead; this gas possesses no etching power, and therefore if  $\text{SiO}_2$  or a silicate is present, the methods given above do not serve for the detection of a fluoride.  $\text{SiF}_4$  however produces white fumes in air, and if passed into dilute AmHO yields a colorless flocculent precipitate of  $\text{H}_4\text{SiO}_4$  and gives AmF in solution; the formation of the flocculent precipitate of  $\text{H}_4\text{SiO}_4$  is sufficient proof of the presence of a fluoride, but the AmF itself may be detected in the filtrate from this precipitate by adding  $\text{CaCl}_2$ , filtering off the precipitate of  $\text{CaF}_2$ , and drying and examining it by par. (296).

By mixing  $\text{SiO}_2$  with any substance to be tested for a fluoride the above method may be employed for its detection; use for the reaction a mixture of  $\text{CaF}_2$  and  $\text{SiO}_2$ . A small piece of marble may be dropped into the tube in which the  $\text{SiF}_4$  is evolved, to carry that gas over into the AmHO solution.

The changes which occur are thus expressed :



**298.**  $\text{CaCl}_2$  added to NaF solution\* gives a gelatinous almost transparent precipitate ( $\text{CaF}_2$ ), which becomes more visible on heating or adding AmHO; slightly soluble in HCl, almost insoluble in ( $\text{H}\bar{\text{A}}$ ).

*Hydrogen fluoride, or hydrofluoric acid (HF)*, is a colorless gas, which fumes in the air, etches glass and dissolves easily in water. The solution of HF is acid in reaction and differs from all other acids in dissolving  $\text{SiO}_2$  and insoluble silicates.

FLUOSILICATES ( $-\text{SiF}_6$ ).—Use  $\text{H}_2\text{SiF}_6$ .

**299.**  $\text{BaCl}_2$ : white precipitate ( $\text{BaSiF}_6$ ), which usually appears only on gently warming the solution, and thus differs from  $\text{BaSO}_4$ ; it is somewhat soluble in boiling HCl, but insoluble in alcohol. Put by some of this precipitate for reaction (301).

This precipitate differs from the precipitate of  $\text{BaSO}_4$ , which it somewhat resembles in being rather insoluble in HCl, by being more or less transparent and crystalline; but fluosilicates differ from sulphates also in giving no precipitates with  $\text{Sr}(\text{NO}_3)_2$  or  $\text{Pb}(\text{NO}_3)_2$  and in giving, when heated with excess of AmHO, a flocculent precipitate of  $\text{H}_4\text{SiO}_4$  and AmF in solution (297); fluosilicates differ also by the following very distinctive reactions:

**300.**  $\text{KCl}$ , on warming, gives a very transparent gelatinous precipitate ( $\text{K}_2\text{SiF}_6$ ), which is visible only after being allowed to settle for a time; it is insoluble in alcohol.

**301.** By evaporating  $\text{H}_2\text{SiF}_6$  solution, or heating a fluosilicate with strong  $\text{H}_2\text{SO}_4$ , HF is given off and may be detected by its etching action on glass (294).

Evaporate a little  $\text{H}_2\text{SiF}_6$  [or  $\text{BaSiF}_6$  from (299)],

\* Prepared by boiling in water the mass obtained by fusing for several minutes on platinum foil some powdered  $\text{CaF}_2$  and fusion-mixture, filtering the solution and acidifying the filtrate with  $\text{H}\bar{\text{A}}$ .



after adding strong  $\text{H}_2\text{SO}_4$ , upon a watch-glass to dryness; on washing off the residue the glass will be found to be etched.

**301 a.** A solid substance (*e.g.*, the precipitate of  $\text{BaSiF}_6$  or  $\text{K}_2\text{SiF}_6$ ) may also be examined by any of the methods given above for HF, the most delicate being (296). This method is particularly useful for proving a precipitate produced by  $\text{BaCl}_2$  to consist of  $\text{BaSiF}_6$ .\*

**302.** The above reactions readily distinguish between sulphates and fluosilicates, both of which give precipitates with  $\text{BaCl}_2$ , which are insoluble in boiling HCl. There is only one other class of salts—the selenates—which give with  $\text{BaCl}_2$  a white precipitate, insoluble in HCl; and this precipitate ( $\text{BaSeO}_4$ ), if boiled with strong HCl for some time, evolves Cl and gradually dissolves as  $\text{BaSeO}_3$ , thus differing from  $\text{BaSO}_4$  and  $\text{BaSiF}_6$ ; further, this HCl solution, if warmed with  $\text{H}_2\text{SO}_3$ , yields a reddish precipitate of Se.

*Hydrogen fluosilicate*, or *hydrofluosilicic acid* ( $\text{H}_2\text{SiF}_6$ ), is a strongly acid liquid, volatilizing entirely when heated in platinum as  $2\text{HF} + \text{SiF}_4$ ; by evaporating it in glass the glass is etched. It is precipitated by KCl and by  $\text{BaCl}_2$ .

### REACTIONS OF ORGANIC ACID RADICLES.

The following acid radicles are included here (although the first four are frequently, and the other three always, classed amongst organic compounds), because they are of such frequent occurrence, and their detection is usually a matter of so little difficulty that they may reasonably be appended even to an elementary treatise on inorganic analysis.

#### CYANIDES (—'Cy).—Use KCy.

HCy smells of bitter almonds, the alkaline cyanides smell only faintly of the same.

\* The readiest means of producing an etch with this precipitate is to pour it off into a watch-glass, decant the water, and warm the precipitate for some time with strong  $\text{H}_2\text{SO}_4$ ; on washing the glass and drying it, it will be found to be corroded. When the more delicate process (296) has to be employed, the precipitate should be shaken up with the liquid, poured off into a platinum crucible or leaden cup, the liquid decanted, and the precipitate warmed with strong  $\text{H}_2\text{SO}_4$  as described in (296).

**303.**  $AgNO_3$ ; white precipitate ( $AgCy$ ): show with three separate portions that this precipitate is soluble in  $KCy$  solution added in excess, and in  $AmHO$ , but insoluble in dilute  $HNO_3$ .

If some of the precipitate be filtered off and well washed with hot water until free from  $AgNO_3$  (shown by  $HCl$  causing no milkiness with a few drops of the washing water), then dried, scraped from the filter into a porcelain crucible, and heated strongly,  $Ag$  is left: if this residue on cooling be warmed with a little dilute  $HNO_3$ , it dissolves, giving off red fumes; and a drop of  $HCl$  added to this solution gives a curdy precipitate of  $AgCl$ . This reaction distinguishes  $AgCy$  from  $AgCl$ ,  $AgBr$  and  $AgI$ , which, like it, are precipitated by  $AgNO_3$ , and are insoluble in  $HNO_3$ , since these three precipitates when dried and strongly heated melt without decomposing; hence, after cooling, if  $HNO_3$  be warmed with them, it dissolves out no  $Ag$ , and  $HCl$  added to the  $HNO_3$  gives either no precipitate or a mere milkiness due to imperfect washing of the precipitate.

This method is somewhat tedious and need seldom be used, cyanides being readily detected by the following reactions.

**304.** If some  $FeSO_4$  solution, mixed with several drops of  $Fe_2Cl_6$ , be added to  $KCy$  solution, and the liquid be then made strongly alkaline with  $KHO$  and boiled, then cooled, and  $HCl$  added to acid reaction, "Prussian Blue" ( $Fe'''_4Cfy_3^*$ ) remains as a deep blue precipitate; or, if in small quantity, it dissolves giving a bluish-green solution. If no cyanide had been present, the addition of  $HCl$  would have given a nearly colorless liquid.

**305.** If to some  $KCy$  solution, contained in a test-tube fitted as shown in Fig. 37 (p. 149), dilute  $H_2SO_4$  is added, then a small piece of marble dropped in and the liquid boiled, allowing the steam to pass off into some  $Am_2S$  solution,  $AmCyS$  is formed. This substance gives with  $Fe_2Cl_6$ , a *blood-red* coloration, which may be obtained by pouring the  $Am_2S$  solution out into a porcelain dish, boiling it, allowing it to cool, making acid with  $HCl$ , and adding several drops of  $Fe_2Cl_6$ : a blood-red color, only faintly seen when but little  $KCy$  is used, shows that  $HCy$  has passed into the  $Am_2S$ , pro-

---

\* The symbol  $Cfy$  conveniently represents the group ( $FeCy_4$ ); it is used here to avoid brackets.

ducing AmCyS, and therefore proves the presence of a cyanide.

*Note.*—The marble is used to cause evolution of  $\text{CO}_2$  gas, which lessens the risk of the  $\text{Am}_2\text{S}$  being sucked back during the reaction.

A little  $\text{HgCl}_2$  solution should be added to a few drops of the red solution; if the red color is destroyed it was certainly due to the presence of AmCyS.

**306.** The former test can often be more simply performed by boiling the solution (of KCy or  $\text{HgCy}_2$ ) with  $\text{Am}_2\text{S}$ , which must be added until the yellow color remains after boiling for a short time. [If a dark-colored precipitate forms, this must be filtered off before the color of the filtrate can be seen.] Then filter if necessary, acidify with HCl in a porcelain dish, and add  $\text{Fe}_2\text{Cl}_6$  in small quantity; a blood-red color is produced in the liquid, which is not removed by heat or addition of HCl, but may be destroyed by adding to a few drops of the liquid some  $\text{HgCl}_2$ .

*Note 1.*—On adding HCl to the  $\text{Am}_2\text{S}$  (305, 306), white S separates, unless the yellow liquid has been previously boiled in a porcelain dish until it becomes colorless (filling in distilled water as it evaporates); but this separation of S is advantageous, since it helps to show by its whiteness even a faint reddish tint in the liquid. Methods (304, 305, 306) are exceedingly delicate tests for a cyanide. Methods (305) and (306), if less readily performed than (304), exceed it in delicacy. Method (306) is the more easy of the two to perform, and (305) is only used in cases where after boiling with excess of  $\text{Am}_2\text{S}$ , the filtrate is dark-colored, and therefore the red coloration would be invisible.

*Note 2.*— $\text{HgCy}_2$  does not show the reaction for a cyanide by methods (303, 304, and 305); but by boiling with  $\text{Am}_2\text{S}$ , according to method (306), AmCyS is obtained in the filtrate from the black HgS precipitate, and will give the red coloration with  $\text{Fe}_2\text{Cl}_6$ .

Solid  $\text{HgCy}_2$ , heated in a small tube closed at one end also gives off cyanogen gas, which burns with a peachblossom-colored flame.

*Hydrogen cyanide*, or *hydrocyanic* ("Prussic") acid, is a colorless, volatile, inflammable liquid, smelling like bitter almonds; it is *highly poisonous*, both when inhaled as vapor and swallowed as liquid; it readily mixes with water. The acid is easily detected by  $\text{AgNO}_3$  (303), also by (304), or by formation of AmCyS when the liquid is boiled with  $\text{Am}_2\text{S}$  (306).

FERROCYANIDES ( -  $'''FeCy_6$ ,\* or -  $'''Cfy$ ). — Use  $K_4FeCy_6 \cdot 3H_2O$ , ( $K_4Cfy \cdot 3H_2O$ ).

307.  $Fe_2Cl_6$ : dark-blue precipitate, "Prussian Blue" ( $Fe_4Cfy_3$ ); insoluble in HCl, soluble in  $H_2C_2O_4$  to a dark-blue liquid; changed by KHO into brown  $Fe_2Ho_6$ .

*Note.*—The solubility in  $H_2O$  and insolubility in HCl are shown by pouring off portions, heating them with HCl and  $H_2O$  respectively, filtering, and noting whether the filtrate is blue; a third portion is mixed with KHO, and becomes brown.

308.  $FeSO_4$ : light-blue precipitate ( $Fe''K_2Cfy$ ), which quickly becomes darker on standing in the air, or immediately on addition of Cl-water or Br-water, or on warming with  $HNO_3$ . It is insoluble in HCl.

309.  $CuSO_4$ : a chocolate-colored precipitate ( $Cu''Cfy$ ), insoluble in  $H\bar{A}$ .

310.  $AgNO_3$ : white precipitate ( $Ag_4Cfy$ ), insoluble in  $HNO_3$  and in  $AmHO$ . If heated with  $HNO_3$  this precipitate changes to orange-red  $Ag_6Cfy$ , soluble in  $AmHO$ .

FERRICYANIDES [ -  $^I(FeCy_6)_2$ , or -  $^ICfy_2$ ]. — Use  $K_6(FeCy_6)_2$ , ( $K_6Cfy_2$ ).

311.  $Fe_2Cl_6$ : dark-greenish or brown coloration; no precipitate is formed, as may be seen by adding much water to the liquid.

312.  $FeSO_4$ : dark-blue precipitate, "Turnbull's Blue," ( $Fe''_3Cfy_2$ ); insoluble in acids; blue color destroyed by KHO.

313.  $CuSO_4$ : yellowish precipitate.

314.  $AgNO_3$ : orange-colored precipitate ( $Ag_6Cfy_2$ ), insoluble in  $HNO_3$ , soluble in  $AmHO$ . Frequently a white residue is left on adding  $AmHO$ , consisting of  $Ag_4Cfy$ , and showing that  $K_4Cfy$  was present in the  $K_6Cfy_2$  solution.

\* The latter part of this formula is conveniently written  $Cfy$  to avoid using brackets.

SULPHOCYANIDES ( $- \text{'CyS}$ ).—Use  $\text{KCys}$ .


**315.**  $\text{Fe}_2\text{Cl}_6$ : blood-red coloration, but no precipitate; the color is destroyed by  $\text{HgCl}_2$  solution, but not by  $\text{HCl}$ .

**316.** Cyanides are usually detected in analysis by methods (304, 305, 306). Ferro- and ferricyanides can be found and distinguished by their first three tests, or by the precipitate given by  $\text{AgNO}_3$  in the solution acidified with  $\text{HNO}_3$ ; this precipitate, if a ferricyanide is present, will have a more or less decided orange-red color; on filtering, or decanting, and warming the precipitate with  $\text{AmHO}$ , white  $\text{Ag}_4\text{Cfy}$  will be left if a ferrocyanide was present, and on acidifying the filtrate with  $\text{HNO}_3$  orange-red  $\text{Ag}_6\text{Cfy}_2$  will precipitate. A sulphocyanide is found by (315).

The three following acid radicles and their salts differ from the foregoing by being charred and emitting a smell of burning when heated in the solid state; oxalates, however, do so only to a slight extent. If the salt be one of the alkalies or of the alkaline earths, the residue left after ignition contains a carbonate which effervesces with an acid, whereas the original substance did not do so.

OXALATES ( $- \text{'C}_2\text{O}_4$ , or  $- \text{'O}$ ).—Use  $\text{Am}_2\text{C}_2\text{O}_4 \cdot 2\text{H}_2\text{O}$ , ( $\text{Am}_2\text{O} \cdot 2\text{H}_2\text{O}$ ).

**317.** Strong  $\text{H}_2\text{SO}_4$  heated with solid  $\text{Am}_2\text{O} \cdot 2\text{H}_2\text{O}$  causes an effervescence of gas consisting of a mixture of  $\text{CO}$  and  $\text{CO}_2$ :



The  $\text{CO}$  is detected by burning with a blue flame when a light is applied, the  $\text{CO}_2$  by rendering a drop of lime-water on a glass rod milky.


No blackening or charring is produced by strong  $\text{H}_2\text{SO}_4$ ; oxalates differ in this respect from the salts of all other organic acids.

**318.**  $\text{CaCl}_2$  ( $\text{CaSO}_4$ , or lime-water), added to  $\text{Am}_2\text{C}_2\text{O}_4$  solution (made by diluting some of the reagent solution in a test-tube with an equal quantity of water), gives a white precipitate ( $\text{CaC}_2\text{O}_4$ ), insoluble in  $\text{AmHO}$  and in  $\text{HA}$ , soluble in  $\text{HCl}$  or  $\text{HNO}_3$ .

**318 a.** If this precipitate is filtered off, a portion placed on a watch-glass will not effervesce with dilute HCl; but if the remainder is dried and heated to dull redness on a platinum foil, it is converted without blackening into  $\text{CaCO}_3$ , which effervesces with dilute HCl.

*Note.*—This is a general test for oxalate of K, Na, Ba, Sr, or Ca; oxalates of these metals leave *carbonates* on ignition which effervesce with an acid.

**319.** Heated with  $\text{MnO}_2$  and  $\text{H}_2\text{SO}_4$ , solid oxalates give off  $\text{CO}_2$ :



*Note.*—Since commercial  $\text{MnO}_2$  usually contains some carbonate, it will give off  $\text{CO}_2$  when treated with  $\text{H}_2\text{SO}_4$ . If the  $\text{MnO}_2$  to be used is not known to be free from carbonate the best way to perform the test is to add to some of the  $\text{MnO}_2$  in a test-tube a little  $\text{H}_2\text{SO}_4$ , and heat to boiling; if  $\text{CO}_2$  is found in the tube by a glass rod with a drop of lime-water on its end continue to boil, occasionally drawing fresh air into the test-tube by pushing down a glass tube near to the surface of the liquid and sucking out the air; as soon as no  $\text{CO}_2$  is detectable in the tube after boiling for a short time, and the liquid is distinctly acid, the  $\text{Am}_2\text{C}_2\text{O}_4$  is added, and the mixture again heated,  $\text{CO}_2$  will be rapidly given off, and may be detected by the ordinary methods.

If the  $\text{MnO}_2$  gives no  $\text{CO}_2$  with  $\text{H}_2\text{SO}_4$  proceed at once to add the  $\text{Am}_2\text{C}_2\text{O}_4$ , heat and test for  $\text{CO}_2$ .

*Hydrogen oxalate*, or *oxalic acid* ( $\text{H}_2\text{C}_2\text{O}_4$ , or crystallized  $\text{H}_2\text{C}_4\text{O}_4 \cdot 2\text{H}_2\text{O}$ ), when heated in a glass tube closed at one end partly sublimes unaltered, and is partly decomposed, yielding vapors which cause coughing; it yields reactions in pars. (317) and (319). It dissolves in water to an acid, poisonous liquid.

TARTRATES.  $\left\{ \begin{array}{l} \text{For a neutral tartrate use } \text{Na}_2\bar{\text{T}}. \\ 2\text{H}_2\text{O or } \text{KNa}\bar{\text{T}}. \text{ For an acid} \\ (-\text{''C}_4\text{H}_4\text{O}_6, \text{ or } -\text{''}\bar{\text{T}}). \left\{ \begin{array}{l} \text{tartrate use } \text{H}_2\bar{\text{T}} \text{ or } \text{NaH}\bar{\text{T}}. \end{array} \right. \end{array} \right.$

**320.** A solid tartrate or solid  $\text{H}_2\bar{\text{T}}$  when heated strongly in a tube chars, giving off a smell of burnt sugar.

**321.** A solid tartrate or solid  $\text{H}_2\bar{\text{T}}$  heated with strong

$\text{H}_2\text{SO}_4$  blackens, giving off a mixture of  $\text{CO}$ ,  $\text{CO}_2$ , and  $\text{SO}_2$ .

**322.**  $\text{AgNO}_3$  gives with solution of a neutral tartrate a white curdy precipitate ( $\text{Ag}_2\bar{\text{T}}$ ), soluble in  $\text{AmHO}$  and in  $\text{HNO}_3$ . Allow some of the above precipitate ( $\text{Ag}_2\bar{\text{T}}$ ) to settle in a test-tube, and pour off as much of the liquid from above it as possible; then pour very dilute  $\text{AmHO}$  drop by drop into the tube until the precipitate, after being shaken up with the liquid, is nearly but *not quite* dissolved: then heat the test-tube gradually and gently (best by immersing it in a beaker of cold water, and slowly raising its temperature to  $60^\circ$ ), a brilliant mirror of  $\text{Ag}$  will form on the interior of the tube. If the liquid, instead of being slowly raised to  $60^\circ$ , be quickly boiled, the  $\text{Ag}$  deposits as a black precipitate, which is not so characteristic.

**323.**  $\text{CaCl}_2$  precipitates from a neutral tartrate solution white or crystalline  $\text{Ca}\bar{\text{T}}$ : the precipitate often separates only after shaking the liquid well and letting it stand for a time; it is soluble in acids, and its formation is retarded by  $\text{AmCl}$  and other  $\text{Am}$  salts. This precipitate is soluble in strong cold  $\text{KHO}$ ,\* is reprecipitated by boiling, but dissolves again more or less completely on cooling.

Decant the liquid from the precipitate and pour upon the precipitate a little very dilute  $\text{AmHO}$  in a test-tube, drop in a crystal of  $\text{AgNO}_3$  and warm carefully as in par. (322). A silver mirror forms on the part of the glass on which the  $\text{AgNO}_3$  rests.

**324.**  $\text{KCl}$  (or better  $\text{K}\bar{\text{A}}$ ) if added to solution of  $\text{H}_2\bar{\text{T}}$  or an acid tartrate, gives on stirring or shaking a white crystalline precipitate ( $\text{KHT}$ ). If  $\text{H}\bar{\text{A}}$  is added at the same time, the precipitate may also be obtained with neutral tartrates. Addition of alcohol assists the formation of the precipitate, which is soluble in alkalies and in mineral acids.

---

\* A milkiness remaining, however, if the potash solution contains carbonate.

**325.** If a few drops of  $\text{Fe}_2\text{Cl}_6$  solution be added to  $\text{H}_2\bar{\text{T}}$  or  $\text{KNa}\bar{\text{T}}$  solution, then  $\text{KHO}$  (or  $\text{AmHO}$ ) until the liquid is strongly alkaline, no precipitate forms. The precipitation of several other metals (*e. g.*,  $\text{Al}$ ,  $\text{Mn}$ ,  $\text{Cu}$ ) by alkalies is similarly prevented by  $\text{H}_2\bar{\text{T}}$ : this is owing to the formation of a double tartrate of the alkali and the other metal, which salt is not decomposed by an alkali.

*Hydrogen tartrate*, or *tartaric acid* ( $\text{H}_2\bar{\text{T}}$ ), occurs in colorless crystals which do not alter in the air; they taste pleasantly acid, and dissolve in water and in alcohol. They are detected by heating them alone or with strong  $\text{H}_2\text{SO}_4$  (320, 321): if dissolved and neutralized, they also yield the reactions in (322, 323).

ACETATES ( $-\text{'C}_2\text{H}_3\text{O}_2$ , or  $-\bar{\text{A}}$ ).—Use  $\text{Na}\bar{\text{A}}.3\text{H}_2\text{O}$ .

**326.** Solid  $\text{Na}\bar{\text{A}}$  if strongly heated blackens, giving off a peculiar smell and a vapor (acetone) which burns with a bright flame when kindled.

**327.** Heated with strong  $\text{H}_2\text{SO}_4$  vapor of  $\text{H}\bar{\text{A}}$  is given off, recognized by its pungent smell (that of vinegar). If alcohol be first added to the  $\text{Na}\bar{\text{A}}$  then strong  $\text{H}_2\text{SO}_4$ , and the mixture warmed, a fragrant smell of ethyl-acetate ( $\text{Et}\bar{\text{A}}$ ) is emitted; it is most decided when the liquid is shaken after cooling for a short time.

**328.**  $\text{Fe}_2\text{Cl}_6$  added in small quantity to a little  $\text{Na}\bar{\text{A}}$  dissolved in water, gives a deep-red color: on boiling the solution made neutral but not alkaline with  $\text{AmHO}$  (if necessary), all the  $\text{Fe}$  separates as a light-brown precipitate and the liquid becomes colorless. The red color of the solution before being boiled is changed to yellow by addition of  $\text{HCl}$  (difference from the coloration due to sulphocyanide).

*Hydrogen acetate* or *acetic acid* ( $\text{H}\bar{\text{A}}$ ).—"Glacial acetic acid" is at  $17^\circ \text{C}$ . a colorless crystalline substance, possessing (especially when heated) a very pungent smell,



and giving inflammable vapor: it is strongly acid. It dissolves in water readily, imparting its peculiar smell to the liquid in a mild degree. The acid is completely volatile without decomposition.

---

**329.** The above organic acid radicles may be specially tested for by any of their reactions; (318) being generally used to test for an oxalate in solution, (317) or (319) when it is solid; a tartrate is best detected in solution by reaction (322) or (323), or if solid by (320); an acetate is found by (326) or (327).

---

\* \* \* After completing the reactions for the acid radicles the student may proceed to try the preliminary examinations of solid substances for metals and acid radicles, as directed in paragraphs (375-377), unless he is intending only to learn the analysis of simple salts, when he proceeds to Section V (330 *et seq.*).

.

## SECTION V.

**ANALYSIS OF SIMPLE SUBSTANCES CONTAINING  
ONE METAL AND ONE ACID RADICLE.\***

*Note.*—This section may be neglected by a student who is proceeding through the senior analytical course.

**330.** The ordinary method of proceeding to detect the metal or acid radicle or both in a simple substance, is to make a few preliminary experiments, which will usually give some idea as to what substance is present, and will occasionally even detect metal or acid radicle, or both of them, with certainty. If the composition of the substance has not been satisfactorily proved by this preliminary examination, the next process is to endeavor to dissolve the substance, if it is not already in the liquid form, in water or acids; then by adding to the solution certain group-reagents in the proper order of succession, it is found to which Analytical Group the metal present belongs. The following table gives the analytical groups for metals, each group being headed by its distinctive number, name, and group-reagent :

---

\* Two metals may occasionally be present, as when an alum, or tartar emetic [ $\text{K}(\text{SbO})\bar{\text{T}}$ ], is given for analysis; also there may be two acid radicles in the liquid, given for analysis, since a salt insoluble in water is occasionally dissolved in an acid; but the student should always be informed of this if it in any way complicates the analysis. It will not usually do so, since in the most commonly occurring case, that of an alkaline earth phosphate dissolved in  $\text{HCl}$ , the  $(\text{---}\text{PO}_4)$  is detected in Group III, and accordingly no other acid radicle is looked for.

TABLE SHOWING THE ANALYTICAL CLASSIFICATION OF THE METALS.

*Note*—It will be noticed that there are two divisions of Groups II and III. In analysis it is usual to separate the two divisions of Group II, after having separated the whole group from the remaining groups; the two divisions of Group III are by some analysts precipitated together and subsequently separated; they are, however, often precipitated separately, and treated as distinct groups.

Metals whose chlorides are insoluble in water and dilute acids, and which are therefore precipitated from their solutions by hydrochloric acid.	Metals whose sulphides are insoluble in dilute acids, and which are therefore precipitated from acid solutions by sulphuretted hydrogen.	Metals which are precipitated, either as hydrates or as sulphides, from their solutions made alkaline with ammonia, by ammonium sulphide or sulphuretted hydrogen.	Metals whose carbonates are precipitated by addition of ammonium carbonate.	Metals whose solutions are not precipitated by any of the foregoing group-reagents in the presence of ammonium chloride.
<div>GROUP I. <i>Hydrochloric Acid Group.</i> <i>Silver Group.</i></div>	<div>GROUP II. <i>Sulphuretted Hydrogen Group.</i></div> <div><div>GROUP II A. <i>Copper Group.</i> Sulphides insoluble in NaHO and in <math>\text{Am}_2\text{S}</math>.</div><div>GROUP II B. <i>Arsenic Group.</i> Sulphides soluble in NaHO and in <math>\text{Am}_2\text{S}</math>.</div></div>	<div>GROUP III. <i>Ammonia and Ammonium-Sulphide Group.</i></div> <div><div>GROUP III A. <i>Ammonia Group.</i> <i>Iron Group.</i></div><div>GROUP III B. <i>Ammonium Sulphide Group.</i> <i>Zinc Group.</i></div></div>	<div>GROUP IV. <i>Ammonium Carbonate Group.</i> <i>Barium Group.</i></div>	<div>GROUP V. <i>Potassium Group.</i></div>
[Lead, . . . . (Pb)] Silver, . . . . (Ag) Mercurous, . . (Hg)	2. Mercuric, (Hg) Lead, . . . . (Pb) Bismuth, . . (Bi) Copper, . . . (Cu) Cadmium, (Cd)	3. Tin, . . . . (Sn) Arsenic, . . (As) Antimony, (Sb)	4. Aluminium, (Al) Iron, . . . . (Fe) Chromium, (Cr)	5. Zinc, . . . . (Zn) Manganese, (Mn) Nickel, . . . (Ni) Cobalt, . . . (Co)
			6. Barium, . . . (Ba) Strontium, . . (Sr) Calcium, . . . (Ca)	7. Ammonium, . . (NH <sub>4</sub> ) Potassium, . . (K) Sodium, . . . (Na) Magnesium, . . (Mg)

After discovering to which analytical group the metal present belongs, it is identified by the properties or appearance of the group-precipitate, and its presence confirmed, if necessary, by special tests made on the precipitate or on another portion of the "original solution." \*

The examination for the acid radicle, if it has not been already detected during the examination for the metal, is carried out in a less systematic manner. A few preliminary experiments are first performed, and if these do not detect the acid radicle, it is found, partly by a systematic examination including that described above for the detection of the metal, but more commonly by special tests.

A substance insoluble in water and in acids is examined according to the special directions in (367-369).

Since in these analyses only one metal and one acid radicle are to be looked for, the examination for the metal is carried no further when one metal has been detected with certainty, and the examination for the acid radicle is similarly at an end when the presence of one such radicle has been satisfactorily proved. It must, however, be understood that substances given for analysis are liable to contain impurities which will yield only slight corresponding precipitates; if in the course of analysis, therefore, *only a slight precipitate* is obtained, proceed until some substance is detected in larger quantity, entering against the small precipitate "Trace of —."

The separation and washing of a precipitate by "decantation" (23 *b*, 24 *b*) is recommended wherever it is possible, as this process is more easily performed than that of filtration (23 *a*): if a precipitate, however, refuses to settle after heating and shaking the liquid, filtration must usually be resorted to.

---

\* By the "original solution" is meant the solution of the substance, or the substance itself if a liquid is under analysis, to which no reagent has been added.

For examples of substances to be given for analysis by Section V, see par. 537.

\* \* *Note*.—The numbers in brackets refer to the paragraphs in the text in which tests or processes to be employed by the student are fully described.

*Caution*.—In every analysis each result must be fully entered in the note-book *as soon as the experiment has been performed*. The form of entry being the same as that adopted for the directions in the text.

The substance given for analysis may be either a solid or a liquid. If it is a solid examine by A (331 *et seq.*), if a liquid by B (334 *et seq.*).

---

#### A.—THE SUBSTANCE GIVEN FOR ANALYSIS IS A SOLID.

**331.** Make a careful examination of the properties of the substance, noting down its color, its smell if it possesses any; also its form, whether it is in powder, crystals, or non-crystalline pieces. Note carefully also any appearance or property which may strike you on examining the substance with the naked eye or with the assistance of a lens.

If the substance possesses a metallic lustre or appearance, work as directed under (366); if it is not metallic in appearance, proceed to ascertain its solubility by the directions contained in par. 332, and, according to the result thus obtained, conduct the further analysis of the substance by par. 333 or by par. 367.

## SOLUBILITY OF THE SUBSTANCE.

**332.** Since the method of examination of a solid substance depends upon whether it is soluble or insoluble in water or acids, the following experiments are tried.

Place a small quantity of the finely powdered substance in a broad test-tube or a boiling-tube, fill the tube about one-third with water, and heat to boiling:

1. <i>The substance dissolves.</i>	<i>The substance does not dissolve:</i> heat a fresh portion with strong HCl to boiling, if it does not dissolve, add water and boil:		
After examining a portion of the solid substance by the preliminary tests (333) proceed to examine the color and reaction of the liquid to litmus by (334, I, II), and refer to (335) for directions for analysis. The examination for the acid radicle will be found in paragraphs (346-358 b).	2. <i>The substance dissolves.</i> (See note below.)	<i>The substance does not dissolve:</i> put aside the test-tube containing the substance with HCl, and boil a fresh portion of the substance with strong HNO <sub>3</sub> ; if it is not dissolved, add water and boil:	
	<i>Absence of Group I.</i>	3. <i>The substance dissolves.</i>	<i>The substance does not dissolve:</i> mix together the contents of the two tubes in which the substance was heated with HCl and with HNO <sub>3</sub> , and heat the mixture.
	After examining a portion of the solid substance by the preliminary tests (333) proceed to examine the HCl solution (after noting its color, see 334, I) for the metal by (336-345) Examine a fresh portion for the acid radicle by (343 <i>et seq.</i> ).  [Note.—The results mentioned in (346) may be here obtained, if any one of them occurs it should be noted down; observe especially whether any effervescence occurs with HCl.]	After examining a portion of the solid substance by the preliminary tests (333) examine the HNO <sub>3</sub> solution (after noting its color, see (334, I) for the metal by (336 <i>et seq.</i> ) taking care after testing for Group I by HCl to boil down considerably in a porcelain dish after adding a little strong HCl,* and then add dilute HCl before testing for Group II by H <sub>2</sub> S.	4. <i>The substance dissolves.</i>
		After examining a portion of the solid substance by the preliminary tests (333 <i>et seq.</i> ) examine the solution in HNO <sub>3</sub> and HCl mixed, as directed in the preceding column (332, 3) for the metal, and a fresh portion of the substance for acid radicles by (346 <i>et seq.</i> ).	5. <i>The substance does not dissolve.</i>  Examine a fresh portion of the substance by (367).

\* This method of proceeding removes HNO<sub>3</sub>, which would cause with H<sub>2</sub>S a deposit of S.

**THE SUBSTANCE IS SOLUBLE IN WATER OR ACIDS.**

**333. PRELIMINARY EXAMINATION.**

The table is thus arranged: in the first column is placed the experiment to be made; in the second a list of the results, any one or more of which may be observed; and in the third the inferences to be drawn from the results. Confirmatory or supplementary experiments are occasionally entered.

A more complete preliminary examination will be found in (387-408).

Experiment.	Observation.	Inference.														
I. Heat a portion of the substance in an ignition-tube, or in a small dry test-tube.	1. The substance does not change, . . .	Absence of volatile, fusible and organic substances, and of water.														
	2. The substance changes color:															
	<table><tr><td><i>Ye low,</i></td><td>hot.</td><td><i>White,</i></td><td>cold.</td></tr><tr><td><i>Yellow-brown,</i></td><td>"</td><td><i>Yellow,</i></td><td>"</td></tr><tr><td><i>Dark red or</i></td><td rowspan="2">}</td><td><i>Red brown,</i></td><td>"</td></tr><tr><td><i>brown.</i></td><td></td><td></td></tr></table>	<i>Ye low,</i>	hot.	<i>White,</i>	cold.	<i>Yellow-brown,</i>	"	<i>Yellow,</i>	"	<i>Dark red or</i>	}	<i>Red brown,</i>	"	<i>brown.</i>		
<i>Ye low,</i>	hot.	<i>White,</i>	cold.													
<i>Yellow-brown,</i>	"	<i>Yellow,</i>	"													
<i>Dark red or</i>	}	<i>Red brown,</i>	"													
<i>brown.</i>																
	3. The substance fuses easily, . . .	Probable presence of a salt of Na or K, possibly also of Ba, Sr, Ca, Mg.														
	4. The substance sublimes, . . .	Presence of a salt of NH <sub>4</sub> , As, or Hg; possibly also of H <sub>2</sub> C <sub>2</sub> O <sub>4</sub> , of S (yellow), and of I (purple vapor). NH <sub>4</sub> is found by the following confirmatory test,—As is detected in Exp. V, Hg may be found by (147), or its detection may be omitted here.														
<i>Confirmatory.</i> —If a sublimate forms, boil another portion of the substance with KHO solution.	A smell of NH <sub>3</sub> is evolved, and white fumes appear when a rod moistened with strong HCl is held to the mouth of the tube.	Presence of NH <sub>4</sub> . If NH <sub>4</sub> is found here, the examination for the metal need not be further continued.														
	5. The substance blackens, and gives off CO <sub>2</sub> , which makes a drop of lime-water on a glass rod milky.	Presence of an organic acid radicle, probably A or T or O; the last is known by a smell of burnt sugar.														
	If the cool residue in the tube effervesces with HCl, whereas the original substance did not.	The organic acid radicle is united with K, Na, Ba, Sr, Ca, or Mg.														

PRELIMINARY EXAMINATION—*Continued.*

Experiment.	Observation.	Inference.
	<p>6. Gas or vapor is given off: smell the vapor:  Red nitrous fumes, recognized by their smell, .  <math>\text{SO}_2</math>, known by its smell and by turning <math>\text{K}_2\text{Cr}_2\text{O}_7</math> solution green.</p> <p>Cl, Br, I, known by smell and color: Br and I by action on starch-paste, and Cl by bleaching moist litmus.  A slip of wood with a spark at its end, glows brightly or bursts into flame.</p>	<p>From nitrate of a heavy metal.  From combustion of free S or a sulphide; possibly also from a sulphate or sulphite.  From certain chlorides, bromides, or iodides, also free I.  O from chlorate or nitrate, or <math>\text{N}_2\text{O}</math> from decomposition of <math>\text{NH}_4\text{NO}_3</math>.</p>
II. Dip into the powdered substance a moistened loop of platinum-wire, and hold the loop in the Bunsen flame; moisten with strong HCl, and again hold in the flame.	<p>1. The flame is not colored, . . . . .</p> <p>2. The flame is colored:  <i>Intense yellow,</i>  <i>Pale-violet,</i> appearing <i>violet-red</i> through the indigo-prism,  <i>Red,</i> { Appearing <i>green</i> through the indigo-prism,            { Appearing <i>intense red</i> through the indigo-prism, . . . . .  <i>Yellowish-green,</i>  <i>Green,</i> { <i>Bright-green,</i> with blue centre after moistening with HCl,            { <i>Green,</i> . . . . .            { <i>Pale-blue</i> (livid), . . . . .  <i>Blue,</i> { <i>Vivid-blue,</i> . . . . .</p> <p>The gas burns at the mouth of the tube; note the color of the flame:  <i>Pale grayish-green,</i> . . . . .</p> <p><i>Bright, white,</i> . . . . .</p> <p><i>Purple,</i> . . . . .</p>	<p>Probable absence of the substances enumerated below.</p> <p>Presence of Na.  " K.  " Ca.  " Sr.  " Ba.  " Cu.  " <math>\text{B}_2\text{O}_3</math>.  " As, Sb, Pb.  " <math>\text{CuCl}_2</math>.</p> <p>Presence of <math>\text{NH}_4</math>, probably from strongly heated <math>\text{NH}_4\text{NO}_3</math>.  Presence of <math>\text{PH}_3</math>, probably from a hypophosphite; the unburnt gas smells of garlic.  Presence of Cy from a cyanogen compound, probably <math>\text{HgCy}_2</math>.</p>
III. Heat a portion of the substance in a cavity scooped in charcoal, in the blow-pipe flame.	<p>1. The charcoal "deflagrates," or burns rapidly.  2. A white mass is left in the charcoal, which does not fuse, and shines brightly when strongly heated.</p>	<p>Presence of a nitrate or chlorate.  Presence of Ba, Sr, Ca, Mg, Al, or Zn.</p>



PRELIMINARY EXAMINATION—*Continued.*

Experiment.	Observation.	Inference.														
<i>Confirmatory.</i> — Detach a small portion of the unfused white mass, place it on red litmus-paper and moisten it with a drop of water.	The red paper becomes blue in a short time where the substance rests.	Presence of Ba, Sr, Ca, or possibly of Mg.														
<i>Confirmatory.</i> —If the moist residue is not alkaline to test-paper, moisten the remainder on the charcoal with $\text{Co}(\text{NO}_3)_2$ solution and heat again strongly in the blowpipe flame.	A blue residue, . . . . .	Presence of Al, or possibly of a phosphate, silicate, or borate.														
	A pink residue, . . . . .	Presence of Mg.														
	A green residue, . . . . .	" Zn.														
	3. The residue is not white, and does not consist of globules of metal; neither is there any incrustation formed upon the charcoal; proceed to Exp. IV.															
	4. The residue contains metallic globules or an incrustation forms on the charcoal; proceed to Exp. V, omitting Exp. IV.															
IV. Fuse a small quantity* of the substance into a clear borax bead first in the inner then in the outer flame, noting in both cases the color of the bead whilst hot and when cold.	The bead is colorless when heated in both flames.	Absence of the metals below.														
	The bead is colored:															
	<table border="1"> <thead> <tr> <th>In outer flame.</th><th>In inner flame.</th></tr> </thead> <tbody> <tr> <td>Blue, hot; green, cold.</td><td>Red or colorless.</td></tr> <tr> <td>Blue, hot and cold.</td><td>Blue, hot and cold.</td></tr> <tr> <td>Brown or dingy purple, hot; light-brown, cold.</td><td>Gray or black, hot and cold, . . . }</td></tr> <tr> <td>Brown, hot; yellow, cold.</td><td>Bottle-green, hot and cold, . . . }</td></tr> <tr> <td>Green,† hot and cold.</td><td>Green, hot and cold, . . . }</td></tr> <tr> <td>Purple, hot and cold.</td><td>Colorless, hot and cold, . . . }</td></tr> </tbody> </table>	In outer flame.	In inner flame.	Blue, hot; green, cold.	Red or colorless.	Blue, hot and cold.	Blue, hot and cold.	Brown or dingy purple, hot; light-brown, cold.	Gray or black, hot and cold, . . . }	Brown, hot; yellow, cold.	Bottle-green, hot and cold, . . . }	Green,† hot and cold.	Green, hot and cold, . . . }	Purple, hot and cold.	Colorless, hot and cold, . . . }	Presence of Cu. " Co. " Ni. " Fe.† " Cr. " Mn.
In outer flame.	In inner flame.															
Blue, hot; green, cold.	Red or colorless.															
Blue, hot and cold.	Blue, hot and cold.															
Brown or dingy purple, hot; light-brown, cold.	Gray or black, hot and cold, . . . }															
Brown, hot; yellow, cold.	Bottle-green, hot and cold, . . . }															
Green,† hot and cold.	Green, hot and cold, . . . }															
Purple, hot and cold.	Colorless, hot and cold, . . . }															
<i>Confirmatory.</i> — The presence of Cr or Mn may be confirmed by fusing a portion of the substance on platinum foil or wire with $\text{Na}_2\text{CO}_3$ and $\text{KNO}_3$ .	A yellow mass on cooling, . . . . .	" Cr.														
	A bluish-green mass on cooling, . . . . .	" Mn.														

\* It is necessary to adjust carefully the quantity of substance fused into the bead; if too much is used, the bead will often appear opaque; if too little is employed, one of the metals may escape detection. It is best at first to fuse only a minute quantity into the bead, adding more if no color, or but a faint color, is produced.

† Cr if present as a chromate gives a bead brown whilst hot, after having been fused in the outer blowpipe flame; but this brown color is not reproduced in the case of Cr when the bead, after becoming green by being heated in the inner flame, is again heated in the outer flame.

PRELIMINARY EXAMINATION—*Continued.*

Experiment.	Observation.	Inference.																				
<p>V. Mix the substance in a small cavity on wood-charcoal with <math>\text{Na}_2\text{CO}_3</math> and <math>\text{KCy}</math>, and heat strongly in the inner blowpipe flame for several minutes.</p> <p><i>Confirmatory.</i>—If metallic globules are obtained, detach one and strike it sharply on the bottom of an inverted mortar with the pestle, and note whether it is crushed to powder (<i>brittle</i>) or merely flattened out into a cake (<i>malleable</i>).</p> <p>If the globules are white and malleable, take one upon the point of a pen-knife and see if it will mark paper as a black lead-pencil does.</p>	<p>Metallic scales or globules are obtained, or an incrustation on the charcoal:</p> <table> <tr> <th><i>Globules.</i></th> <th><i>Incrustation.</i></th> </tr> <tr> <td>White and brittle.</td> <td>White, . . . .</td> </tr> <tr> <td>White and brittle.</td> <td>Yellow, . . . .</td> </tr> <tr> <td>Red and malleable.</td> <td>None, . . . .</td> </tr> <tr> <td>White and malleable marking paper easily.</td> <td>Yellow, . . . .</td> </tr> <tr> <td>White and malleable not marking paper, and readily fusible.</td> <td>None, . . . .</td> </tr> <tr> <td>White and malleable not marking paper, fusible only with difficulty.</td> <td>None, . . . .</td> </tr> <tr> <td>None, . . . .</td> <td> <div> Yellow whilst hot, white when cold; becomes green when moistened with <math>\text{Co}(\text{NO}_3)_2</math> solution and reheated in the outer flame. </div> </td> </tr> <tr> <td>None, . . . .</td> <td>Brown, . . . .</td> </tr> <tr> <td>None, . . . .</td> <td>White; on smelling the charcoal, a smell of onion is perceived.</td> </tr> </table>	<i>Globules.</i>	<i>Incrustation.</i>	White and brittle.	White, . . . .	White and brittle.	Yellow, . . . .	Red and malleable.	None, . . . .	White and malleable marking paper easily.	Yellow, . . . .	White and malleable not marking paper, and readily fusible.	None, . . . .	White and malleable not marking paper, fusible only with difficulty.	None, . . . .	None, . . . .	<div> Yellow whilst hot, white when cold; becomes green when moistened with <math>\text{Co}(\text{NO}_3)_2</math> solution and reheated in the outer flame. </div>	None, . . . .	Brown, . . . .	None, . . . .	White; on smelling the charcoal, a smell of onion is perceived.	<p>Presence of Sb.</p> <p>Bl.</p> <p>Cu.</p> <p>Pb.</p> <p>Sn.</p> <p>Ag.</p> <p>Zn.</p> <p>Cd.</p> <p>As.</p>
<i>Globules.</i>	<i>Incrustation.</i>																					
White and brittle.	White, . . . .																					
White and brittle.	Yellow, . . . .																					
Red and malleable.	None, . . . .																					
White and malleable marking paper easily.	Yellow, . . . .																					
White and malleable not marking paper, and readily fusible.	None, . . . .																					
White and malleable not marking paper, fusible only with difficulty.	None, . . . .																					
None, . . . .	<div> Yellow whilst hot, white when cold; becomes green when moistened with <math>\text{Co}(\text{NO}_3)_2</math> solution and reheated in the outer flame. </div>																					
None, . . . .	Brown, . . . .																					
None, . . . .	White; on smelling the charcoal, a smell of onion is perceived.																					

After the completion of the preliminary tests for the metal, a solution of the substance is made according to the directions in par. (332), and this solution is examined as there directed.

## B.—THE SUBSTANCE FOR ANALYSIS IS A LIQUID.

### 334. PRELIMINARY EXAMINATION.

*Note.*—The experiments or observations which are to be made on the liquid are stated in succession in the first column: opposite to these there are drawn out in the second column the results, any one of which may be obtained; and corresponding to these in the third column are given the inferences to be drawn from each of these results.

Observation or experiment to be made.	Result of observation or experiment.	Inference to be drawn from the result.
<p>I. Note the color of the liquid.</p> <p><i>Note.</i>—If the color has to be observed by night it should be examined by the white light obtained by burning a piece of magnesium ribbon. Delicate tints are best seen by looking through some thickness of the liquid at a sheet of perfectly white paper.</p>	<p><i>Pink</i> { Very delicate, . . . . .           { Intense, . . . . . <i>Green</i> { Very delicate, . . . . .           { Intense, . . . . . <i>Yellow</i> { Reddish-yellow, . . . . .           { Light yellow, . . . . . <i>Blue,</i> . . . . . <i>Violet</i> (or purple), . . . . . The liquid is colorless, . . . . .</p>	<p>Presence of Mn or dilute Co solution. Presence of Co. " Fe" or dilute Ni or Cr solution. " Ni, Cr, or Cu solution. " Fe" or a bichromate. " Neutralchromate. " Cu. " Cr or a Permanganate. Absence of Co, Ni, Cr, Cu, etc.</p>
<p>II. Dip pieces of red and blue litmus-paper into the liquid.</p>	<p><i>It is neutral,</i> (Not affecting the color of either paper.) . . . . .  <i>It is acid,</i> (Turning blue paper red.) . . . . .  <i>Note.</i>—If, on adding <math>\text{Na}_2\text{CO}_3</math> to a part of the solution and warming, no effervescence occurs, free acid and acid salt are absent.  <i>It is alkaline,</i> (Turning red paper blue.) . . . . .  <i>Note.</i>—If the color of the paper becomes only slightly changed, it indicates, probably, the presence of an alkaline salt.</p>	<p>Absence of free acids and alkalis, and of all salts but some of those of Am, Na, K, Mg, Ba, Sr, Ca, Ag.  Presence of a free acid, of an acid salt, or of a salt with acid reaction.  Presence of a hydrate of K, Na, Am, Ca, Sr, or Ba, or of a salt with alkaline reaction.</p>

PRELIMINARY EXAMINATION—*Continued.*

Observation or experiment to be made.	Result of observation or experiment.	Inference to be drawn from the result.
<p>III. Evaporate (21) some of the liquid upon a thin watch-glass,* heated very gently by supporting it some inches above a small flame upon a piece of wire gauze, or by means of a sand or water bath. If a residue remains on the glass bring this gradually into the flame and heat it more strongly.</p>	<p><i>A residue is left,</i> . . . . .</p> <p>On being strongly heated the residue blackens, and emits a smell of burning. . . . The smell resembles that of burning sugar.</p> <p><i>Note.</i>—If the cold residue after ignition effervesces with a drop of HCl, whereas the unignited residue did not, the organic acid radicle is united with Ba, Sr, Ca, Mg, Na, or K.</p> <p>[The results in Exp. I, 1, 2, 3, 5, par. 333, may be here observed, and should be noted.]</p> <p><i>No residue is left,</i> . . . . .</p> <p><i>Note.</i>—If the liquid is also colorless, tasteless, and without action on litmus-paper, it must consist only of distilled water, and no further examination of it need be made. If the liquid is alkaline <math>\text{NH}_3</math> must be looked for; if it is acid, <math>\text{H}_2\text{CO}_3</math>, <math>\text{H}_2\text{SO}_4</math>, <math>\text{HCl}</math>, <math>\text{HNO}_3</math>, <math>\text{H}_2\text{SO}_4</math>, and other volatile acids only need be tested for.</p>	<p>Presence of some dissolved solid.</p> <p>Pres. of <math>\bar{\text{A}}</math>, <math>\bar{\text{O}}</math>, or <math>\bar{\text{T}}</math>. Presence of <math>\bar{\text{T}}</math>.</p> <p>Absence of any dissolved solid.</p>
<p>IV. Add excess of strong NaHO or KHO solution and boil (51).</p>	<p>Smell of <math>\text{NH}_3</math>; moist red litmus is turned blue, and a rod moistened with strong HCl produces white fumes when held in the mouth of the test-tube.</p>	<p>Presence of Am. No further test need be tried for a metal if Am is detected here.</p>
<p>V. Dip a loop of platinum wire into the liquid, and hold it in the Bunsen flame (32); note the color of the flame, then moisten the loop with a drop of strong HCl, and hold again in the flame.</p>	<p>The flame is colored:</p> <p><i>Intense yellow,</i> <i>Pale violet,†</i> violet red when seen through the indigo-prism, . . . . .</p> <p><i>Red,</i> { Appearing <i>green</i> through the indigo-prism, . . . . .           { Appearing <i>intense red</i> through the indigo-prism, . . . . .</p> <p><i>Green,</i> { Yellowish-green, . . . . .           { Bright-green, . . . . .</p> <p><i>Blue,</i> { Pale (livid), . . . . .           { Vivid, . . . . .</p>	<p>Presence of Na.</p> <p>" K.</p> <p>" Ca.</p> <p>" Sr.</p> <p>" Ba.</p> <p>" <math>\text{B}_2\text{O}_3</math>, Cu (usually blue in centre).</p> <p>" As, Sb, Pb.</p> <p>" <math>\text{CuCl}_2</math>.</p>

\* A piece of thin glass from the side of a broken flask is a cheap and useful substitute for a watch-glass.

† This color is often rendered somewhat indefinite by the presence of a trace of Na. K is then found by the indigo-prism (49) or spectroscope.

PRELIMINARY EXAMINATION—*Continued.*

Observation or experiment to be made.	Result of observation or experiment.	Inference to be drawn from the result.														
VI. Make a clear borax bead in a platinum wire loop (31); dip it into a small quantity of the liquid contained in a watch-glass,* then fuse it in the outer and inner blowpipe flames; observe the color of the bead carefully both whilst hot and when cold.	<p>The bead is colorless both hot and cold, } after having been heated in both flames, }</p> <p>The bead is colored :</p> <table><tr><td>In outer flame.</td><td>In inner flame.</td></tr><tr><td><i>Blue</i>, hot; <i>green</i>, cold.</td><td><i>Red</i> or <i>colorless</i>, . .</td></tr><tr><td><i>Blue</i>, hot and cold.</td><td><i>Blue</i>, hot and cold, . .</td></tr><tr><td><i>Brown</i> or <i>dingy</i> } <i>purple</i>, hot; <i>tight</i> } <i>brown</i>, cold, }</td><td><i>Gray</i> or <i>black</i>, . . .</td></tr><tr><td><i>Brown</i>, hot; <i>yellow</i>, } cold, }</td><td>{ <i>Bottle-green</i>, hot } and cold, }</td></tr><tr><td><i>Green</i>,† hot and cold.</td><td><i>Green</i>, hot and cold, .</td></tr><tr><td><i>Purple</i>, hot and cold.</td><td><i>Colorless</i>, . . .</td></tr></table>	In outer flame.	In inner flame.	<i>Blue</i> , hot; <i>green</i> , cold.	<i>Red</i> or <i>colorless</i> , . .	<i>Blue</i> , hot and cold.	<i>Blue</i> , hot and cold, . .	<i>Brown</i> or <i>dingy</i> } <i>purple</i> , hot; <i>tight</i> } <i>brown</i> , cold, }	<i>Gray</i> or <i>black</i> , . . .	<i>Brown</i> , hot; <i>yellow</i> , } cold, }	{ <i>Bottle-green</i> , hot } and cold, }	<i>Green</i> ,† hot and cold.	<i>Green</i> , hot and cold, .	<i>Purple</i> , hot and cold.	<i>Colorless</i> , . . .	<p>Absence of the metals below.</p> <p>Presence of Cu. " Co. " Ni. " Fe.† " Cr. } See " Mn. } VI a.</p>
In outer flame.	In inner flame.															
<i>Blue</i> , hot; <i>green</i> , cold.	<i>Red</i> or <i>colorless</i> , . .															
<i>Blue</i> , hot and cold.	<i>Blue</i> , hot and cold, . .															
<i>Brown</i> or <i>dingy</i> } <i>purple</i> , hot; <i>tight</i> } <i>brown</i> , cold, }	<i>Gray</i> or <i>black</i> , . . .															
<i>Brown</i> , hot; <i>yellow</i> , } cold, }	{ <i>Bottle-green</i> , hot } and cold, }															
<i>Green</i> ,† hot and cold.	<i>Green</i> , hot and cold, .															
<i>Purple</i> , hot and cold.	<i>Colorless</i> , . . .															
VI a. <i>Confirmatory.</i> —The presence of Cr and Mn may be confirmed, if detected by the borax bead, by moistening a mixture of a little solid $\text{Na}_2\text{CO}_3$ and $\text{KNO}_3$ on platinum foil, with a few drops of the original solution, drying by gently heating it at a short distance above the flame, and then fusing by directing the blowpipe flame on the lower surface of the foil.	<p>A yellow mass on cooling (105), . . .</p> <p>A bluish-green mass on cooling (119), . .</p>	<p>Presence of Cr.</p> <p>" Mn.</p>														

## EXAMINATION FOR THE METAL.

**335.** If the original liquid leaves a residue on evaporation (334, III), and is neutral or acid to litmus-paper (334, II), examine as directed under I (336 *et seq.*); if it is alkaline work by II (359).

The method of analysis given below will be best understood by a reference to the Table of Analytical Classification of the metals (p. 186): the list of metals possibly present, which is entered here at the head of

\* If the solution is very dilute, a portion of it should be evaporated nearly to dryness in a small porcelain dish before the bead is dipped into it; if after dipping the bead once and fusing, no color is seen, it should be wetted once more with the solution and dried in the flame, and this process should be several times repeated before again fusing the bead.

† Cr, if present as a chromate, gives a bead *brown* while hot after having been fused in the outer blowpipe flame, but becoming permanently green in both flames after having been heated in the inner flame.

each Group Table, will be seen to correspond with the groups in that table.

### I. THE LIQUID IS NEUTRAL OR ACID TO LITMUS.

**336. Examination for Group I (Pb, Ag, Hg').**—To a portion of the liquid add HCl in excess (35 a), and note any change which occurs (see Note 1). If a white precipitate forms,\* add more HCl and heat for a moment to boiling, then cool at once; if any precipitate remains, work as directed under the left-hand column; if no precipitate remains, refer to the right-hand column (note 2).

<p><i>A white precipitate forms</i> (see note 3), and is either not dissolved when boiled with HCl, or dissolves and reappears on cooling the liquid (showing the <i>Presence of Pb</i>). The precipitate may consist of AgCl, PbCl<sub>2</sub>, Hg<sub>2</sub>Cl<sub>2</sub>.</p> <p>Let the precipitate settle and decant the liquid (see note 4); boil the precipitate in the test-tube with AmHO (see note 5):</p>		<p><i>No precipitate forms</i>, or a precipitate forms in the cold, but disappears on boiling after addition of more HCl, and does not reappear on cooling the liquid:</p> <p><i>Absence of Ag, and probably of Pb and Hg'.</i></p> <p><i>Note.</i>—Hg' may possibly be passed over in this group, since Hg<sub>2</sub>Cl<sub>2</sub> is dissolved as HgCl<sub>2</sub> very readily when warmed with HCl and HNO<sub>3</sub>, and slowly when boiled with HCl alone. Hence it is most safe to examine some of the <i>original</i> precipitate (before it has been boiled with HCl) by the first column.</p> <p>Pass on to (337).</p>	
<p>1. <i>The precipitate dissolves</i>, at least in part (see note below):</p> <p><i>Presence of Ag.</i> Confirm by adding excess of HNO<sub>3</sub> to the AmHO solution, a milkiness appears:</p> <p><i>Presence of Ag.</i></p> <p><i>Note.</i>—Coagulated AgCl often dissolves only slowly in AmHO.</p>	<p>2. The precipitate does not dissolve:</p> <p><i>The color of the precipitate is unchanged:</i></p> <p><i>Presence of Pb.</i> Confirm by decanting the liquid, dissolving the precipitate in H<sub>2</sub>A, and adding K<sub>2</sub>CrO<sub>4</sub> to the solution; a yellow precipitate:</p> <p><i>Presence of Pb.</i></p> <p><i>Note.</i>—The presence of Pb is also confirmed by the precipitate with HCl dissolving on boiling and reappearing on cooling.</p>	<p><i>The precipitate becomes black:</i></p> <p><i>Presence of Hg' (Mercurosum).</i> Confirm by immersing a bright slip of copper in a portion of the original solution, made acid with a few drops of HNO<sub>3</sub>; a gray deposit forms on the Cu, which becomes white and bright when rubbed:</p> <p><i>Presence of Hg' (Mercurosum).</i></p>	

*Note 1.*—Effervescence may be noticed on the addition of HCl, owing to certain gases being evolved, a list of which will be found in (349); some of them are readily detected by their smell or color, but special tests need not be tried, as they will be afterwards used in the Preliminary Examination for acid radicles (348 *et seq.*); the fact that a gas is evolved should however always be noted down, even if the nature of the gas is not ascertained.

A yellow liquid whose color is due to the presence of a neutral chromate turns reddish-yellow on addition of HCl; this confirms the presence of a chromate.

*Note 2.*—A white precipitate which disappears on boiling after

\* Yellow S may separate from a theiosulphate (see 349; 2); this precipitate requires no further examination.

## 198 EXAMINATION OF A SIMPLE SUBSTANCE. [337.

addition of more HCl, and does not reappear on cooling, may consist of BiOCl or SbOCl. A little of the milky liquid produced by the first addition of HCl should be mixed with H<sub>2</sub>T solution, if the precipitate consists of SbOCl it will disappear, BiOCl is not thus dissolved.

*Note 3.*—A colorless, gelatinous precipitate of hydrated SiO<sub>2</sub> may form here; confirm the presence of SiO<sub>2</sub> by evaporating a portion of the original solution to dryness with HCl, and warming the residue with HCl; a residue insoluble in HCl proves the presence of SiO<sub>2</sub>; filter or decant from the residue of SiO<sub>2</sub>, and employ the HCl solution to detect the metal present by (337 *et seq.*).

*Note 4.*—In this and all subsequent places where decantation of the liquid is recommended, it should be understood that a precipitate will usually settle if the liquid is well shaken or heated and allowed to stand for several minutes. (See 23 *b.*)

*Note 5.*—In case the liquid cannot be completely decanted, add the AmHO in excess to the liquid in which the precipitate is suspended.

**337. Examination for Group II (Hg, Pb, Bi, Cu, Cd, Sn, As, Sb).**—To the above portion of the solution, made acid with HCl and boiled until free from Cl or SO<sub>2</sub>, if these gases are evolved on heating, add water; a milkiness due to precipitation of BiOCl or SbOCl may result, SbOCl being distinguished from BiOCl by a few drops of the liquid becoming clear on addition of H<sub>2</sub>T solution. Whether a milkiness has been produced or not by water, add strong H<sub>2</sub>S-water or pass H<sub>2</sub>S until the liquid after having been well shaken smells strongly of the gas, and heat gently (see Note 1 below); if no precipitate forms, heat to boiling, since Sn (as stannicum) and As (as arsenicum) are only precipitated by H<sub>2</sub>S on boiling; if a precipitate has formed examine it by the following table; if none has appeared, or only a white milkiness due to precipitation of sulphur (see Note 2 below), the absence of Hg, Pb, Bi, Cu, Cd, Sn, As, and Sb is proved; pass on to (338).

*Note 1.*—Certain changes in the color of the precipitate may be noticed if the H<sub>2</sub>S is slowly added; thus Hg'' gives a precipitate which is at first white, then yellow, brown, and ultimately black; Pb, in strongly-acid solutions, gives a red precipitate, becoming black on addition of water and more H<sub>2</sub>S.

*Note 2.*—Characteristic changes of color of the solution often accompany the separation of sulphur; thus

A *reddish-yellow* solution changing to *pale green* indicates a ferric salt.

A *reddish-yellow* solution changing to *green* indicates a chromate.

A *green* solution changing to *colorless* indicates a manganate.

A *purple* solution changing to *colorless* indicates a permanganate.

The precipitated S need not be examined.

Note the color of the precipitate yielded by  $H_2S$ , and examine it according to 1, 2, or 3, below:

<p>1. <i>The precipitate is dark-brown or black</i>; it may consist of <math>SnS</math>, <math>CuS</math>, <math>HgS</math>, <math>Bi_2S_3</math>, <math>PbS</math>. Add pure <math>NaHO</math> in excess to the liquid and precipitate, or to the precipitate alone after decanting the liquid if possible (see note 4, p. 198), and boil:</p>		
<p><i>The precipitate dissolves</i>: to a portion of the <math>NaHO</math> solution add <math>HCl</math> in excess, brown <math>SnS</math> is reprecipitated. Boil another portion after adding a little yellow <math>Am_2S_3</math>, then add <math>HCl</math> in excess, a yellow precipitate of <math>SuS_2</math>: <i>Presence of Sn''</i> (Stannosum). [The presence of <math>Sn''</math> may also be confirmed by adding <math>HgCl_2</math> to a part of the original solution, a white precipitate shows <i>Presence of Sn''</i>.]</p>	<p><i>The precipitate does not dissolve</i>: it may consist of <math>CuS</math>, <math>HgS</math>, <math>Bi_2S_3</math>, <math>PbS</math>. <i>a. The original solution is blue or green</i>, and becomes intensely blue on addition of excess of <math>AmHO</math>: <i>Presence of Cu</i>. Confirm by adding a few drops of <math>H_2SO_4</math> to the original solution, and dipping into it a bright knife-blade or piece of steel, a red film of <math>Cu</math> deposits: <i>Presence of Cu</i>. <i>Note</i>.—To detect small quantities of <math>Cu</math> the <math>K_4Cfy</math> test (in 163) should be used. <i>β. The original solution is colorless</i>. Test in separate portions for <math>Hg</math>, <math>Bi</math>, <math>Pb</math>, by the following special tests:</p>	
<p>After acidifying with <math>HCl</math> (if not already acid), immerse in the solution for a few minutes a bright strip of <math>Cu</math>; a gray film is deposited, which becomes white and brilliant when rubbed: <i>Presence of Hg''</i> (Mercurium). [The <math>SnCl_2</math> test (145) may also be used as confirmatory.]</p>	<p>Pour another portion of the original solution into much cold distilled water; a white precipitate or milkiness: <i>Presence of Bi</i>. <i>Note</i>.—If <math>Bi</math> is suspected and no milkiness appears, add excess of <math>AmHO</math> to the liquid; <math>Bi</math> (if present) falls as hydrate; filter, pour one or two drops of boiling <math>HCl</math> upon the precipitate and let them drop through into large excess of water; a milkiness, on stirring, shows <i>Presence of Bi</i>.</p>	<p>To a portion add <math>H_2SO_4</math>, a white precipitate: <i>Presence of Pb</i>. [The presence of <math>Pb</math> may be confirmed by adding to a part of the original solution <math>AmHO</math> in excess, then <math>HA</math> in excess, then <math>K_2CrO_4</math>. A yellow precipitate shows <i>Presence of Pb</i>.]</p>
<p>2. <i>The precipitate is yellow</i>: it may consist of <math>CdS</math>, <math>As_2S_3</math>, <math>SnS_2</math>. Boil the precipitate with <math>NaHO</math> in excess, after decanting the liquid if possible:</p>		
<p><i>The precipitate does not dissolve</i>: it is bright yellow and pulverulent: <i>Presence of Cd</i>. [The presence of <math>Cd</math> may be confirmed by showing that the precipitate produced by <math>H_2S</math> dissolves on being boiled with dilute <math>H_2SO_4</math>.]</p>	<p><i>The precipitate dissolves</i>: it may consist of <math>As_2S_3</math> or <math>SnS_2</math>; examine portions of the original solution as directed below:</p> <p>Acidify a portion with <math>HCl</math> and place in it a piece of <math>Zn</math> resting on a piece of platinum foil: <math>Su</math> is deposited on the <math>Zn</math>: <i>Presence of Sniv</i>. (Stannicum). [The deposit of <math>Su</math> may be detached from the <math>Zn</math>, and dissolved by boiling it in a test-tube with a little strong <math>HCl</math>, then <math>HgCl_2</math> added; a white precipitate confirms <i>Presence of Sniv</i>.]</p>	<p>Acidify a portion with strong <math>HCl</math>, immerse in it some pieces of bright <math>Cu</math> and boil, a black film is deposited on the <math>Cu</math>: <i>Presence of As</i>. <i>Note</i>.—If the yellow precipitate formed by <math>H_2S</math> appeared at once in the cold, the <math>As</math> is present as <i>Arsenorum</i>; if it formed only on boiling, <i>Arsenicum</i> is present.</p>
<p>3. <i>The precipitate is orange-red</i>: it may consist of <math>Sb_2S_3</math>. Confirm by acidifying a portion of the original solution with <math>HCl</math>, and immersing in it a piece of <math>Zn</math> resting on a slip of platinum foil, a black stain on the platinum (193): <i>Presence of Sb</i>. <i>Note</i>.—It occasionally happens, if the solution has not been sufficiently diluted, that <math>H_2S</math> gives a reddish precipitate with <math>Pb</math> solution; such a precipitate, however, becomes black on adding more <math>H_2S</math>-water.</p>		



**338.**—*Examination for Group III.* (Al, Fe, Cr; Zn, Mn, Ni, Co; Ba, Sr, Ca, Mg as phosphate, fluoride, or oxalate;  $\text{BaCrO}_4$ ).

*Caution.*—It is of the greatest importance to remember that if an organic salt—more especially a tartrate—has been detected in the preliminary examination, and it has not been proved to be a salt of Ba, Sr, Ca, Mg, K, or Na, and must therefore be a salt of one of the metals of Group III, the detection of the metal by the directions given below may be hindered or prevented. It is therefore necessary to destroy the organic acid radicle by evaporating some of the substance to dryness after adding a little strong  $\text{HNO}_3$ , and then gently igniting the residue. The residue, when cool, is dissolved by warming with a few drops of  $\text{HCl}$ , the solution diluted, and *this solution is to be substituted for the original solution in the examination for Group III.*

To another portion of the original solution add first  $\text{AmCl}$  solution, then  $\text{AmHO}$  in excess; note carefully the color and appearance of any precipitate (which is probably due to Al, Fe or Cr, but if white or nearly so may be a phosphate, oxalate, etc.), or the color of the liquid if no precipitate forms (Ni yields a violet-colored liquid); then whether a precipitate has been produced or not, add  $\text{Am}_2\text{S}$  or better  $\text{H}_2\text{S}$ , and boil. (See note below.)

*Note.*—If no precipitate forms on addition of  $\text{AmCl}$  and  $\text{AmHO}$ , or of  $\text{Am}_2\text{S}$ , until the liquid is boiled, when a pale-green precipitate is produced, this precipitate is due to the presence of a chromate, which will have been already detected by the liquid turning from yellow to green on addition of  $\text{H}_2\text{S}$ . The pale-green precipitate need not be further examined, but its formation should be noted as confirmatory of the presence of a chromate. Proceed to examine for the metal present by (342).

If a precipitate forms examine it by the table below.\*

---

\* White S, separated from the  $\text{Am}_2\text{S}$  by the action of a ferricyanide or other oxidizer, may here render the liquid milky: it is recognized by disappearing again when a small quantity of the milky liquid is warmed with more  $\text{Am}_2\text{S}$ . Its appearance should be noted, but it is not to be considered as a third group precipitate.

If no precipitate forms, the absence of Al, Fe, Cr, Zn, Mn, Ni, Co, and of Ba, Sr, Ca, Mg as phosphate, oxalate, and probably as fluoride, also of  $\text{BaCrO}_4$ , is proved; pass on to (342).

Note the color of the precipitate, and examine it accordingly by 1 or 2 below.

1. *The precipitate after addition of  $\text{Am}_2\text{S}$  is black, it may consist of  $\text{FeS}$ ,  $\text{CoS}$ , or  $\text{NiS}$ ; the means of distinguishing these three metals is given below. Note first the color of the original solution (334, 1).*

<p><i>a. The color of the original solution was yellow or reddish brown:—Presence of <math>\text{Fe}^{+++}</math>.</i></p>	<p><i>b. The color of the original solution was pink, or possibly blue, becoming pink on dilution:—Presence of Co.</i></p>	<p><i>c. The color of the original solution was green:—Presence of <math>\text{Ni}</math> or <math>\text{Fe}^{++}</math>.</i></p>
<p>Note the effect which was produced by the addition of <math>\text{AmCl}</math> and <math>\text{AmHO}</math>:</p>		
<p>This is confirmed by the precipitate produced by <math>\text{AmCl}</math> and <math>\text{AmHO}</math> in the original solution having been reddish-brown, and becoming black on addition of <math>\text{Am}_2\text{S}</math>.</p>	<p>The presence of Co may be confirmed by adding excess of <math>\text{KHO}</math> to the original solution, a blue precipitate turning red on boiling shows the presence of Co.</p>	<p><i>A blue solution was produced:—Presence of <math>\text{Ni}</math>.</i></p>
<p>As a confirmatory test add to a portion of the original solution (acidified with <math>\text{HCl}</math> if necessary), a few drops of <math>\text{KC}_2\text{S}</math> solution, a blood-red coloration, destroyed by <math>\text{HgCl}_2</math>, shows:</p>	<p>Co will also have been detected by its borax bead (334, VI).</p>	<p>The presence of Ni may be confirmed by adding excess of <math>\text{KHO}</math> to a part of the original solution; a light-green precipitate shows the presence of Ni.</p>
<p><i>Presence of <math>\text{Fe}^{++}</math> (Ferrous).</i></p>		<p>Ni will also have been detected by its borax bead (334, VI).</p>
		<p><i>A dingy-green precipitate, which, when shaken round in a porcelain dish and left for several minutes exposed to the air, becomes brown:—Presence of <math>\text{Fe}^{++}</math>.</i></p>
		<p>Confirm by adding <math>\text{K}_2\text{C}_2\text{O}_4</math> to some of the original solution (acidified with <math>\text{HCl}</math>, if necessary); a dark-blue precipitate:—Presence of <math>\text{Fe}^{++}</math> (Ferrous).</p>

2. *The precipitate, after addition of  $\text{Am}_2\text{S}$ , is white or light-colored.—It may consist of  $\text{Cr}_2\text{H}_2\text{O}_6$ ,  $\text{Al}_2\text{H}_2\text{O}_6$ ,  $\text{ZnS}$ ,  $\text{MnS}$ ; [or of  $\text{BaCrO}_4$ , or Ba, Sr, Ca, Mg, as phosphate, oxalate, fluoride, silicate, or possibly borate]. The substances*

in brackets can however only be present if the liquid given for analysis was acid in reaction, or if a solid was given which was insoluble in water.

To a small portion of the original solution add KHO drop by drop until a precipitate is produced. (See note below the table.) Note the color and appearance of this precipitate, then add a larger quantity of KHO and stir or shake well; one of the following results (a or b) will be noticed:

a. *The precipitate at first produced by KHO redissolves when the KHO is added in larger quantity; indicating the presence of Al, Zn, or Cr. Note the appearance of the precipitate:*

<p><i>A pale green flocculent precipitate, which is reprecipitated from the solution in KHO. when that liquid is diluted and boiled for several minutes, or when it is mixed with AmCl:—</i>  <i>Presence of Cr.</i></p> <p>Confirm by boiling some of the green KHO solution with lead peroxide (<math>PbO_2</math>); a yellow liquid is obtained, which, if decanted, yields a yellow precipitate on addition of <math>H\bar{A}</math> in excess.</p> <p>The color of the solution and of the borax bead will also have detected Cr.</p> <p><i>Note.</i>—If the Cr was present as a chromate, AmCl and AmHO will have yielded no precipitate; but there will have been a green precipitate on addition of <math>Am_2S</math> and heating.</p>	<p><i>A white precipitate, which is reprecipitated from the KHO solution by addition of a few drops of <math>H_2S</math>-water, but is not reprecipitated by addition of AmCl:—</i>  <i>Presence of Zn.</i></p> <p>Confirmed by no precipitate having been produced by AmCl and AmHO, but a white precipitate by addition of <math>Am_2S</math>, or of <math>H_2S</math> after AmHO.</p>	<p><i>A white flocculent precipitate, not reprecipitated from the KHO solution by addition of a few drops of <math>H_2S</math>-water or by boiling, but reprecipitated immediately if sufficient AmCl solution is added:—</i>  <i>Presence of Al.</i></p> <p><i>Note.</i>—<math>AlPO_4</math> is precipitated here also, and is soluble in KHO, but differs from <math>Al_2HO_6</math> in being reprecipitated from the KHO solution by addition of excess of <math>H\bar{A}</math>; hence, if on acidifying the KHO solution with <math>H\bar{A}</math> the precipitate reappears, test for <math>PO_4</math> by (339, 1); if <math>PO_4</math> is detected the analysis is completed. Further, <math>AlPO_4</math> is soluble only in acids; hence the original liquid must have been acid or the original solid substance insoluble in water, and in the acid solution AmCl and excess of AmHO will have given a colorless gelatinous precipitate resembling <math>Al_2HO_6</math>.</p>
---	--	--

*Note.*—Sometimes a flocculent precipitate is produced on adding AmCl and AmHO to the original solution, and yet no precipitate is produced by KHO; this renders probable the presence of an alkaline silicate. In this case evaporate a portion of the solution to dryness with HCl; warm the residue with HCl: if any residue is left, the *presence of a silicate* is proved; the clear solution is further examined for K or Na by (345).

- b. *The precipitate at first produced by KHO does not re-dissolve on addition of more KHO; the precipitate will have one of the following appearances:*

<p><i>A white or dingy yellow precipitate, rapidly darkening when shaken in the air:</i>  <i>Presence of Mn.</i>  <i>Confirmed</i> in the preliminary examination by producing a green mass when fused with <math>\text{Na}_2\text{CO}_3</math> and <math>\text{KNO}_3</math> and by no precipitate having been produced on addition of <math>\text{AmCl}</math> and <math>\text{AmHO}</math>, unless the solution was exposed for some time to the air.</p>	<p><i>A pale yellow precipitate not darkening in the air:</i> this precipitate will also have been produced by <math>\text{AmCl}</math> and <math>\text{AmHO}</math>, and its color will not have been altered by addition of <math>\text{AmS}</math>:  <i>Presence of BaCrO<sub>4</sub>.</i>  <i>Ba</i> has been already confirmed by the flame coloration (334, V), and a chromate by the change of color of the orange-red liquid to green on adding <math>\text{H}_2\text{S}</math> and heating (337, note 2).</p>	<p><i>A white precipitate, not darkening in the air:</i>  <i>Presence of Ba, Sr, Ca, or Mg, as phosphate, oxalate, etc.</i>  <i>Pass on to (339).</i>  <i>Note.</i>—This precipitate can only be formed if the original solution was acid.</p>
--	--	--

**339.** *The precipitate may consist of Ba, Sr, Ca, or Mg as phosphate, oxalate, borate,\* fluoride, or silicate.*

Examine for the metal and acid radicle as directed below, working through the columns from left to right, unless the presence of an oxalate (333, I, 5) or borate (334, V, or 333, II) has been already proved in the preliminary examination, in which case proceed at once to examine according either to column 2 or 3. If this precipitate is proved to contain a phosphate, oxalate, borate, fluoride, or silicate, the further examination for an acid radicle becomes unnecessary.

---

\* Borates of Ba, Sr, and Ca are rarely precipitated in Group III, since they are soluble in the  $\text{AmCl}$  formed on adding  $\text{AmHO}$  to the  $\text{HCl}$  solution.

<p>1. Add a few drops of the original solution to some <math>\text{AmHMoO}_4</math> solution, and warm gently: a yellow precipitate (277) shows <i>Presence of a phosphate.</i> Examine further for the metal present by (340).</p> <p>If no precipitate is produced by <math>\text{AmHMoO}_4</math>, pass on to column 2.</p>	<p>2. Add <math>\text{AmHO}</math> in excess to a portion of the original solution,* filter, dry the precipitate and heat a portion of it to a dull red heat for several minutes on platinum foil. Place the foil in a test-tube, and pour a few drops of dilute <math>\text{HCl}</math> upon it; if the residue dissolves with effervescence, it shows the <i>Presence of an oxalate.</i></p> <p>Test this <math>\text{HCl}</math> solution for Ba, Sr, Ca, Mg, by (342-344), using it as the original solution.</p> <p>The presence of an oxalate may be confirmed by heating the precipitate* with strong <math>\text{H}_2\text{SO}_4</math>, when <math>\text{CO}</math> and <math>\text{CO}_2</math> will be evolved; and if the substance under analysis is a solid, it will have been found also by (333, I, 5).</p> <p>If the ignited precipitate does not dissolve with effervescence, pass on to column 3.</p>	<p>3. Stir some of the precipitate produced by <math>\text{AmHO}</math>* from column 2, on a watch-glass, with a little water to which a few drops of <math>\text{HCl}</math> have been added, until the precipitate is dissolved. Dip into the liquid a slip of turmeric paper, and dry at steam heat; the slip appears <i>reddish-brown</i>, and becomes <i>dingy green</i> when moistened with <math>\text{AmHO}</math>:— <i>Presence of a borate.</i></p> <p>The presence of a borate may also be detected or confirmed by heating with alcohol and strong <math>\text{H}_2\text{SO}_4</math>, and kindling the alcohol: a green-edged flame shows a borate.</p> <p>Examine for the metal present by (341).</p> <p>If a borate is not found proceed to column 4.</p>	<p>4. Place another portion of the precipitate produced by <math>\text{AmHO}</math>* (column 2) in a leaden cup or platinum crucible, add strong <math>\text{H}_2\text{SO}_4</math>, and warm gently, covering the vessel with a watch-glass, which has been coated with a film of wax and has had characters traced with a sharp point through the wax (296); the glass is etched:— <i>Presence of a fluoride.</i></p> <p>Examine for the metal present by (341).</p> <p>If no fluoride is found proceed to column 5.</p>	<p>5. If phosphate, oxalate, borate and fluoride have been proved to be absent, a silicate is probably present. Test for it by par. 341.</p>
--	--	--	--	--

**340. Explanation of Method.**—In order to detect the Ba, Sr, Ca, or Mg, present as phosphates, the  $\text{PO}_4$  must be removed from the solution and the metal left as chloride. This is effected by neutralizing all  $\text{HCl}$  and leaving the solution acidified only by free  $\text{H}\bar{\text{A}}$ ; by *gradual* addition of  $\text{Fe}_2\text{Cl}_6$  and boiling, all  $\text{PO}_4$  is precipitated as  $\text{FePO}_4$ , which (though soluble in  $\text{HCl}$ ) is perfectly insoluble in  $\text{H}\bar{\text{A}}$ . On filtering,  $\text{FePO}_4$  remains upon the filter, and

\* If a solid is being examined, a portion of the original solid is employed for this test, instead of the precipitate produced by  $\text{AmHO}$  in the  $\text{HCl}$  solution.

the filtrate contains the metal (Ba, Sr, Ca, or Mg) as chloride, and is perfectly free from phosphate.

*Method.*—Add to a portion of the cold original solution AmHO, drop by drop, whilst constantly shaking the liquid, until a slight precipitate is formed, which does not disappear by shaking; then add H $\bar{A}$  in slight excess (35 a), the precipitate will redissolve: dissolve in the cold liquid a little solid Na $\bar{A}$  by shaking, and add Fe<sub>2</sub>Cl<sub>6</sub> drop by drop, until the liquid after being well shaken has a brown or red color; boil and filter immediately. The precipitate is thrown away. To the filtrate AmCl is added, then AmHO until it is just in excess (35 a); if any precipitate forms, boil and filter; the liquid or filtrate, which must smell strongly of NH<sub>3</sub>, is then tested for Ba, Sr, Ca, and Mg, by (342–344), this liquid taking the place of the original solution referred to in (342). See note.

*Note.*—If any precipitate is formed in (342) by Am<sub>2</sub>CO<sub>3</sub>, it must be well washed several times with boiling water to remove Na, since else the intense yellow coloration of Na would mask that of Ba, Sr, or Ca.

**341.** A portion of the original substance is placed in an evaporating basin, which is then nearly filled with dilute HCl, and the liquid is entirely boiled away; the dish is once more filled with acid, and again evaporated to dryness: the residue in the dish is then warmed with dilute HCl, and filtered if any residue is left. (See note, p. 206.)

*Solution:* the HCl solution is examined for Ba, Sr, Ca, Mg by (342–344); this solution being treated as is there directed for the original solution.

*Residue:* if any residue remains insoluble in HCl, it proves:

*Presence of a silicate.*  
(See note.)

*Note*—The evaporation to dryness with HCl removes  $H_2BO_3^*$  and HF, which volatilize, whilst  $H_2SiO_3$  is left insoluble in HCl, and therefore remains as an insoluble powder on afterwards warming with HCl; this residue of  $SiO_2$  also remains, however, if HF was present, being caused by the corrosion of the dish; since, however, HF if present has already been found; no notice will then be taken of the residue.

### 342. Examination for Group IV.

To a portion of the original solution add  $AmCl$ ,  $AmHO$  in excess, and then  $Am_2CO_3$  solution; warm gently:

<p>A white precipitate forms, which may consist of <math>BaCO_3</math>, <math>SrCO_3</math>, <math>CaCO_3</math>. Heat and filter, or let stand for a time and decant; dissolve the precipitate by pouring upon it a little hot dilute HCl, and proceed to examine the solution by (343).</p>	<p>No precipitate forms:— Absence of Ba, Sr, Ca. Pass on to (344).</p>
---	--

343. Dip a loop of platinum wire into the HCl solution, and hold the wire in the Bunsen flame. If the flame coloration thus obtained is not decisive, employ the tests placed at the foot of each column as confirmatory:

<p>A yellowish-green flame:†</p> <p>Presence of Ba.</p> <p>Confirmatory.—Add <math>CaSO_4</math> to a perfectly cold portion of the HCl solution, an immediate precipitate shows Presence of Ba.</p>	<p>A crimson flame, appearing deep red through the indigo-prism:</p> <p>Presence of Sr.</p> <p>Confirmatory.—Add <math>CaSO_4</math> to a perfectly cold portion of the HCl solution, no precipitate; but an immediate precipitate on boiling:</p> <p>Presence of Sr.</p>	<p>A red flame, appearing dingy green through the indigo-prism:</p> <p>Presence of Ca.</p> <p>Confirmatory.—Add <math>CaSO_4</math> to a portion of the HCl solution and boil; no precipitate.</p> <p>To another portion add <math>AmHO</math> in excess, and <math>Am_2C_2O_3</math>: white precipitate:</p> <p>Presence of Ca.</p>
--	---	--

\* Several evaporations are necessary in order to remove  $H_2BO_3$  entirely if it is present in large quantity; but its presence interferes only with the flame colorations of Ba, Sr, and Ca, and not with the liquid tests, and is therefore not of serious consequence.

† A brilliant green coloration may be seen following a yellowish-green or red coloration; it is due to the presence of  $H_2BO_3$ .

**344. Examination for Group V.**

To the liquid, in which  $\text{Am}_2\text{CO}_3$  has produced no precipitate, add  $\text{Na}_2\text{HPO}_4$ , and if a precipitate does not form at once warm the liquid gently, shake or stir it well, and let it stand for some time; a white crystalline precipitate shows *presence of Mg*. If no precipitate forms, even after standing, Mg is absent; proceed to (345).

**345.** Into a portion of the original solution, acidified (if not already acid) with HCl, dip a loop of platinum wire, and hold it in the Bunsen flame; if no distinct coloration is obtained, boil down some of the original solution in a porcelain dish nearly to dryness, add a few drops of HCl, and again try the flame coloration:

*A pale violet flame* (sometimes tinged yellow by a trace of Na), which appears violet or red through the indigo-prism:—*Presence of K*. Confirm by stirring some of the original solution, boiled down, if very dilute, with  $\text{PtCl}_4$  and a few drops of HCl: a *yellow* precipitate shows *presence of K*. Addition of alcohol much aids the formation of the precipitate.  
[NaHT may be substituted for  $\text{PtCl}_4$ , as being more economical.]

*Intense yellow flame*: which shows no red or violet color through the indigo-prism:—*Presence of Na*.

*Note.*—If the preceding examination has not detected a metal, no commonly occurring metal can be present in any quantity.

### EXAMINATION FOR THE ACID RADICLE IN A SOLID SUBSTANCE OR SOLUTION.

**346.** Several acid radicles will usually have been already detected during the examination for the metal. Thus, on acidifying with HCl and warming (336) the acid radicles mentioned in (349) would be detected.

**346 a.** On adding  $\text{H}_2\text{S}$ -water and warming (337), the



following acid radicles will have been detected by the following changes :

- |                                |  |  |
|--------------------------------|--|--|
| a. Change of color.            | $\left\{ \begin{array}{l} \text{CrO}_4: \text{ a yellow or orange-red solution, becoming green,} \\ \text{"MnO}_4: \text{ a green solution, becoming colorless,} \\ \text{'MnO}_4: \text{ a purple solution, becoming colorless,} \end{array} \right.$   | $\left. \vphantom{\left\{ \begin{array}{l} \text{CrO}_4: \text{ a yellow or orange-red solution, becoming green,} \\ \text{"MnO}_4: \text{ a green solution, becoming colorless,} \\ \text{'MnO}_4: \text{ a purple solution, becoming colorless,} \end{array} \right\}} \right\} \text{ and depositing white S, which makes the liquid milky.}$ |
| b. Formation of a precipitate. | $\left\{ \begin{array}{l} \text{"SnO}_3: \text{ a dingy-yellow precipitate, usually forming only when heat is applied.} \\ \text{"AsO}_3: \text{ a bright - yellow precipitate, forming at once in the cold.} \\ \text{"AsO}_4: \text{ a bright - yellow precipitate, forming at once if the liquid is boiled; but only after a long time in the cold solution.} \\ \text{"SbO}_4: \text{ an orange-red precipitate.} \end{array} \right.$ |  |

Also  $\text{PO}_4$ ,  $\text{BO}_3$ ,  $\text{F}$ ,  $\bar{\text{O}}$ ,  $\text{SiO}_3$ , may have been detected, if in combination with Ba, Sr, Ca, or Mg (339-341).

If any one of the above acid radicles has been found and confirmed by a special test, together with the metal with which it is combined, any further examination for an acid radicle is unnecessary. If no acid radicle has yet been found, proceed according to (347 *et seq.*).

347. Since the metal present has been detected already, the examination for the acid radicle may be usually much simplified by considering the solubility in water or in acids of the different saline compounds of this metal. Thus a salt of the metal which is insoluble in water cannot possibly be present in a neutral solution; a salt insoluble in water and acids cannot be present in a liquid with either acid or neutral reaction. A reference to the Table of Solubilities (455-458, p. 271) will thus frequently limit very considerably the number of acid

radicles that can possibly be present; and in cases where this number is very small, the analyst may dispense with the full analytical course drawn out below, and proceed at once to detect the acid radicle by trying special tests for those acid radicles only which can possibly be present.

### PRELIMINARY EXAMINATION FOR THE ACID RADICLE.

**348.** Pour dilute  $\text{H}_2\text{S}_2\text{O}_4$  in excess upon the substance in a test-tube, and observe the effect produced;\* then heat to boiling:

**349.** A gas is evolved, shown by the occurrence of effervescence or by a characteristic smell; see the six cases below:

<p>1. <i>Colorless gas without smell.</i> A drop of lime-water, held in the tube upon the end of a glass rod becomes milky:— <i>Presence of a carbonate.</i></p>	<p>2. <i>Gas smelling of burning sulphur.</i> A drop of <math>\text{K}_2\text{Cr}_2\text{O}_7</math> solution, held in the tube upon the end of a glass rod, becomes green:— <i>Presence of a sulphide.</i> The <math>\text{SO}_2</math> gas is given off only after standing for a time or on warming, and yellow S separates:— <i>Presence of a theiosulphate.</i></p>	<p>3. <i>Gas with a most fetid smell.</i> A slip of filter-paper moistened with alkaline solution of <math>\text{PbA}_2</math>, and held in the tube, is blackened:— <i>Presence of a sulphide.</i></p>
<p>4. <i>Yellow gas, smelling of Cl.</i> A slip of moist litmus-paper held in the tube is bleached:— <i>Presence of a hypochlorite.</i></p>	<p>5. <i>Reddish-brown fumes are evolved</i>, best seen by looking down the tube at white paper. On adding a little cold dilute <math>\text{H}_2\text{O}_2</math> and <math>\text{FeSO}_4</math> solution to a fresh portion, a deep-brown liquid:— <i>Presence of a nitrite.</i></p>	<p>6. <i>Smell of <math>\text{HCy}</math>.</i> Test for a cyanide and a ferro-, ferri- and sulphocyanide in the original substance by (354, 355). <i>No gas is evolved:—</i> Absence of carbonate, sulphite, theiosulphate, sulphide, hypochlorite, nitrite. <i>Pass on to (350).</i></p>

**350.** Add strong  $\text{H}_2\text{SO}_4$  in small quantity to a fresh portion of the substance, mix well by shaking, heat the liquid and note whether any of the results enumerated below occurs;\* if none of them is noticed, examine the tube by (351).

\* If Pb, Ba, Sr, or Ca is present, the addition of  $\text{H}_2\text{SO}_4$  causes a precipitate of the insoluble sulphate: this in no way interferes with the results to be observed and may be disregarded.

<p>1. <i>The liquid becomes yellow or red (see column 5), and a chlorous smell is given off: on warming the tube, a crackling noise or slight explosion is produced:—</i>  <i>Presence of a chlorate.</i></p> <p><i>Confirmatory.</i>—To a portion of the cold original solution* add a few drops of dilute indigo solution, then some <math>H_2SO_4</math>, the blue color of the indigo is destroyed:—  <i>Presence of a chlorate.</i></p>	<p>2. <i>A smell of vinegar is given off:—</i>  <i>Presence of an acetate.</i></p> <p><i>Confirmatory.</i>—Add to a fresh portion of the substance some alcohol, then strong <math>H_2SO_4</math>, and heat; a fragrant smell is given off:—  <i>Presence of an acetate.</i></p>	<p>3. <i>Acid fumes are given off, but the liquid is not colored: drop in some small pieces of Cu and heat to boiling, reddish-brown fumes are evolved:—</i>  <i>Presence of a nitrate.</i></p> <p><i>Confirmatory.</i>—Add to the liquid* in a test-tube about an equal volume of strong <math>H_2SO_4</math>, cool thoroughly, then pour a little strong <math>FeSO_4</math> solution gently down the inside of the tube so as to form a distinct layer on the surface of the acid liquid, a brown ring at the surface of contact of the two layers shows  <i>Presence of a nitrate.</i></p>
<p>4. <i>A mixture of CO and <math>CO_2</math> gases is given off; found by a drop of lime-water held in the tube being turned milky, and by the CO burning with a blue flame when kindled:—</i>  <i>Presence of an oxalate.</i></p>	<p>5. <i>A reddish-brown or violet color is produced in the acid, and no explosion or crackling occurs on warming:—</i>  <i>Prob. presence of a bromide or iodide.</i></p> <p>If on heating the mixture, colored vapor is evolved (see note), hold in the vapor the end of a glass rod covered with moist starch powder: the starch becomes brown:—  <i>Presence of a bromide.</i></p> <p>The starch becomes blue:—  <i>Presence of an iodide.</i></p> <p><i>Note.</i>—The vapor is much more easily obtained by mixing the substance with <math>MnO_2</math> powder before adding <math>H_2SO_4</math> and heating.</p>	<p>6. The substance blackens and evolves gases, amongst which <math>SO_2</math> is recognized by its smell and action on <math>K_2Cr_2O_7</math> solution:—  <i>Prob. presence of a tartrate.</i></p> <p>Confirm by (358).</p> <p>7. If none of the preceding acid radicles is found, add alcohol to the mixture of the substance with strong <math>H_2SO_4</math>, heat to boiling, and kindle the alcohol, it burns with a green edged flame:—  <i>Presence of a borate.</i></p> <p>Confirm by (357).</p>

**351.** After heating the tube containing the strong  $H_2SO_4$  and substance, let it stand for some time, then rinse it out, dry the inside thoroughly; if it is seen to be corroded and dimmed:—*Presence of a fluoride* is shown. The presence of a fluoride, if decisively indicated here, requires no confirmation.

\* Obtained by boiling the substance if a solid with water, and filtering or decanting the liquid, if the substance does not completely dissolve.

## ACTUAL EXAMINATION FOR THE ACID RADICLE.

**352.** The examination for the metal, or the preliminary examination for the acid radicle, will have detected any one of the commonly occurring acid radicles, except those whose detection is provided for below (353-358).

For the tests given below, use separate portions of the original substance.

**353.** Acidify a portion of the liquid (or if a solid, boil it) with HCl and add  $\text{BaCl}_2$ , or if Pb, Hg', or Ag has been found, acidify with  $\text{HNO}_3$  and add  $\text{Ba}(\text{NO}_3)_2$ , since HCl and chlorides would precipitate the above metals as chlorides. Either no precipitate forms—pass on to (354),—or a white precipitate forms which does not disappear on boiling: *Presence of a sulphate.*

**354.** Acidify a portion of the liquid, or if a solid is being examined heat the solid, with  $\text{HNO}_3$ ; filter off any precipitate or residue which may form, and add  $\text{AgNO}_3$  solution: either no precipitate forms (see note on p. 212), showing the absence of 'Cl, 'Br, 'I, 'Cy''' ( $\text{FeCy}_6$ ), '( $\text{Fe}_2\text{Cy}_{11}$ ), and (CyS)', pass on to (356); or a precipitate forms, which does not disappear on warming, see (355):

**355.** A precipitate forms. Note its color; and according as it is white, yellow, or orange, examine it by 1, 2, or 3:

<p><b>1. The precipitate is white, and may consist of <math>\text{AgCl}</math>, <math>\text{AgCy}</math> (<math>\text{Ag}_2\text{FeCy}_6</math>, or <math>\text{AgCyS}</math>). To a portion of the original solution add <math>\text{HCl}</math> in excess, and then <math>\text{Fe}_3\text{Cl}_6</math>:</b></p>		
<p><b>A deep blue precipitate:</b> <i>Presence of a ferrocyanide.</i></p>	<p><b>A blood-red coloration.</b> <i>Presence of a sulphocyanide.</i></p>	
<p><b>If <math>\text{Fe}_3\text{Cl}_6</math> has yielded neither a precipitate nor coloration (indicating absence of a ferro-, ferri-, and sulphocyanide), decant the liquid from the precipitate caused by <math>\text{AgNO}_3</math>, pour upon the precipitate some strong <math>\text{HNO}_3</math> and boil:</b></p>		
<p><b>The precipitate remains undissolved:</b> <i>Presence of a chloride.</i></p>	<p><b>The precipitate dissolves:</b> <i>Presence of a cyanide.</i> Confirm the presence of a cyanide in a portion of the original solution by heating it with <math>\text{FeSO}_4</math>, <math>\text{Fe}_3\text{Cl}_6</math>, and <math>\text{KHO}</math>, and acidifying with <math>\text{HCl}</math>. A blue precipitate:—<i>Presence of a cyanide.</i></p>	
<p><b>2. The precipitate is yellow, and may consist of <math>\text{AgBr}</math> or <math>\text{AgI}</math>. Test some of the original solution for an iodide by adding to it starch solution and <math>\text{KNO}_3</math> solution, and then acidifying (if not acid) with <math>\text{HCl}</math>: a deep blue coloration:</b>  <i>Presence of an iodide.</i></p>	<p><b>If an iodide is not thus detected, add to another portion of the original solution several drops of <math>\text{Cl}</math> water; if the liquid becomes yellow, and on being shaken with <math>\text{CS}_2</math> colors that liquid reddish-brown, it shows:</b>  <i>Presence of a bromide.</i></p>	<p><b>3. Orange-red precipitate.</b> To a portion of the original solution add <math>\text{HCl}</math> in excess, then <math>\text{FeSO}_4</math> solution; a deep blue precipitate shows:  <i>Presence of a ferri-cyanide.</i></p>

*Note.*—If  $\text{Hg}$  has been found,  $\text{HgCy}_2$  may be present, notwithstanding the formation of no precipitate with  $\text{AgNO}_3$ . It is detected by (306, note 2, p. 178). If  $\text{HgCy}_2$  is not present pass on to (356).

**356.** Acidify some of the original solution if it is not already acid, with dilute  $\text{HNO}_3$ ; filter if necessary, add a small quantity of this acid solution to some  $\text{AmHMoO}_4$  solution, stir well and warm gently:

<b>A yellow precipitate shows presence of a phosphate.</b>	<b>If no precipitate forms, heat to boiling; the formation of a yellow precipitate shows presence of an arsenate.</b> <b>If no precipitate forms even on boiling the liquid, pass on to (357).</b>
--	---

*Note.*—The presence of a phosphate or arsenate may be confirmed by dissolving the yellow precipitate, separated by filtration or decantation, by pouring upon it a little  $\text{AmHO}$ ; then adding to this

## 357, 358 a.] EXAMINATION FOR ACID RADICLE. 213

solution  $\text{AmCl}$  and  $\text{MgSO}_4$ , a crystalline precipitate will form which is to be filtered off and washed with a little cold water; on pouring upon this precipitate on the filter several drops of  $\text{AgNO}_3$  solution, the precipitate becomes *yellow* if phosphate is present, and *brown* if arsenate is present. The presence of arsenate is further confirmed by a yellow precipitate ( $\text{As}_2\text{S}_3$ ) appearing with  $\text{H}_2\text{S}$  (337) when the liquid is boiled.

**357.** Acidify a portion of the original solution with  $\text{HCl}$ , dip into it a piece of turmeric-paper, and dry the paper at a gentle heat; if the paper becomes *reddish-brown* when dry, and changes to *dingy-green* or *blue-black* when moistened with  $\text{AmHO}$ , the *presence of a borate* is shown. If no reaction is obtained pass on to (358).

**358.** The acid radicles, *one* of which may still be present, are 'T, ' $\text{C}_2\text{O}_4$ , and 'F. If the metal found belongs to Groups I, II, III, or IV, the examination must be begun at (358 b); if the metal present is K, Na, or  $\text{NH}_4$ , examine by (358 a).

**358 a.** Make the solution just alkaline with  $\text{AmHO}$ , then add  $\text{CaCl}_2$ , [or  $\text{Ca}(\text{NO}_3)_2$ ] solution in some quantity, shake well and let it stand if no precipitate forms at once; the preliminary examination will usually have indicated which of these acid radicles is present, and thus show according to which of the three columns below, the precipitate is to be further tested:

I. A crystalline precipitate, appearing after some time, shows probable presence of a tartrate.

(Confirm either by 1 or 2 below:

1. Decant the liquid from the precipitate, add a little water and a single drop of  $\text{AmHO}$ , drop in a small crystal of  $\text{AgNO}_3$  and warm very carefully; a mirror is formed below the crystal, *presence of a tartrate*. This test may be tried also with the original solution by (322).
2. On acidifying some of the neutral original solution with  $\text{H}\bar{\text{A}}$ , and adding alcohol and  $\text{K}\bar{\text{A}}$  (unless K or  $\text{NH}_4$  has been detected in the substance), and stirring or shaking well, the formation of a crystalline precipitate shows *presence of a tartrate*.

II. A white pulverulent precipitate shows probable presence of an oxalate.

The presence of an oxalate, already indicated by the test in 350, 4, must be confirmed either by warming some of the original substance with  $\text{MnO}_2$  and  $\text{H}_2\text{SO}_4$ , and testing for  $\text{CO}_2$  gas (319); or by the more tedious method of filtering off and drying some of the above precipitate of  $\text{CaC}_2\text{O}_4$  and testing for a carbonate with  $\text{HCl}$  (318 a).

III. A gelatinous flocculent precipitate shows probable presence of a fluoride.

Confirm by filtering and warming some of the precipitate (or of the substance, if solid) in a small cup of lead or platinum, with strong  $\text{H}_2\text{SO}_4$ ; fumes will be evolved which etch a properly prepared watch-glass (296).

**358 *b*.** An oxalate or fluoride of Ba, Sr, Ca, or Mg will have been already detected in par. 339. A tartrate of any one of these metals is found by gently heating the finely powdered substance (if it is a solid, or neutralizing the solution, if a liquid) with *very dilute* AmHO and a crystal of AgNO<sub>3</sub>; the formation of a mirror shows *presence of a tartrate*. If a metal of Groups I, II, or III has been found, the metal must be separated before testing for the acid radicle.

If the metal belongs to Group I or II, make the original solution just acid with HCl, and add H<sub>2</sub>S solution or pass H<sub>2</sub>S (500 *a*) into the liquid for about five minutes; filter off the precipitate and add more H<sub>2</sub>S to the filtrate; if no more precipitate forms\* boil in a porcelain dish until the liquid ceases to smell of H<sub>2</sub>S, and examine by (358 *a*).

If the metal belongs to Group III add AmCl, AmHO till alkaline and Am<sub>2</sub>S; boil and filter, add to the filtrate Am<sub>2</sub>S; if no more precipitate forms\* add H $\bar{A}$  in excess, boil, filter off S if necessary, and examine by (358 *a*).

THE LIQUID FOR ANALYSIS IS ALKALINE (334, II).

**359.** The following special course of examination should be pursued if the liquid under analysis is alkaline in reaction, since it is possible in such a case that *two metals* may be present, one of them as a constituent of the acid radicle. In the preceding course the detection of *one* metal only is provided for.

An alkaline solution may contain the oxide of any one of the metals Pb, Zn, As, Sn, Sb, Al, or Cr, together with the oxide of one of the metals K or Na; it may also contain K or Na as silicate. The detection of such substances is provided for in paragraph 360. Certain other salts of K, Na, and NH<sub>4</sub>, which also have an alkaline reaction, may however be analyzed by the usual plan (336 *et seq.*), as will be seen below (360 *et seq.*).

---

\* If any further precipitate forms add more of the reagent, filter, and test again by adding more of the reagent; this addition of reagent, filtration, and testing the filtrate is repeated until no further precipitate forms on addition of the reagent.

**360.** To a small portion of the alkaline solution add several drops of dilute  $\text{HNO}_3$ ; either no precipitate will form (see column 2, p. 216), or a precipitate forms; in the latter case note the color and appearance of the precipitate, and proceed to test further as directed in 1 of table, p. 216.

**360 a.** The presence of Na, K or  $\text{NH}_4$  will probably have been already detected in the preliminary examination;  $\text{NH}_4$  or Na, if thus found, require no further confirmation.

The presence of K will usually not require confirmation, but if any doubt is felt concerning its presence, confirm by adding to a part of the original solution excess of HCl, evaporating to dryness, stirring and warming the residue with a *few drops* of water and several drops of HCl, decanting or filtering if necessary from any undissolved residue, and stirring several drops of the cold, clear liquid on a watch-glass with  $\text{PtCl}_4$  solution for some time; a yellow precipitate forming on the lines rubbed by the rod confirms the *presence of K*.



<p>1. <i>A precipitate forms on addition of the first few drops of <math>\text{HNO}_3</math>.</i> To another portion of the solution add <math>\text{H}_2\text{S}</math>-water drop by drop, and note if any precipitate forms, then add <math>\text{HCl}</math> in excess: if a precipitate is formed (see 8 below), if no precipitate appears (see 4 below):</p> <p>2. <i>No precipitate is produced by addition of <math>\text{HNO}_3</math></i> except possibly a slight milkiness due to the separation of <math>\text{S}</math>. Acidify another portion with <math>\text{HCl}</math> and employ it as the original solution for the examination for the metal by pars. 336 <i>et seq.</i></p> <p>It must be remembered that in this case a <i>silicate</i> may be present.</p> <p>3. <i>A precipitate is formed by addition of <math>\text{H}_2\text{S}</math> and excess of <math>\text{HCl}</math>:</i> note its color and appearance:</p>					
<p><i>A black precipitate:</i> <i>Presence of Pb.</i></p> <p>Confirm by adding to another portion of the solution <math>\text{H}_2\text{A}</math> in excess, then <math>\text{K}_2\text{CrO}_4</math>; a yellow precipitate: <i>Presence of Pb.</i></p>	<p><i>A white precipitate forms on addition of <math>\text{H}_2\text{S}</math>, but is dissolved by the <math>\text{HCl}</math>:</i> <i>Presence of Zn.</i></p> <p>Confirm by shaking well, and letting the precipitate settle; on decanting the liquid and boiling the precipitate with strong <math>\text{HCl}</math>, it is not dissolved: <i>Presence of As as arsenite.*</i></p>	<p><i>A bright yellow very flocculent precipitate:</i> <i>Presence of As as arsenide.</i></p> <p>Confirm by shaking well, and letting the precipitate settle; on decanting the liquid and boiling the precipitate with strong <math>\text{HCl}</math>, it dissolves readily: <i>Presence of Sn as stannate.†</i></p>	<p><i>A yellow precipitate, not separating in large flocks:</i> <i>Presence of Sn as stannate.</i></p> <p>Confirm by warming, allowing the precipitate to settle, decanting the liquid if possible, or filtering, and boiling the precipitate with strong <math>\text{HCl}</math>; a piece of <math>\text{Zn}</math> is resting; a black stain on the <math>\text{Pt}</math> appears after a few minutes: <i>Presence of Sb as antimonate.</i></p>	<p><i>An orange-red precipitate:</i> <i>Presence of Sb.</i></p> <p>Confirm by adding <math>\text{HCl}</math> in excess to a portion of the original solution, and boiling; pour some of this solution, when cool, upon a piece of platinum foil on which a piece of <math>\text{Zn}</math> is resting; a black stain on the <math>\text{Pt}</math> appears after a few minutes: <i>Presence of Sb as antimonate.</i></p>	<p><i>A semitransparent gelatinous precipitate.</i> <i>Presence of a silicate.</i></p> <p>Confirm by evaporating some of the original solution to dryness after acidifying it with <math>\text{HCl}</math>; warm the residue in the dish with <math>\text{HCl}</math>; an insoluble powder is left: <i>Presence of a silicate.</i></p>
Proceed to (360 a).	Proceed to (360 a).	Proceed to (360 a).	Proceed to (360 a).	Proceed to (360 a).	Proceed to (360 a).

<p>4. <i>No precipitate is formed by addition of <math>\text{H}_2\text{S}</math> and excess of <math>\text{HCl}</math>:</i></p> <p><i>The original solution was colorless:—</i>acidify a portion with <math>\text{HCl}</math> and evaporate to dryness, then warm the residue with <math>\text{HCl}</math>:</p> <p><i>The residue dissolves completely:—</i> Add to the solution <math>\text{AmH}_2\text{O}</math> in excess, a white gelatinous precipitate shows: <i>Presence of Al.</i> Proceed to (360 a).</p>		<p><i>The original solution was colorless:—</i>acidify a portion with <math>\text{HCl}</math> and evaporate to dryness, then warm the residue with <math>\text{HCl}</math>:</p> <p><i>The residue is not completely dissolved:</i> <i>Presence of a silicate.</i> <i>The residue is not completely dissolved:</i> <i>Presence of a silicate.</i> Proceed to (360 a).</p>
--	--	--

4. *No precipitate is formed by addition of  $\text{H}_2\text{S}$  and excess of  $\text{HCl}$ :*

*The original solution was green:*

*Presence of Cr.*

*The presence of Cr is confirmed* by the formation of a pale green precipitate on addition of the first few drops of  $\text{HNO}_3$ . It will probably also have been found in the preliminary examination by the borax bead and by fusion with  $\text{KNO}_3$  and  $\text{Na}_2\text{CO}_3$ . Its presence may be further confirmed by adding to some of the original solution excess of  $\text{KHO}$ , then  $\text{PbO}_2$  and boiling, on acidifying with  $\text{H}_2\text{A}$  yellow  $\text{PbAs}_2$  is precipitated.

Proceed to (360 a).

\* Tests (180, 189) may also be used here with a portion of the original solution.

† Confirmed also by  $\text{AmHO}$  giving a precipitate in the solution formed by adding excess of  $\text{HCl}$  and boiling, whereas it gives no precipitate with  $\text{As}_2\text{O}_3$  solution similarly prepared.

*The original solution was colorless:*—acidify a portion with  $\text{HCl}$  and evaporate to dryness, then warm the residue with  $\text{HCl}$ :

*The residue dissolves completely:*—Add to the solution  $\text{AmHO}$  in excess, a white gelatinous precipitate shows:  
*Presence of Al.*

*The residue is not completely dissolved:*  
*Presence of a silicate.*

Proceed to (360 a).

Proceed to (360 a).

## THE SUBSTANCE POSSESSES METALLIC LUSTRE.

**366.** This shows the probable absence of an acid radicle. The only non-metallic elements likely to be present are free carbon as graphite, free iodine, or combined sulphur, since several sulphides are decidedly metallic in appearance.\* The presence of the metals Ni, Co, Fe in the free state, and of  $\text{Fe}_3\text{O}_4$ , may be shown by the substance being attracted by a magnet. Pb, or graphite, will be recognized by marking paper when rubbed upon it, as a black lead-pencil does; the mark due to Pb disappears if wetted with dilute  $\text{HNO}_3$ .

The preliminary examination may be advantageously modified and curtailed as shown below.

Experiment.	Observation.	Inference.
I. Heat a portion of the substance in a small ignition-tube (10).	1. A black lustrous mirror forms, . 2. A yellow sublimate forms, melting, when heated, to brown drops, 3. Violet vapor is given off, condensing to black scales on the sides of the tube, . If the substance sublimes entirely in violet vapors, . . .	Presence of As. Presence of S. Presence of I. Iodine alone is present.
II. Heat a portion in the inner blowpipe flame in a cavity scooped on a piece of wood-charcoal.	1. Smell of garlic, . 2. If an incrustation forms, refer to par. (333 V), where in the second and third column will be found the appearance of the incrustation and the metal whose presence it indicates. 3. Smell of burning sulphur, . . 4. A green flame coloration, . .	Presence of As. Presence of S. Presence of Cu.
III. Place a portion of the substance in a piece of hard glass tube, open at both ends, and heat it strongly in the Bunsen flame, and afterwards in the blowpipe flame, if necessary (Fig. 35, p. 132).	1. A gas is evolved from the upper end which smells of burning sulphur, and turns a piece of filter-paper moistened with $\text{K}_2\text{Cr}_2\text{O}_7$ solution green, . . 2. The substance burns completely but slowly away, glowing when strongly heated. This effect is best seen by heating it strongly on Pt foil by directing the blowpipe flame on the under surface of the foil, .	Presence of S. Presence of C.

\* The most common sulphides which possess metallic lustre are "copper pyrites," and "iron pyrites," which resemble brass in appearance; and "galena," resembling lead.

After completing the preliminary examination proceed as directed under (332), using however strong HCl and strong  $\text{HNO}_3$  instead of the dilute acids.

No acid radicle need be tested for unless S has been found in the preliminary examination, when in a portion of the  $\text{HNO}_3$  solution  $\text{H}_2\text{SO}_4$  is tested for by  $\text{BaCl}_2$ ; if a white precipitate forms insoluble on boiling, it confirms the *presence of S*.

---

### THE SUBSTANCE IS INSOLUBLE BOTH IN WATER AND IN ACIDS.

**367.** It may consist of any one of the following substances, those included in brackets being less likely to occur than the others, since they are soluble in sufficient water or acid on heating. As will be seen below the color serves to give some notion as to what substance is present.

1. *White*:  $\text{BaSO}_4$ ,  $\text{SrSO}_4$ ,  $(\text{CaSO}_4)$ ,  $\text{PbSO}_4$ ,  $(\text{PbCl})_2$ ,  $\text{AgCl}$ ,\*  $\text{SiO}_2$ ,  $(\text{Al}_2\text{O}_3 \text{ ignited})$ ,  $\text{SnO}_2$ ,  $\text{Sb}_2\text{O}_3$ ,  $\text{Sb}_2\text{O}_4$ ,  $\text{CaF}_2$ .
2. *Dark-colored*: ignited  $\text{Fe}_2\text{O}_3$  (dark-brown or black), ignited  $\text{Cr}_2\text{O}_3$  (dark-green),  $\text{FeCr}_2\text{O}_4$  and ignited  $\text{PbCrO}_4$  (dark-brown), C (black).
3. *Yellow*: S,  $\text{AgBr}$ ,\*  $\text{AgI}$ ,\*  $(\text{PbI}_2)$ .

An insoluble silicate may also be present.

For a fuller description of these substances see (475). According to the color of the substance examine it by (368 ; 1, 2, or 3).

**368.** The preliminary tests given below will indicate or confirm the composition of the substance very rapidly, and may be substituted for those described in (333).

1. *The substance is white* (see 367, 1): Pour upon a small portion of it a drop of  $\text{Am}_2\text{S}$ ; if it blackens, the presence of Pb or Ag is probable; if it becomes yellow or orange-red, the presence of Sn or Sb is probable.

---

\* Darkens quickly in sunlight, and slowly in ordinary daylight.

Experiment.	Result.	Inference.
I. Moisten a clean loop of platinum-wire, immerse it in the powdered substance, and heat the adhering powder for a short time strongly in the <i>inner</i> blowpipe flame; moisten the loop with a drop of strong HCl, and hold it in the Bunsen flame.	A <i>yellowish-green</i> flame.  A <i>crimson</i> flame, appearing <i>deep red</i> through the indigo-prism. A <i>reddish</i> flame, appearing <i>dusky green</i> through the indigo-prism.	<i>Presence of Ba.</i> [Test another portion for $\text{SO}_4$ by (III), and see note below.] <i>Presence of Sr</i> [ " " ]  <i>Presence of Ca</i> [ " " ] If $\text{SO}_4$ is not found, test for $\text{F}^-$ by (351 and 296). <i>Note.</i> —The above result may be confirmed by (369).
II. Mix some of the finely powdered substance with powdered $\text{Na}_2\text{CO}_3$ and $\text{K}_2\text{Cr}_2\text{O}_7$ in a small cavity made in a piece of wood-charcoal, and heat the mixture in the inner blowpipe flame.	White metallic <i>malleable</i> globules, which if taken upon the point of a pen-knife are found to mark paper as black lead does, and if dissolved in $\text{HNO}_3$ give a white precipitate with $\text{H}_2\text{SO}_4$ ; a yellow incrustation is also formed on the charcoal.  White metallic <i>malleable</i> globules, which do not mark paper; they dissolve in HCl, the solution giving a white precipitate with $\text{HgCl}_2$ . Slight incrustation or none. White metallic <i>brittle</i> globules, which, if dissolved in boiling HCl, give an orange-red precipitate with $\text{H}_2\text{S}$ ; white incrustation. White scales, no incrustation; the metal is insoluble in HCl, and if dissolved in $\text{HNO}_3$ gives no precipitate with $\text{H}_2\text{SO}_4$ , but a white precipitate with HCl; the original substance is instantly blackened by a drop of $\text{Am}_2\text{S}_3$ , and is dissolved on being warmed with $\text{AmHO}$ (212-216).	<i>Presence of Pb</i> as chloride, sulphate or chromate. Test for chloride by boiling some of the substance with much water, adding $\text{HNO}_3$ and $\text{AgNO}_3$ ; white precipitate. <i>Presence of PbCl_2.</i> Test for sulphate by (III). (See also 369.) Test for chromate by fusing in a porcelain crucible, with fusion mixture; yellow mass on cooling: <i>Presence of PbCrO_4.</i>  <i>Presence of SnO_2.</i>  <i>Presence of Sb_2O_3 or Sb_2O_4.</i>    <i>Presence of AgCl.</i> [See also (369 a) ].
III. Fuse a portion of the substance mixed with $\text{Na}_2\text{CO}_3$ in fine powder, in a cavity scooped upon a piece of wood-charcoal, in the inner blowpipe flame, <i>produced from a spirit-lamp flame.</i>	The cooled mass, if detached from the charcoal, placed upon a bright silver coin, and moistened with water, gives, when crushed with the blade of a knife, a black stain.	<i>Presence of a sulphate.</i>

*Note.*—The presence of  $\text{PbSO}_4$  may be confirmed by pouring upon the original substance  $\text{H}\bar{\text{A}}$ , then excess of  $\text{AmHO}$ , and boiling; the substance will dissolve: acidify a portion of the solution with  $\text{H}\bar{\text{A}}$ , and add  $\text{K}_2\text{CrO}_4$ , a yellow precipitate: *Presence of Pb*. Acidify another portion with  $\text{HCl}$ , and add  $\text{BaCl}_2$ , a white precipitate insoluble on boiling: *Presence of  $\text{SO}_4$* .

If none of the substances mentioned in the above table have been detected, test for  $\text{Al}_2\text{O}_3$  by heating the substance strongly on charcoal in the outer blowpipe flame after moistening it with  $\text{Co}(\text{NO}_3)_2$  solution, a blue mass shows *presence of  $\text{Al}_2\text{O}_3$* .

If  $\text{Al}_2\text{O}_3$  is not found fuse some of the substance in a clear bead of  $\text{NaAmHPO}_4$ , it floats undissolved: *Presence of  $\text{SiO}_2$* . Proceed to (369 b).

2. *The substance is dark-colored.* (See 367, 2.)

Place the substance on a piece of platinum foil and heat the foil strongly on its under surface by the blowpipe flame :

<p><i>The substance burns away slowly but completely:</i></p> <p><i>Presence of C.</i></p>	<p><i>The substance does not burn away; place upon it three or four times as much powdered <math>\text{Na}_2\text{CO}_3</math> and <math>\text{KNO}_3</math>, and fuse for some time.</i></p>
	<p>1. The substance dissolves, forming a yellow mass when cold:—<i>Presence of <math>\text{Cr}_2\text{O}_3</math></i>.</p> <p>2. The substance remains undissolved as a dark-brown powder, and the mass on cooling is white:—<i>Presence of <math>\text{Fe}_2\text{O}_3</math></i>.* Confirm by boiling the mass with water, filtering and fusing the brown residue into a borax bead (100).</p>

3. *The substance is yellow* (367, 3).

Heat a portion of it strongly in a small ignition-tube; one of the following results will occur :

<p><i>It fuses and sublimes, the sublimate being yellow and melting to reddish-brown drops when heated:</i></p> <p><i>Presence of S.</i></p> <p><i>Note.</i>—If the substance sublimes entirely it consists only of sulphur.</p>	<p><i>It fuses but does not sublime.</i></p> <p>Examine a portion of the substance for <math>\text{Ag}</math> and <math>\text{Pb}</math> by (368 II), and another portion for <math>\text{Br}</math> and <math>\text{I}</math> by (369 a).</p>
--	--

\*  $\text{FeCr}_2\text{O}_4$  shows both reactions 1 and 2, since it contains both  $\text{Fe}$  and  $\text{Cr}$ .

**369.** Fuse a portion of the substance, mixed with two or three times as much fusion mixture, for several minutes on a piece of platinum foil (or if Pb is present, in a porcelain crucible) in the blowpipe flame; boil the cool mass with water until only a white powder remains undissolved, filter or decant:

<i>Filtrate:</i> acidify with HCl and add $\text{BaCl}_2$ , a white precipitate which does not disappear on boiling: <i>Presence of <math>\text{SO}_4</math>.</i>	<i>Residue:</i> after washing several times with boiling water dissolve it in a little hot HCl and examine the solution for the metal by (336 <i>et seq.</i> ).
--	---

**369a.** Fuse a portion of the finely powdered substance for about five or ten minutes with two or three times as much fusion mixture in a porcelain crucible in the blowpipe flame; as soon as the substance is cool, boil it with water for some time, filter, or decant if possible:

<i>The residue may be used to confirm the presence of Ag, if the presence of that metal is not fully proved in (368 II). Wash the residue well, dissolve it by warming with a little dilute <math>\text{HNO}_3</math>, add HCl, shake well, let the precipitate settle, decant, and show the precipitate to be soluble in AmHO.</i>	<i>The clear solution is acidified with <math>\text{HNO}_3</math> and divided into two portions.</i>	
	1. To the first portion add $\text{AgNO}_3$ ; a perfectly white precipitate, easily soluble in excess of AmHO, shows <i>Presence of chloride.</i> If the precipitate is yellow and not easily soluble in AmHO, see next column.	2. To the second portion add $\text{CS}_2$ sufficient to form a large drop at the bottom of the liquid, then Cl-water drop by drop, constantly shaking up the liquid and $\text{CS}_2$ well together:  <div style="display: flex; justify-content: space-between;"> <div> <i>The <math>\text{CS}_2</math> is colored reddish-brown: Presence of bromide.</i> </div> <div> <i>The <math>\text{CS}_2</math> is colored violet showing: Presence of iodide. [The presence of HI may be confirmed by (269)].</i> </div> </div>

**369b.** Fuse a portion of the powdered substance with fusion mixture, as directed in (369a). Boil the cold crucible in a porcelain dish with dilute  $\text{HNO}_3$  until the mass is detached entirely, then evaporate the liquid to complete dryness: warm the cool residue with dilute  $\text{HNO}_3$ , an insoluble residue shows or confirms the *presence of  $\text{SiO}_2$ .*

The solution, filtered if necessary from  $\text{SiO}_2$ , is tested for the metal present (336 *et seq.*), bearing in mind that K and Na cannot possibly be present in the insoluble substance, since they form soluble silicates.

## ANALYSES OF SIMPLE SALTS, SHOWING HOW TO ENTER RESULTS.

### A. Liquid given for Analysis.

#### *Preliminary Examination for the Metal.*

Experiment.	Observation.	Inference.
1. Noted the color of the liquid.	Blue . . . . .	Presence of Cu.
2. Dipped into it a piece of blue litmus-paper.	The paper was turned red, but on addition of $\text{Na}_2\text{CO}_3$ no effervescence occurred . . .	Presence of a salt with acid reaction.
3. Evaporated a few drops slowly upon a watch-glass.	Pale blue residue left . .	Presence of some dissolved solid.
Ignited the residue strongly.	Blackened without smell of burning . . .	Absence of organic acid radicles.
4. Added KHO solution and boiled.	No smell of $\text{NH}_3$ . . .	Absence of $\text{NH}_4$ .
5. Dipped a loop of platinum wire into the solution and held it in the flame,	<i>Bright green</i> flame; on moistening the wire with strong HCl and again holding it in the flame it gave a blue coloration . . .	Presence of Cu.
6. Dipped a clear borax bead into a portion of the solution, dried it by a gentle heat, dipped it again and fused in the outer and inner blow-pipe flame.	In outer flame the bead was <i>green</i> whilst hot, <i>blue</i> when cold. In inner flame it became colorless. }	Presence of Cu.

#### *Examination for the Metal.*

To a portion of the liquid added HCl :

No precipitate. Absence of Group I.	<p>Added strong <math>\text{H}_2\text{S}</math>-water :</p> <p><i>A brownish-black</i> precipitate. Presence of Sn'' Hg'', Bi, Pb, or Cu.</p> <p>Allowed the precipitate to settle, poured off as much of the liquid as possible, and boiled the precipitate with excess of NaHO, it remained undissolved.</p> <p>Since the original solution was blue, a clean knife-blade was dipped into a portion of it acidified with <math>\text{H}_2\text{SO}_4</math>, red copper was deposited on the steel :</p> <p style="text-align: center;"><i>Presence of Cu.</i></p>
--	--

*Preliminary Examination for the Acid Radicle.*

Experiment.	Observation.	Inference.
1. Added dilute $H_2SO_4$ and warmed.	No gas was evolved.	Absence of carbonate, sulphite, sulphide, and nitrite.
2. Added strong $H_2SO_4$ and warmed.	No gas was evolved. And no red fumes on adding Cu clippings.	Absence of chlorate. Absence of nitrate.

*Examination for the Acid Radicle.*

Added HCl and  $BaCl_2$ ; a white precipitate was produced which did not disappear on boiling: *Presence of sulphate.*

*Found Cu,  $SO_4$ .*

*B. Liquid given for Analysis.**Preliminary Examination for the Metal.*

Experiment.	Observation.	Inference.
1. Noted the color of the liquid.	Light yellow.	Presence of a neutral chromate.
2. Dipped red litmus-paper into the liquid.	It was turned <i>faintly</i> blue.	Probable presence of an alkali salt.
3. Evaporated a few drops to dryness upon a watch-glass. Ignited the residue strongly.	A yellow residue remained. No blackening.	Presence of some dissolved solid. Absence of organic acid radicle.
4. Boiled a portion with excess of KHO.	No smell of $NH_3$ .	Absence of $NH_4$ .
5. Dipped a loop of platinum wire into the solution, and held it in the Bunsen flame.	A pale violet flame, appearing reddish-violet through the indigo-prism.	Presence of K.
6. Dipped a clear borax-bead into the solution, and fused in the outer and inner blowpipe flames.	Outer flame: <i>brown</i> , hot; <i>green</i> , cold. Inner flame: <i>green</i> , hot and cold. <i>Brown</i> color not reappearing in the outer flame.	Presence of Cr as a chromate.

*Examination for the Metal.*

Since the liquid was alkaline in reaction, added to a



small portion of it a few drops of  $\text{HNO}_3$ , until it became acid, this produced no precipitate.

To another portion added  $\text{HCl}$ , the liquid turned from yellow to orange-red, indicating presence of a neutral chromate :

No precipitate. Absence of Group I.	Then added $\text{H}_2\text{S}$ -water and warmed :		
	The liquid became green, and white S was deposited :  <i>Presence of a chromate.</i>	To another portion of the original solution added $\text{AmCl}$ , then $\text{AmHO}$ ; no precipitate formed ; then added $\text{Am}_2\text{S}$ , and boiled :	
		<i>No precipitate</i> in the cold, but on boiling, a green flocculent precipitate gradually formed, evidently consisting of $\text{Cr}_2\text{H}_2\text{O}_4$ reduced from the chromate.	Added to another portion of the solution $\text{AmCl}$ , $\text{AmHO}$ , and $\text{Am}_2\text{CO}_3$ :
			<i>No precipitate.</i> Absence of Group IV.  Added to the same portion $\text{Na}_2\text{HPO}_4$ and stirred :  <i>No precipitate.</i> Absence of <i>Mg</i> .

Since K was found by the flame coloration in the preliminary examination, its presence was confirmed by stirring a portion of the solution, to which a few drops of  $\text{HCl}$  had been added, with  $\text{PtCl}_4$  on a watch-glass ; a yellow crystalline precipitate forming on the lines rubbed by the rod : *Presence of K.*

*Found K,  $\text{CrO}_4$ .*

#### C. A white crystalline substance given for Analysis.

Boiled a small portion with water ; it dissolved completely.

*Preliminary Examination for the Metal.*

Experiment.	Observation.	Inference.
1. Heated in small dry test-tube.	The substance gave off red fumes and oxygen gas, which inflamed a glowing splinter of wood. It left a yellow residue,	Presence of a nitrate of a heavy metal. Presence of Pb, Sn, or Bi.
2. Dipped a loop of moistened platinum-wire into the powdered substance, moistened with strong HCl and held in the Bunsen flame.	A pale blue flame, . .	Presence of As, Sb, Pb.
3. Heated a small portion in a cavity on wood charcoal in the inner blowpipe flame.	Deflagration occurred, A white malleable globule remained, which easily marked paper. A yellow incrustation, . . . .	Presence of a nitrate or chlorate.
4. Fused on charcoal in the inner blowpipe flame with $\text{Na}_2\text{CO}_3$ and KCy.	Same result as with 3, .	Presence of Pb. Presence of Pb.

*Examination for the Metal in Solution.*

Boiled a portion of the powdered substance with water, cooled, added HCl to a part of the clear solution, a white precipitate was formed, which disappeared on boiling, but appeared again in the crystalline form on cooling the liquid (*probable presence of Pb*).

Confirmed the presence of Pb by decanting the liquid and boiling the precipitate with  $\text{AmHO}$ , it remained white and did not dissolve; decanted the ammoniacal liquid, dissolved the precipitate in a little  $\text{H}\bar{\text{A}}$ , and added  $\text{K}_2\text{CrO}_4$ , a yellow precipitate: *Presence of Pb*.

Since a nitrate was found in the preliminary examination, its presence was confirmed by adding strong  $\text{H}_2\text{SO}_4$  to the aqueous solution of the substance, and pouring solution of  $\text{FeSO}_4$  carefully upon the cooled liquid in a test-tube; a brown ring formed on the surface of the acid: *Presence of  $\text{NO}_3$* .

Also some of the solution when heated with Cu and strong  $\text{H}_2\text{SO}_4$ , evolved reddish-brown fumes: *Presence of  $\text{NO}_3$* .

*Found:  $\text{Pb}, \text{NO}_3$ .*

D. *A white powder emitting no smell, given for Analysis.*

Boiled some of the powdered substance with water, it did not dissolve: added a few drops of strong HCl and boiled, the substance dissolved completely (absence of Group I) without effervescence.

*Preliminary Examination for the Metal.*

Experiment.	Observation.	Inference.
1. Heated in a small dry test-tube.	No change, . . . .	Absence of volatile and fusible substances, and of water of crystallization, etc.
2. Dipped a moistened loop of platinum-wire into the powdered substance, moistened the powder with strong HCl, and held it in the Bunsen flame.	No flame coloration, .	Probable absence of K, Na, Ba, Sr, Ca, Cu, etc.
3. Heated a portion of the powder strongly on charcoal in the inner blowpipe flame.	The substance did not fuse. A white luminous residue, which, when moistened on red litmus-paper, showed an alkaline reaction,	Absence of alkali salt.  Presence of Ba, Sr, Ca, Mg.

*Examination of the Solution.*

To a portion of the HCl solution added  $H_2S$  water; no precipitate:—*Absence of Group II.*

To another portion of the HCl solution added  $AmCl$  and  $AmHO$  in excess, a white flocculent precipitate; then added  $Am_2S$ , the color of the precipitate was unchanged.

To a few drops of the HCl solution added KHO drop by drop, until a white flocculent precipitate formed, then added excess of KHO and boiled, the precipitate did not dissolve, and it did not become discolored in the air:—*Presence of Ba, Sr, Ca, Mg as phosphate, oxalate, etc.*

Added a few drops of the original HCl solution to some  $AmHMoO_4$  solution in a test-tube, and warmed gently, a yellow precipitate:—*Presence of  $PO_4$ .*

To another portion of the cold HCl solution added  $AmHO$ , drop by drop, until a precipitate formed which did not dissolve when shaken, then added  $H\bar{A}$  until this

precipitate dissolved when the liquid was well shaken. Some small crystals of  $\text{NaA}$  were then dissolved in the cold liquid by shaking it, and  $\text{Fe}_2\text{Cl}_6$  added until the liquid appeared red; it was then boiled and immediately filtered. To the filtrate  $\text{AmHO}$  was added in excess (no precipitate), then  $\text{Am}_2\text{CO}_3$ , a white precipitate formed:—*Presence of Ba, Sr, or Ca.*

Filtered and dissolved the precipitate by pouring upon it a few drops of boiling  $\text{HCl}$ ; dipped a loop of platinum-wire into the solution, and held it in the Bunsen flame, a reddish flame, appearing dingy-green through the indigo-prism:—*Presence of Ca.*

A few drops of the solution boiled with  $\text{CaSO}_4$  gave no precipitate:—absence of Ba and Sr, therefore the presence of Ca was confirmed by adding to the rest of the solution excess of  $\text{AmHO}$ , then  $\text{Am}_2\text{C}_2\text{O}_4$ , a white precipitate:—*Presence of Ca.*

*Found Ca,  $\text{PO}_4$ .*

*E. A white odorless powder given for Analysis.*

Boiled a portion of the substance with water, it did not dissolve: added  $\text{HCl}$  and heated again, it dissolved completely (absence of Group I) without effervescence.

*Preliminary Examination.*

Experiment.	Observation.	Inference.
1. Heated a portion in a small dry test-tube.	The substance blackened slightly; it evolved $\text{CO}_2$ which turned a drop of lime-water milky. The residue in the test-tube effervesced with $\text{HCl}$ , whereas the original substance did not.	Presence of an organic salt of K, Na, Ba, Sr, Ca, or Mg.
2. Held a portion of the substance on a loop of platinum-wire in the Bunsen flame; then moistened with $\text{HCl}$ and again held in the flame.	A red coloration, appearing <i>dirty green</i> through the indigo-prism.	Presence of Ca.
3. Heated on charcoal in inner blowpipe flame.	A white luminous mass remained, which when moistened on red litmus-paper, turned it blue.	Presence of Ba, Sr, Ca, or Mg.

*Examination for the Metal.*

Added to a portion of the HCl solution strong  $\text{H}_2\text{S}$ -water.

<p><i>No precipitate:</i> Absence of Group II.</p>	<p>To another portion of the HCl solution added <math>\text{AlnCl}</math> then <math>\text{AmHO}</math> in excess, a <i>white precipitate formed</i>; added <math>\text{Am}_2\text{S}</math>, the color of the precipitate remained unaltered.</p> <p>To another portion of the solution added <math>\text{KHO}</math>, drop by drop, until a precipitate formed, then more <math>\text{KHO}</math> and boiled; a white precipitate insoluble in <math>\text{KHO}</math>, and not darkening in the air:—<i>Presence of oxalate, phosphate, etc., of alkaline earths.</i></p> <p>Since the presence of an oxalate, of an alkali, or alkaline earth-metal was indicated in the preliminary examination, a portion of the original substance was heated on platinum foil, then dissolved in HCl, and to the solution excess of <math>\text{AmHO}</math> and of <math>\text{Am}_2\text{CO}_3</math> was added. The liquid was warmed and put aside till the precipitate subsided, then the liquid was decanted and the precipitate dissolved in a few drops of HCl.</p> <p>A platinum wire dipped into this HCl solution gave a <i>red</i> flame coloration, appearing <i>dusky green</i> through the indigo-prism:—<i>Presence of Ca.</i> [Confirmed by another portion of the HCl solution giving a white precipitate with excess of <math>\text{AmHO}</math> and <math>\text{Am}_2\text{C}_2\text{O}_4</math>.]</p>
--	--

The presence of  $\text{C}_2\text{O}_4$  was confirmed by heating some of the original substance with strong  $\text{H}_2\text{SO}_4$ : gases were evolved which rendered milky a drop of lime-water, and burnt with a blue flame, showing presence of  $\text{CO}_2$  and  $\text{CO}$ :—*Presence of  $\text{C}_2\text{O}_4$ .*

*Found Ca,  $\text{C}_2\text{O}_4$ .*

F. A yellow metallic-looking substance given for Analysis.

*Preliminary Examination :*

Experiment.	Observation.	Inference.
1. Heated a portion of the substance strongly in a small tube closed at one end.	A yellow sublimate formed, which melted into brown drops; a smell of burning S was evolved, and a piece of paper dipped into $K_2Cr_2O_7$ solution, when placed in the mouth of the tube, became green, . . . . A brown residue left, . .	Presence of S. Probable presence of Fe.
2. Heated strongly on charcoal in the inner blowpipe flame.	Strong smell of burning S. The residue fused into a dark-colored globule, which, when cold, was strongly attracted by the magnet,	Presence of S.  Probable presence of Fe.
3. Heated a fragment strongly in a glass tube, open at both ends, and held obliquely in the flame.	A gas was evolved with a suffocating smell, and which turned $K_2Cr_2O_7$ paper green, . . . .	$SO_2$ from combustion of S.

*Solution of substance.*

Warmed the powdered substance with a mixture of strong  $HNO_3$  and  $HCl$ , it dissolved completely; boiled down in an evaporating-dish, adding strong  $HCl$  several times; when nearly dry, diluted with water.

*Examination of the solution for the Metal.*

Since the substance dissolved entirely in  $HCl$ , Group I is absent; to a portion of the solution added  $H_2S$  water:

The solution became colorless, and white S was precipitated: <i>Probable presence of a ferric salt.</i>	To another part of the solution added $AmCl$ , then excess of $AmHO$ ; a brown flocculent precipitate formed, <i>probable presence of <math>Fe'''</math></i> ; then added $Am_2S$ , and heated; the precipitate became black, and on settling left a yellow solution:— <i>Absence of Ni.</i> Confirmed the presence of Fe by adding to a fresh part of the original solution $K_4FeCy_6$ solution, a dark-blue precipitate:— <i>Presence of <math>Fe'''</math>.</i>
--	--

Confirmed the presence of S by adding to a part of the original solution  $\text{BaCl}_2$ , a white precipitate formed which did not disappear on boiling:—*Presence of  $\text{SO}_4$  derived from oxidation of sulphur.*

*Found  $\text{Fe}_2\text{S}_3$ .*

*G. A white powder given for Analysis.*

Boiled a small portion with water, it did not dissolve.

Boiled another portion with  $\text{HCl}$ , it did not dissolve: added  $\text{HNO}_3$  and heated again, the substance was not dissolved.

Hence the substance is insoluble in water and in acids, and since it is white, it can only consist of one of the following substances:

$\text{BaSO}_4$ ,  $\text{SrSO}_4$ ,  $(\text{CaSO}_4)$ ,  $\text{PbSO}_4$ ,  $(\text{PbCl}_2)$ ,  $\text{AgCl}$ ,  $\text{SiO}_2$ ,  $\text{Al}_2\text{O}_3$ ,  $\text{SnO}_2$ ,  $\text{Sb}_2\text{O}_3$ ,  $\text{CaF}_2$ .

*Preliminary Examination:*

Experiment.	Observation.	Inference.
1. Took some of the powder up with a loop of moist platinum-wire, ignited it strongly in the inner blowpipe flame, moistened it with $\text{HCl}$ , and held in the Bunsen flame.	A yellowish-green coloration.	Presence of Ba.
2. Fused a portion with $\text{Na}_2\text{CO}_3$ on charcoal in the inner blowpipe flame, placed the cooled mass on a bright shilling, and moistened it with a drop of water.	A black stain on the coin.	Presence of S.

*Examination of the Solution.*

Fused some of the powder with fusion mixture on platinum foil: boiled the cold mass with water, filtered.

*Filtrate*, made acid with HCl and added  $\text{BaCl}_2$ , a white precipitate, which did not disappear on boiling the liquid: *Presence of  $\text{SO}_4$ .*

*The residue on the filter* was well washed with boiling water, and boiling dilute HCl poured upon it, it dissolved completely:

Absence of Group I.	To a portion added strong $\text{H}_2\text{S}$ -water and heated:		
	No precipitate. Absence of Group II.	To another portion added AmCl, then AmHO in excess, then $\text{Am}_2\text{S}$ , and boiled:	
		No precipitate/s. Absence of Group III.	To another portion of the original solution added AmCl, AmHO, and $\text{Am}_2\text{CO}_3$ :
			A white precipitate: Presence of Ba, Sr, or Ca. A platinum-wire loop dipped into the original solution, gave a yellowish-green tint to the Bunsen flame: <i>Presence of Ba.</i>

*Found Ba,  $\text{SO}_4$ .*



## SECTION VI.

## FULL ANALYTICAL COURSE AND TABLES.

AFTER having tried the reactions for the members of the analytical groups, and learnt for each group how to detect its members occurring singly or two or more together, solutions containing a single metal belonging to any of the groups, or members of two or more of these groups mixed, will readily be analyzed.

**370.** *It is necessary first to separate the metals present into groups*, by adding the group-reagents successively in such an order that each precipitates its own group only, leaving in solution the members of all other groups which may be present. This is effected by adding the group-reagents in the order directed in the General Table (419); refer to this and to the Group Table on p. 186. HCl, which is the first reagent added, precipitates only Group I, leaving Groups II, III, IV, and V in solution; hence, if a precipitate is obtained on adding excess of HCl, after filtering it off any member or members of Group I present will be obtained upon the filter, whilst the members of all other groups remain in the filtrate. On passing  $H_2S$  into the heated filtrate,\* members of Group II alone will be precipitated and may be filtered off; the filtrate,\* after removal of  $H_2S$  and oxidation of any Fe present by boiling with  $HNO_3$ , may be mixed with AmCl and excess of AmHO; the AmHO will precipitate any members of Group III A, Group III B, and Mg remaining dissolved by the AmCl; the addition of Am $_2$ S to the filtrate\* will separate Group III B as sulphides. Group IV is precipitated from

\* Or solution, supposing no precipitate to have been obtained, and filtration therefore to have been unnecessary.

the filtrate\* by the last group-reagent  $\text{Am}_2\text{CO}_3$ , and the filtrate\* is examined for Group V, the members of which not being precipitated by any group-reagent will now remain in solution if they were originally present.

Since each group must be *entirely* precipitated by its own group-reagent it is necessary, if any precipitate is formed, to add each group-reagent in "excess" before proceeding to add the next; unless the presence of the reagent in excess is ascertained before proceeding, much confusion may be caused.

**371. Addition of Reagents in Excess.**—The most general method for ascertaining the presence of a reagent in excess is to add a few drops more of the reagent to the clear liquid obtained either by letting the precipitate settle or by filtering off a small quantity of it; if any further precipitate is formed the filtered portion must be returned, more of the reagent must be added, and the clear liquid again tried with a few additional drops of the reagent, this process being repeated until no further precipitate is caused.

In certain cases this method of proceeding is rendered unnecessary, the excess of a reagent being detectable by its smell after thoroughly mixing up the solution and blowing out the air above it; this is the case for example with  $\text{H}_2\text{S}$  and  $\text{AmHO}$ . In other cases the excess is seen by the color of the filtrate, as with yellow  $\text{Am}_2\text{S}$  for example.

**372.** Since certain organic substances hinder or prevent the precipitation of the members of Group III A by  $\text{AmHO}$ , it is necessary, if the preliminary examination has shown the presence of organic matter, to destroy it by evaporation and ignition before proceeding to precipitate Groups III and IV; evaporation to dryness and gentle ignition are also necessary to separate  $\text{SiO}_2$ , which if remaining in solution might be mistaken for  $\text{Al}_2\text{Ho}_6$ . But if it is known that neither organic substances nor  $\text{SiO}_2$  are present, a considerable saving of

---

\* Or solution, supposing no precipitate to have been obtained, and filtration therefore to have been unnecessary.

time may be effected by boiling the  $\text{H}_2\text{S}$  filtrate until it no longer smells of  $\text{H}_2\text{S}$ , then continuing to boil for several minutes after adding a few drops of strong  $\text{HNO}_3$ ; this process may replace the evaporation to dryness and ignition.

**373.** The ordinary method of analysis drawn out in the general table and in the tables for examination of the group precipitates will present little difficulty if the directions therein given are strictly complied with. Table III c will perhaps be found a little more difficult, and should not be attempted until some proficiency has been obtained by practice on the other group tables. Table III d, which must be employed under conditions fully explained in (460), should not be used until considerable proficiency has been attained.

**374.** It will be seen that the full course of the analysis is arranged so as not only to prove what is present, but also that all else is absent; this, except in special cases, is the object in view.

The process of analysis is much hastened by using separate portions of the solution instead of carrying on the examination throughout with one part of the solution; the plan in the general table is however preferable, since it gives at once a notion of the relative quantities of the different substances present.

For an example of how to write out the results see pars. (488-493); it will be seen that the form adopted is that of the analytical tables.

---

#### PRELIMINARY EXAMINATION OF SOLID SUBSTANCES.

**375.** Before proceeding to employ the full course of analysis for substances of complex composition, the student will do well to examine a few solid substances for metals and acid radicles by the tables given in paragraphs 387-418.\* For the full account of how to try these tests for any particular substance the student must refer back to the preceding reactions in Section IV.

---

\* For examples of substances to be analyzed by the Preliminary Examinations, see paragraph 538.

The results of these analyses are entered in the form of the table as shown in the examples in paragraphs (488) and (489). *The entries must be made during the progress of analysis.*

**376. Preliminary Examination for Metals.**—A little study of this table (387 *et seq.*) will show that it contains three main tests marked Exps. I, II, and III; these stand in the first column: in the second column are placed the results which may be noticed on trying these experiments; the student should glance through these so as to be forewarned of what he has to look for. The main classification of the results which may be observed is indicated by large type letters, the subsidiary divisions are marked by small numbers, and the individual results are simply placed in succession in a vertical column. Occasionally an observation is made which it is advisable to confirm by an additional experiment: this “confirmatory” test is then entered in the first column, just below the observation to be confirmed, and is marked “confirm.” to distinguish it from the main experiment; its results are placed beside it in the second column: the student must understand that any one or more of the results noted in the second column may be looked for.

A few salts and mixtures such as those mentioned in (538) should be subjected to a careful preliminary examination by the student, and the results obtained carefully entered.

**377. Preliminary Examination for Acid Radicles.**—This table (410-418) exactly resembles in arrangement the table for metals which precedes it, and no explanation is therefore necessary. The student should examine by it some solid substances (538) for acid radicles only.

**378.** A few solid substances may now be tested for both metals and acid radicles by the two preliminary tables.

---

SIMPLE SUBSTANCES, DETECTION OF ACID RADICLES  
AND METALS BY PRELIMINARY TESTS AND BY  
EXAMINATION OF THE SOLUTION.

**379.** After having become familiar with the preliminary examination of solid substances the student may

analyze some simple substances containing only one or two metals and acid radicles. The analytical course drawn out for the examination of complex substances may be at once employed for these more simple analyses, as the student is thus enabled to prove not only the presence of any particular metals or acid radicles, but also the absence of all other commonly occurring metals and acid radicles.

He will by this means be gradually introduced to the analytical course, and after satisfactorily analyzing some simple substances, may proceed to mixtures of gradually increasing difficulty or complexity.

At first no mixture should be attempted which will yield a precipitate containing a phosphate in Group III, and members of Group III A and III B should not be present together in the same solution, also substances containing organic matter and  $\text{SiO}_2$  should be avoided; the examination of the filtrate after passing  $\text{H}_2\text{S}$  is thus very much simplified.

The following is a description of the course to be pursued in making the complete analysis of a complex substance.

### GENERAL COURSE OF ANALYSIS FOR SOLID AND LIQUID SUBSTANCES.

THE substance given for analysis\* will be either a liquid or a solid, since the analysis of gases forms a department which is best studied by itself.

Unless it is known that cyanogen is not present in the substance, it must be tested for in a small portion by the "Prussian blue" or AmCyS tests (304 or 306); if found, refer to (482 *et seq.*) for the preparation of the solution and the method of analysis: if cyanogen is not present, the substance is examined by (380, 381), or by (382 *et seq.*), according as it is liquid or solid.

Solid substances require to be dissolved before they can be fully analyzed;  $H_2O$ ,  $HCl$ ,  $HNO_3$  and aqua regia are the solvents usually employed.

#### A. THE SUBSTANCE IS A LIQUID.

380. 1.—Test it with both blue and red litmus-papers, one of the following results will be obtained:

<i>It is neutral</i> (not changing either paper).	<i>It is acid</i> (turning <i>blue</i> litmus <i>red</i> ).	<i>It is alkaline</i> (turning <i>red</i> litmus <i>blue</i> ).
Absence of acids and alkalis and of salts with acid or alkaline reaction. The salts of Ag and Mg and certain salts of Am, Na, K, Ba, Sr, and Ca are the only neutral soluble salts.	Presence of an acid, or salt with acid reaction.	Presence of a hydrate of Am, Na, K, Ba, Sr, or Ca, or of a salt with alkaline reaction.

381. 2.—Evaporate a few drops of the liquid upon a piece of platinum foil or a piece of thin glass,† smelling it occasionally:

\* For examples of substances to be analyzed, see par. 539.

† A piece of the side of a broken flask will serve well.

**No residue is left.**

The liquid must consist of some volatile substance, probably of water, either pure or containing certain gases or volatile substances dissolved in it (e. g.  $\text{NH}_3$ ,  $\text{HCl}$ ,  $\text{Br}$ , etc.), which are detected by their smell, by the action of the liquid on litmus-paper, or by special tests. If the liquid has no action on litmus, no smell, and leaves no residue on evaporation, and is tasteless, it is pure water.

**A residue is left:**—Examine the liquid by the further preliminary tests in (334) or evaporate a portion of the solution to dryness in a porcelain dish, avoiding heating the substance after it is dry; and examine it by the preliminary tables for metals and acid radicles (387–418).

Examine the larger portion of the remaining solution for metals by the general table which follows (419), and the remainder of the solution for acid radicles by (440 *et seq.*).

**B. THE SUBSTANCE IS A SOLID.****382. B. a. IT IS NOT A METAL.**

Finely powder it in a mortar, using an agate mortar if the substance is very hard. A part of this powder is examined by the preliminary tables for metals and acid radicles (387–418).

**383.** A large part of the remainder\* is boiled in a flask or boiling-tube with distilled water, if it dissolves, see (384): if the powder does not dissolve, allow the liquid to stand still until the undissolved portion has settled, then decant through a filter:

**Filtrate:**—It is usually possible to tell whether anything has been dissolved by the water by observing whether it has acquired any color or action upon litmus-paper. If there is any doubt, try whether a residue remains when a few drops are evaporated; also, whether  $\text{Na}_2\text{CO}_3$  causes any precipitate. If none of these tests proves the presence of dissolved matter, the water solution may be neglected.

If anything is in solution, it is kept and marked.

Solution I.

**Residue in flask:**—Boil with dilute  $\text{HCl}$  (385), allow any residue to settle, decant through the filter used above, and boil the residue with strong  $\text{HCl}$ , let settle, decant through the filter:

**Filtrate.**

**Residue:**—Heat with dilute, then with strong  $\text{HNO}_3$ , as was directed above for  $\text{HCl}$ ; decant:

**Filtrate.**

The residue may be warmed for a short time with a mixture of strong  $\text{HNO}_3$  with three times as much  $\text{HCl}$ ; if this does not dissolve it, decant the acid mixture.

**Residue:** Wash well with water, dry, and examine as directed (475 *et seq.*) for substances insoluble in water and acids.

Solution II.

Solution III.

Solution IV.

\* If the whole of the substance left after completing the Preliminary Examination has to be employed in making the solution,

**384.** If at any of the preceding stages the substance dissolves entirely, proceed at once to examine the solution by the General Table (419). The Solutions I, II, III, IV, need rarely be examined separately; a general method is to add to I some  $\text{HNO}_3$  (see 420), and then some  $\text{HCl}$  to the clear liquid; add also to the  $\text{HNO}_3$  solution (III) some  $\text{HCl}$ ; any precipitate thus caused by  $\text{HCl}$  is examined by Table I, and any precipitate caused then on mixing the solutions I, II, III, IV, is examined by (475) as an insoluble substance. It is best to boil down the solutions III and IV considerably before mixing them with I and II, in order to get rid of the  $\text{HNO}_3$  as much as possible, then dilute with dilute  $\text{HCl}$ ; any precipitate caused by  $\text{HCl}$  being examined by Table I. On cooling the hot  $\text{HCl}$  solution crystals often separate; these consist probably of  $\text{PbCl}_2$ ; they may be filtered off and dissolved in a little boiling water; if on addition of  $\text{K}_2\text{CrO}_4$  to this solution of the crystals a yellow precipitate forms, it shows the *presence of Pb*.

*Note* —If a complex solid mixture is given for analysis, and the only object is to obtain a solution as rapidly as possible, it may be at once boiled with aqua regia for a few minutes, then filtered: the residue is examined as a substance insoluble in water and acids (475 *et seq.*), and the filtrate is evaporated nearly to dryness: dilute  $\text{HCl}$  is added, and the solution examined by the General Table (419).

$\text{HNO}_3$  seldom requires to be used in dissolving a substance, and when necessary should be employed only in small quantity, since it is liable to cause a precipitate of S from  $\text{H}_2\text{S}$  (423).

**385.** Any changes which occur whilst preparing the solution should be noted down; notice more particularly whether any gases are given off on heating with  $\text{HCl}$ ; many of these gases may be detected by their smell, but a closer examination of them may be neglected, as they are obtained in the preliminary examination for acid

---

the method given below must be employed; if, however, plenty of the substance is in hand, the method given under (332) may be used, the process of solution will thus be much more rapidly executed.



radicles (411). If the gases come off from the aqueous solution on adding HCl, or on mixing the HCl and aqueous solutions, they must have been present combined with metals as soluble salts, probably salts of alkalies; if they are evolved on treating the residue insoluble in water with HCl, they were present as insoluble salts.

---

**386. B. b. THE SUBSTANCE IS JUDGED FROM ITS METALLIC LUSTRE TO BE A METAL OR ALLOY.**

Examine it according to the directions (467).

---

#### PRELIMINARY EXAMINATION FOR METALS.

**387.** The substance is required in the solid state for this examination; hence if a solution is being analyzed, some of it must be evaporated to dryness and the dry residue employed (see 381).

Carefully note down the appearance and physical properties of the substance: whether it is crystalline or amorphous;\* its hardness,† color, smell, and anything which strikes you on a careful examination aided by a pocket lens.

The following three experiments, together with some confirmatory ones (marked "Confr.") when necessary, are then to be tried on small quantities of the finely powdered substance, and the results obtained noted down.

For a further explanation of the form of the table, see paragraph 376.

---

\* A body, if crystalline, consists of particles which have a definite and similar shape: the particles are amorphous if they show no similar and definite shape.

† The hardness of a substance may be roughly tested by trying to scratch it with the point of a penknife: if very easily scratched, it is soft; if the knife will not scratch it, but simply marks it as a lead-pencil would do paper, it is very hard (*e. g.*, quartz).

Experiment.	Observation.	Inference.																			
388. Exp. I.—Heat a small quantity of the substance in a small test-tube or piece of hard glass tubing drawn off at one end.	a. The substance does not change.	Absence of organic substances which blacken and give off a smell of burning; of volatile substances which sublime; of substances which are altered by ignition; and of water combined with or absorbed by the substance, which would be evolved as steam, and form drops on the side of the tube.																			
	<p>b. The substance changes.</p> <p>1. <i>It changes color:</i></p> <table><tr><td><i>Yellow,</i></td><td><i>hot</i></td><td><i>white,</i></td><td><i>cold.</i></td></tr><tr><td><i>Yellow, brown,</i></td><td>"</td><td><i>yellow,</i></td><td>"</td></tr><tr><td><i>Yellow-brown,</i></td><td>"</td><td><i>yellow,</i></td><td>"</td></tr><tr><td><i>Dark red or</i></td><td><i>"</i></td><td><i>red-brown,</i></td><td><i>"</i></td></tr><tr><td><i>brown.</i></td><td><i>}</i></td><td></td><td></td></tr></table> <p>The substance blackens—</p> <p>a. Emitting a smell of burning. The black carbon may be burnt off by strongly heating on platinum foil with the blowpipe flame.</p> <p>Smell of acetone. Smell of burning sugar.</p> <p>b. Emitting no smell of burning. The color is not removed by ignition as under a.</p> <p>2. <i>It fuses, and becomes solid again on cooling.</i></p> <p>3. <i>It sublimes.</i></p>	<i>Yellow,</i>	<i>hot</i>	<i>white,</i>	<i>cold.</i>	<i>Yellow, brown,</i>	"	<i>yellow,</i>	"	<i>Yellow-brown,</i>	"	<i>yellow,</i>	"	<i>Dark red or</i>	<i>"</i>	<i>red-brown,</i>	<i>"</i>	<i>brown.</i>	<i>}</i>		
<i>Yellow,</i>	<i>hot</i>	<i>white,</i>	<i>cold.</i>																		
<i>Yellow, brown,</i>	"	<i>yellow,</i>	"																		
<i>Yellow-brown,</i>	"	<i>yellow,</i>	"																		
<i>Dark red or</i>	<i>"</i>	<i>red-brown,</i>	<i>"</i>																		
<i>brown.</i>	<i>}</i>																				
<p><i>Confirmatory.</i>—The sublimate is carefully examined, if necessary, with the aid of a lens.</p> <p>389. <i>Confirmatory.</i>—If a sublimate has been obtained, but not otherwise, heat another portion of the substance, mixed with about three times as much Na<sub>2</sub>CO<sub>3</sub> in fine dry powder and a little KCy, in a tube closed at one end, in the manner described under par. 147.</p>	<p>White, crystalline (sparkling) sublimate. White, non-crystalline sublimate. Black sublimate, becomes red when rubbed.</p> <p>Yellow sublimate.</p> <p>Gray mirror, which, when rubbed with a splinter of wood or a glass rod, or when examined by a lens, is seen to consist of globules of Hg. Blackish-brown shining mirror, and smell of garlic; no globules.</p>																				

Experiment.	Observation.	Inference.
390. <i>Confirmatory.</i> —Mix also some of the substance with soda-lime in a mortar and moisten; or pour a little strong KHO solution upon it, and heat to boiling.	NH <sub>3</sub> is given off, known by its smell, and by turning moist red litmus-paper blue or turmeric-paper brown.  <i>Note.</i> —The change in color is only a trustworthy indication if it occurs very soon, as the NH <sub>3</sub> in laboratory air often causes it to occur after a time.  Sublimate of reddish drops, } which are yellow when cold. } Violet vapor, cooling to blackish crystals. 4. <i>The substance gives off water.</i> It fuses first, then gives off water, and again becomes solid if the heat is continued. It swells up considerably whilst giving off its water.	Presence of NH <sub>3</sub> .  S, free or from certain polysulphides. Presence of I.  Presence of water, absorbed or combined. Water of crystallization.  Borax and certain borates, alums, and phosphates. Probably NH <sub>4</sub> compounds.
391. <i>Confirmatory.</i> —Examine the drops of water on the sides of the tube with blue and red litmus-papers.	The water is alkaline.  The water is acid.  5. <i>Gas or vapor is given off.*</i> <i>a. It is without smell.</i>	Presence of volatile acids, <i>e. g.</i> , HNO <sub>3</sub> , HCl, H <sub>2</sub> SO <sub>4</sub> .
392. <i>Confirmatory.</i> —Introduce a burningsplinter of wood into the tube.	It burns more brightly, and if introduced with a spark at the end is inflamed. The flame is extinguished.	O, from chlorates, nitrates, peroxides, etc. CO <sub>2</sub> or N.
393. <i>Confirmatory.</i> —Introduce a glass rod, drawn out if necessary, and moistened with lime-water, into the tube.	The lime-water turns milky.  The lime-water does not turn milky.  <i>b. The gas or vapor has a smell.</i>	CO <sub>2</sub> from carbonates or oxalates. N (from AmNO <sub>3</sub> probably).
394. <i>Confirmatory.</i> —Introduce a glass rod or slip of paper moistened with K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> .	Smell of burning S. The K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> turns green; moist blue litmus is also reddened by the gas.  Red nitrous fumes are given off, known by their peculiar smell, and by not coloring starch paste orange red.  Smell resembling Cl:	SO <sub>2</sub> from combustion of free S, or from acid sulphites, hyposulphites, reduction of sulphates or oxidation of sulphides. Nitrates of heavy metals ( <i>e. g.</i> , of Pb, Bi, Hg, etc.).
395. <i>Confirmatory.</i> —Introduce a glass rod carrying a piece of moist litmus.	Yellowish gas, which bleaches moist litmus.	Cl from certain chlorides.

\* This will usually be shown by effervescence of the substance, or the gas or vapor is detected by the color or smell; but the gas or vapor is often only found on testing for it specially by (392) and (393).

Experiment.	Observation.	Inference.
396. <i>Confirmatory.</i> — Introduce a glass rod with moist starch powder on its end, obtained by wetting a glass rod and dipping it into starch powder.	Brown vapor, which colors the moist starch powder orange red. Violet vapor, which colors starch powder (or better, starch solution) blue. <i>c. The gas burns at the mouth of the tube with a:—</i> Fale grayish-green flame.  Bright white flame.  <i>Purple flame.</i>	Br from certain bromides. I, free or from certain iodides.  NH <sub>3</sub> probably from strongly heated NH <sub>4</sub> NO <sub>3</sub> . PH <sub>3</sub> probably from a hypophosphite; the unburnt gas smells of garlic. Cy, from a cyanide, probably HgCy <sub>2</sub> .
397. Exp. II. — Dip a moistened loop of platinum-wire, which gives no color to the Bunsen flame, into the substance, and hold the loop with the adhering powder near the top of the Bunsen flame or in the inner blowpipe flame. Then moisten with a drop of strong HCl and heat again. Continue the heating for some time.  <i>Note.</i> —If sillicic acid is present, the colorations for K and Na usually show only after ignition of the powdered substance with powdered CaSO <sub>4</sub> .	1. The substance colors the flame: <i>Intense yellow.</i>    <i>Pale violet, reddish-violet through cobalt glass or indigo-prism.</i>  <i>Yellowish-green.</i>  <i>Crimson, deep-red through cobalt glass or indigo-prism.</i> <i>Orange-red, dingy green through indigo-prism.</i> <i>Green.</i>   <i>Blue.</i>  2. The substance does not color the flame.	Na. If a violet or red color is seen on examining the flame through cobalt glass or the indigo-prism, either K or Sr, or both, are also present.  K.  Ba. Certain other substances also give green colorations. Sr. Ca. Cu, B <sub>2</sub> O <sub>3</sub> . Cu shows green outside and blue in the centre, after being moistened with HCl. As, Sb, Pb, CuCl <sub>2</sub> . CuBr <sub>2</sub> . Probable absence of the above substances.
398. Exp. III. — Heat the substance in a small cavity scooped in a piece of wood-charcoal, in the blowpipe flame.	A. The substance decrepitates (crackles). B. The substance deflagrates (causes charcoal to burn rapidly). C. The substance fuses easily, and is absorbed by the charcoal, or forms a liquid bead. D. An infusible residue is left on the charcoal. 1. <i>The residue is white and very luminous.</i>	NaCl and certain other crystalline salts. Chlorates, nitrates, etc. Salts of alkalis and certain salts of the alkaline earths.
399. <i>Confirmatory.</i> — Detach a portion of the cool residue from the charcoal, place it upon a piece of red litmus-paper, and moisten with a drop of water.	The paper turns blue, showing the residue to be alkaline.	Probably BaO, SrO, CaO, MgO, Al <sub>2</sub> O <sub>3</sub> , ZnO (or SiO <sub>2</sub> ). BaO, SrO, CaO (MgO).

Experiment.	Observation.	Inference.
400. <i>Confirmatory</i> .—Moisten the residue on the charcoal when cool, with several drops of $\text{Co}(\text{NO}_3)_2$ solution, and heat again strongly in the outer blowpipe flame.	A <i>blue</i> residue, the color of which does not disappear with intense heat. A <i>pink</i> residue. A <i>green</i> residue.	$\text{Al}_2\text{O}_3$ (and some phosphates, silicates, and borates). $\text{MgO}$ . $\text{ZnO}$ .
401. <i>Confirmatory</i> .—Heat a small quantity of the substance in a clear colorless borax bead, first in the outer, then in the inner blowpipe flame.	2. <i>The residue is colored.</i> Color of bead. In outer flame. In inner flame. <i>Green</i> , hot. } <i>Red or colorless</i> , cold. } <i>Blue</i> , cold. } <i>Blue</i> , hot and cold. } <i>Blue</i> , hot and cold. } <i>Violet</i> , hot. } <i>Gray or opaque</i> . } <i>Yellow</i> , cold. } <i>Brown-red</i> , hot. } <i>Green</i> , hot and cold. } <i>Yellow or colorless</i> , cold. } <i>Green</i> , hot and cold. } <i>Green</i> , hot and cold. } <i>Reddish-purple</i> , hot and cold. } <i>Colorless</i> , hot and cold. }	Cu. Co. Ni. Fe. Cr. } Mn. } See (402).
402. <i>Confirmatory</i> .—The presence of Mn and Cr, if suspected, may be confirmed by fusing the substance with some powdered $\text{Na}_2\text{CO}_3$ and $\text{KNO}_3$ on platinum foil.	<i>Light-yellow</i> mass on cooling. <i>Blue-green</i> mass on cooling.	Cr. Mn. (See 406.)
403. <i>Confirmatory</i> .—Heat also some of the substance, mixed or covered with powdered $\text{K}_2\text{CO}_3$ and $\text{Na}_2\text{CO}_3$ (or $\text{K}_2\text{S}_2\text{O}_8$ ), on charcoal in the inner blowpipe flame.	3. <i>The residue is colored, or metallic scales or globules are seen.</i> The substance is reduced to the metallic state. (See 407.)  a. Without incrustation. [Brilliant <i>white</i> metal. <i>Yellow</i> metal. <i>Red</i> scales or globules. <i>Gray</i> powder, attracted by a magnetized knife-blade. (See 407.)  b. With incrustation:  Globules. Incrustation. <i>White</i> , malleable. None, or very slight.  " <i>brittle</i> . <i>Orange</i> , hot; } <i>yellow</i> , cold. } " <i>malleable</i> . <i>Yellow</i> , hot and cold. }	Au, Ag, Cu, Pb, Sn, Sh, Bi; [Co, Ni, Fe, Mn form only gray powders]. Au, Ag, Cu, Co, Ni, Fe, Mn. Ag. Au. Cu. Fe, Co, Ni, Mn.
404. <i>Confirmatory</i> .—Detach several of the globules with the point of a knife and strike them on the bottom of an inverted mortar with the pestle: if they flatten to a cake they are <i>malleable</i> , if crushed to a powder they are <i>brittle</i> .  <i>Note</i> .—The appearance and malleability of one reduced metal is frequently much altered by the presence of another metal.	" <i>brittle</i> . " <i>malleable</i> .  " <i>brittle</i> ; giving white fumes.	Sn: differs from Pb in being difficult to fuse into globules, and in not marking paper. Bi. Pb. The globules of Pb differ from those of Sn by marking paper as black lead does, if taken on the point of a knife. Sb.

Experiment.	Observation.		Inference.
	Globules. <i>White</i> or none.	Incrustation. <i>Yellow</i> , hot; <i>white</i> , cold.	} Zn.
	" "	<i>Red-brown</i> , easily volatili- zied when heated in the flame.	
	None.	<i>White</i> , and white fumes; smell of garlic.	} Cd.
			} As.
405. Exp. IV.—Fuse some of the substance, mixed with $\text{Na}_2\text{CO}_3$ , on charcoal in the inner blowpipe flame produced from a spirit-lamp; remove the fused mass when cold, place it on a bright silver coin, moisten with a drop of water, crush the mass, and let stand for several minutes.	On rinsing off the substance, the coin is found to be stained black (408). Also, if a drop of $\text{HCl}$ be placed upon the mass, a smell of $\text{H}_2\text{S}$ is perceived, and lead-paper is blackened.		Presence of S, free or combined.

*Note.*—Since this reaction serves to detect S and sulphur acid radicles, it more properly belongs to the Preliminary Examination for acid radicles.

#### NOTES TO THE PRECEDING PRELIMINARY TABLE.

**406.** The green color produced by Mn conceals the light yellow of Cr, but on boiling the residue with water it dissolves, giving a pink or purple solution, the color of which is best seen on filtering; this color is due to the presence of  $\text{KMnO}_4$ , and shows Mn to be present. On acidifying this solution with  $\text{H}^+\text{A}$  and boiling again for several minutes, the purple color is destroyed; and on filtering, the yellow color due to the Cr is seen: the presence of Cr may be further confirmed by the formation of a yellow precipitate in the acid solution on addition of  $\text{PbA}_2$ .

**407.** The metal is best separated and examined by detaching the mass when cold from the charcoal, and powdering it by crushing in an agate mortar or on a watch-glass with a little water, letting stand for a short time, and then quickly pouring off the water down a glass rod or pestle which is wetted and pressed against its edge; by several times repeating this operation, the heavier metallic particles alone are left in the mortar or watch-glass.

**408.** This stain is removed by rubbing the coin with a little lime made into a paste with water.

**409. Note.**—When the substance to be examined contains several bodies, they frequently more or less mask one another's reactions: thus Co if mixed with Fe will give a bead green whilst hot and blue when cold, thus resembling Cu, but differing in remaining blue in the inner flame: hence the composition of many complex mixtures is only roughly indicated by the preliminary examina-

tion, and must be confirmed and established in the wet way. Many substances, more particularly minerals, however, can be completely analyzed by a careful preliminary examination.

### PRELIMINARY EXAMINATION FOR ACID RADICLES.

**410.** The substance is required for this examination in the state of powder or of *strong* solution.

If the substance given for analysis is a solution, part of it may be evaporated to dryness: the dry residue is then finely powdered and subjected to the following tests. If several acid radicles and metals are present, the reactions may be more or less perfectly concealed or altered, hence failure in obtaining a certain reaction does not necessarily prove the absence of an acid radicle.

Experiment.	Observation.	Inference.
<b>411. EXP. I.</b> —Treat some of the substance with dilute HCl and note the result, then heat (416).	<i>One or more of the following gases may be evolved:</i>	
	A colorless gas without smell, which turns milky a drop of lime-water on the end of a glass rod.	CO <sub>2</sub> from a carbonate.
	A gas of suffocating smell, which turns a drop of K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub> solution green. Yellow S is precipitated at the same time.	SO <sub>2</sub> from a sulphite. SO <sub>2</sub> and S from a theio-sulphate.*
	A gas with fetid odor which blackens a drop of PbAc <sub>2</sub> solution, or a piece of lead-paper. (See 417.)	H <sub>2</sub> S from a sulphide; or possibly from a sulphite or hyposulphite, if Zn or other reducing agent be present.
	Reddish fumes.	N <sub>2</sub> O <sub>3</sub> from a nitrite.
	A yellowish green gas, of suffocating smell, which bleaches moist litmus-paper, is evolved:	
	a. In the cold; and also evolved on adding HAc.	Cl from hypochlorites.
	b. Only when heated.	Cl from action of HCl on oxidizing substances, e.g., MnO <sub>2</sub> , chromates, nitrates, chlorates, &c.
	A colorless gas, smelling of bitter almonds.	HCy from a cyanide.

\* Formerly called a "hyposulphite."

Experiment.	Observation.	Inference.
412. Exp. II.—Heat another portion gently with strong $H_2SO_4$ .	Any gas or gases evolved in Exp. I will probably be evolved also by strong $H_2SO_4$ ; no further attention need be paid to them; but in addition to these one or more acid gases may be given off. A gas is evolved with pungent smell, which fumes in the air, and renders milky a drop of $AgNO_3$ solution made acid with $HNO_3$ . 1. The liquid does not become brown or violet or give off colored vapor when heated. 2. Brown vapors are given off on heating, which color a little moist starch powder, orange-red. 3. Violet vapors are given off, when the mixture is heated, which color a drop of starch-paste blue. 4. A heavy fuming suffocating gas is evolved, and the liquid behaves as if the glass were greasy; on rinsing out the tube, and drying it thoroughly, the inside is seen to be corroded and dimmed. 5. The acid fumes are reddish, usually seen only on heating after adding a few fragments of Cu.	$HCl$ , $HBr$ , $HI$ , $(HF)^*$ from chloride, bromide, iodide (fluoride).  Probably $HCl$ from a chloride, confirmed by heating with $MnO_2$ and $H_2SO_4$ , when $Cl$ is given off (418).  $Br$ from a bromide.†  $I$ from an iodide.‡ $HF$ from a fluoride. <i>Note.</i> —The $HF$ acting upon the silica of the glass evolves $SiF_4$ , which is detected by holding a moistened glass rod in the gas, $SiO_2$ is deposited as a gelatinous film upon it.  Presence of a nitrate. <i>Note.</i> —If iodine is present it must be separated by $Cu_2SO_4$ (266) before a nitrate can be detected.
413. <i>Confirmatory</i> .—Drop into the hot liquid a few small pieces of copper.		
414. <i>Confirmatory</i> .—Boil some of the substance with water (or take some of the solution if a liquid), add about twice as much strong $H_2SO_4$ , cool, and pour solution of $FeSO_4$ carefully in upon the top of the acid liquid.	A dark brown ring or layer forms upon the surface of acid, best obtained on cooling the liquid by immersing the test-tube in cold water.  6. Greenish - yellow gas, smelling like $Cl$ , and exploding when warmed. Confirm by adding indigo solution and $H_2SO_4$ to the colorless aqueous solution of the substance (252).  For detection of a nitrate and chlorate when mixed, see (256).	Presence of a nitrate.        Presence of a chlorate.

\*  $HF$  would also render the drop milky by depositing  $SiO_2$  from the  $SiF_4$ , formed by its action on the glass, but it is distinguished by corroding the glass. (See 412, 4, note.)

†  $Br$  and  $I$  may be more perfectly liberated by mixing the substance with a little  $MnO_2$ , before adding  $H_2SO_4$ , and heating.



248 EXAMINATION FOR ACID RADICLES. [415-418.

Experiment.	Observation.	Inference.
<b>415. Confirmatory.</b> — Add a little alcohol to the substance and heat with $H_2SO_4$ .	7. The substance changes from yellow to green, O being evolved, as is shown by a glowing taper.	Presence of a chromate.
	8. Yellowish-green gas, with suffocating smell, which bleaches moist litmus.	Cl from a hypochlorite (detected already in 411), or from a chloride in presence of $MnO_2$ , etc.
	9. The same gas as in 8, but colored reddish-brown by nitrous fumes:	A chloride in presence of a nitrate or nitrite.
	9a. Or colored by $CrOCl_2$ .	A chloride in presence of a chromate.
	10. Substance does not blacken, but evolves CO (burning with a blue flame) and $CO_2$ (turning a drop of lime-water milky).	Presence of an oxalate.
	11. Substance blackens, and $SO_2$ is smelt.	Presence of a tartrate.
	12. A smell of acetic acid (vinegar) is noticed.	$\overline{HA}$ from an acetate.
	Strong fragrant odor.	Ethyl-acetate from an acetate.

NOTES ON THE PRECEDING TABLE.

**416.** If HCl has been employed in dissolving the substance, the acid radicles detected by the action of HCl will have been already noticed in preparing the solution for metals (383).

**417.** The S present in sulphides which are not decomposed by dilute HCl, is detected by fusing the powdered substance with two or three times as much fusion mixture in a covered crucible, boiling with a little water, and placing a drop upon a piece of bright silver (e. g., a silver coin), when a dark stain will be produced; or by adding excess of an acid to the solution, when  $H_2S$  will be evolved. Smaller quantities of S may be detected by the formation of a black color or precipitate on adding  $PbA_2$  to the solution.

**418.** Chlorides treated in this manner evolve Cl, recognized by its color, smell, and bleaching action on litmus-paper; but this test for chlorides is reliable only if the precautions stated in (259) are attended to, since many samples of  $MnO_2$  evolve Cl when warmed alone with  $H_2SO_4$ .

### ACTUAL EXAMINATION FOR METALS.

After the preceding Preliminary Examinations have been made, the substance is further examined in a more systematic manner in solution. This is often called "The Examination in the Wet Way."

---

#### 419. GENERAL TABLE FOR SEPARATION OF METALS INTO GROUPS BY GROUP-REAGENTS.

*Note.*—If the substance for analysis be a liquid with *alkaline* reaction, it requires special treatment (see 420); if its reaction be *neutral* or *acid*, it may be at once examined as directed below.

In this table (p. 250), and also in the group tables which follow, it has been supposed that all metals are present: if a metal or group of metals is absent, no precipitate will be produced when the reagent is added for its detection; and hence no filtration is necessary, the solution being then treated as the filtrate.

<p>Add to the greater part of the "original solution" <math>\text{HCl}</math>; if any precipitate is produced, continue to add <math>\text{HCl}</math> as long as it causes any further precipitate (421), warm and filter: The precipitate may contain—</p> <p><math>\text{PbCl}_2</math>—white.  <math>\text{AgCl}</math>—white.  <math>\text{Hg}_2\text{Cl}_2</math>—white.          (See 422.)</p> <p>Examine the precipitate by Table I (433).</p>	<p>The filtrate or solution, which must give no further precipitate on addition of several drops of <math>\text{HCl}</math> (423), is diluted with water (425), unless it is already dilute; then saturated with <math>\text{H}_2\text{S}</math> by passing the gas through the solution† (500 a), and warmed gently (426 and 426 a) and filtered.</p> <p>The precipitate may contain—</p> <p><math>\text{HgS}</math>—black,  <math>\text{PbS}</math>—black,  <math>\text{Bi}_2\text{S}_3</math>—black,  <math>\text{CuS}</math>—black,  <math>\text{CdS}</math>—yellow,</p> <p>Insoluble in <math>\text{KHO}</math> and in <math>\text{Am}_2\text{S}</math>.</p>	<p>Boil the solution or the filtrate (which has been diluted and had <math>\text{H}_2\text{S}</math> passed again into it, to make sure that Group II is completely precipitated), in a porcelain dish until it ceases to smell of <math>\text{H}_2\text{S}</math> (427), then add a little strong <math>\text{HNO}_3</math> and boil for several minutes, concentrating if necessary.† [If the presence of <math>\text{SiO}_2</math> is suspected, or if organic substances are likely to be present (the presence of organic substances will usually have been indicated by the body blackening in Exp. 388) the solution must here be evaporated to complete dryness, and the residue gently ignited in the dish, then dissolved (after cooling) by warming with a little strong <math>\text{HCl}</math> and adding water; <math>\text{SiO}_2</math>, if present, will remain undissolved (428).]</p> <p>If the presence of a phosphate is suspected, add a few drops of the acid solution to some <math>\text{AmH}_2\text{MoO}_6</math> solution in a test-tube, and warm gently; a yellow precipitate shows the presence of a phosphate, and the rest of the solution must be examined by Table III D (466); if no precipitate is formed with <math>\text{AmH}_2\text{MoO}_6</math> phosphates are absent.</p> <p>Add to the rest of the <math>\text{HCl}</math> solution (which has been freed from <math>\text{H}_2\text{S}</math> as directed above) <math>\text{AmCl}</math>, heat it to boiling, then add <math>\text{AmHO}</math> in excess, boil again and filter off at once any precipitate which may form as quickly as possible, keeping the funnel closely covered with a glass plate (429):</p>	<p>The precipitate may contain—</p> <p>Add to the filtrate or solution, which may be colored (430), <math>\text{Am}_2\text{S}</math> in excess, boil and filter:</p>	<p>The precipitate may contain—</p> <p><math>\text{ZnS}</math>—white.  <math>\text{MnS}</math>—light-pink.  <math>\text{CoS}</math>—black.  <math>\text{NiS}</math>—black.</p> <p>Examine this precipitate at once by Table III B (437).  <i>Note.</i>—If the filtrate is brown, <math>\text{Ni}</math> is often easily detected in it. (See 431.)</p>	<p>Add to the filtrate or solution, which must be distinctly yellow or brown (431), <math>\text{Am}_2\text{CO}_3</math>.          The filtrate may contain <math>\text{Mg}</math>, <math>\text{K}</math>, <math>\text{Na}</math>.</p> <p>Examine by Table V (439).</p>
<p><i>Note.</i>—The absence of <math>\text{Pb}</math> is not proved, if it is not precipitated in this group as <math>\text{PbCl}_2</math> is somewhat soluble, and <math>\text{Pb}</math> may therefore be found only in Group II.</p>	<p><i>Note.</i>—1. This precipitate should not be examined at once as possibly a precipitate may be formed on boiling the filtrate, and this precipitate would have to be added to the above.          2. Unless the liquid has been sufficiently diluted some of the above metals may be afterwards precipitated by <math>\text{Am}_2\text{S}</math> in Group III, and cause much confusion.</p>	<p>Examine the precipitate by Table II (435).</p> <p><i>Note.</i>—1. This precipitate should not be examined at once as possibly a precipitate may be formed on boiling the filtrate, and this precipitate would have to be added to the above.          2. Unless the liquid has been sufficiently diluted some of the above metals may be afterwards precipitated by <math>\text{Am}_2\text{S}</math> in Group III, and cause much confusion.</p>	<p>The precipitate may contain—</p> <p><math>\text{Al}_2\text{H}_6</math>—almost colorless.  <math>\text{Cr}_2\text{H}_6</math>—light-green.  <math>\text{Fe}_2\text{H}_6</math>—reddish-brown.</p> <p>Examine the precipitate by Table III A (436).</p>	<p>The precipitate may contain—</p> <p><math>\text{ZnS}</math>—white.  <math>\text{MnS}</math>—light-pink.  <math>\text{CoS}</math>—black.  <math>\text{NiS}</math>—black.</p> <p>Examine this precipitate at once by Table III B (437).  <i>Note.</i>—If the filtrate is brown, <math>\text{Ni}</math> is often easily detected in it. (See 431.)</p>	<p>Add to the filtrate or solution, which must be distinctly yellow or brown (431), <math>\text{Am}_2\text{CO}_3</math>.          The filtrate may contain <math>\text{Mg}</math>, <math>\text{K}</math>, <math>\text{Na}</math>.</p> <p>Examine by Table V (439).</p>

\* By the "original solution" is meant the solution of the substance (if a solid), or the substance itself (if a liquid), to which no reagent has been added.

† Instead of diluting and passing the gas,  $\text{H}_2\text{S}$  solution may be added.

‡ If silicates, phosphates, and organic substances are known to be absent, the following portion, inclosed in square brackets, may be omitted.

## NOTES TO THE GENERAL TABLE.

**420.** If the liquid under examination is alkaline in reaction, before commencing to examine it by the General Table  $\text{HNO}_3$  must be added in excess; if no precipitate forms, proceed at once to examine the liquid by the general table; if a precipitate is produced, more  $\text{HNO}_3$  is to be added, and the liquid heated; should a precipitate still remain, it is filtered off and the filtrate examined by the general table, commencing with the addition of  $\text{HCl}$ .

The precipitate caused and not dissolved by  $\text{HNO}_3$  may be finely divided sulphur, separated as a *white* powder from a sulphide, or *yellow* from a theiosulphate;\* this precipitate is recognized by its color, and by not being separable by standing or filtration, it may be disregarded; there may also be precipitated  $\text{H}_2\text{SiO}_4$  (gelatinous, almost colorless),  $\text{SnO}_2$  (or metastannic acid, see 420 a),  $\text{Sb}_2\text{O}_5$ ,  $\text{AgCl}$  (all three white),  $\text{SnS}_2$ ,  $\text{As}_2\text{S}_3$  (both yellow),  $\text{Sb}_2\text{S}_3$  (orange); hence a full examination of this precipitate for all that it may contain will involve boiling the precipitate with aqua regia, diluting and filtering; the insoluble residue is examined as a substance insoluble in water and acids (475), and the aqua regia solution is added to the filtrate from the precipitate caused by  $\text{HNO}_3$ , any precipitate caused on thus mixing the solutions being examined by Table I (433).

**420 a.** Metastannic acid is precipitated from a solution of an alkaline metastannate by addition of an acid: it forms a white precipitate, which becomes *golden-yellow* when moistened with  $\text{SnCl}_2$  solution; it is further recognized by its dissolving whilst being washed in the washing-water, but being reprecipitated when the washings run into the acid filtrate.

**421.** Even if  $\text{HCl}$  has been used in dissolving the substance, it is advisable to add a few drops of the acid to make sure that no member of Group I remains.

**422.** It must be remembered that  $\text{HCl}$  precipitates (in addition to  $\text{AgCl}$ ,  $\text{PbCl}_2$ ,  $\text{Hg}_2\text{Cl}_2$ ) strong Ba solutions; this precipitate however disappears on diluting with water and heating, thus differing from the three chlorides mentioned above.

**423.** If  $\text{HNO}_3$  or aqua regia has been used in dissolving the substance, or if the solution smells of  $\text{Cl}$  or  $\text{SO}_2$ , it should be boiled down considerably until after adding  $\text{HCl}$  and boiling, neither of the above gases is smelt; it is then diluted, and  $\text{H}_2\text{S}$  passed whether the dilution has caused any precipitate or not.

If As has been detected in the preliminary examination,  $\text{H}_2\text{SO}_4$  should be added until the liquid smells of  $\text{SO}_2$  after having been boiled for a short time, and the liquid is then heated for some time short of boiling until it no longer smells of  $\text{SO}_2$ . This treatment reduces arsenic or stannic compounds to the arsenious and stannous condition. In the case of As this reduction is to be recommended, since arsenic compounds are not easily precipitated by  $\text{H}_2\text{S}$ ; it is also very desirable to reduce stannic compounds, since  $\text{SnS}_2$  (unlike  $\text{SnS}$ ) readily runs through the filter, and being yellow in color may be passed over as S. It will of course be necessary to

---

\* Often called a "hyposulphite."

ascertain by special tests (187-190 and 201-203), made on the *original solution*, in which state As and Sn were originally present. If any precipitate is produced by boiling with  $\text{H}_2\text{SO}_4$  other than white sulphur, refer to (424).

**424.** Whilst boiling with  $\text{H}_2\text{SO}_4$  some  $\text{H}_2\text{SO}_3$  is usually formed, which may partially or completely precipitate Pb, Ba, Sr as white sulphates;  $\text{H}_2\text{SO}_3$  will also possibly precipitate Au, which causes the cool liquid to appear blue by transmitted and red by reflected light, and on boiling separates as a black powder. Examine any precipitate thus formed by  $\text{H}_2\text{SO}_3$  by Table G (434).

**425.** On addition of water, Bi, Sb, Sn may give white precipitates of their oxychlorides; these precipitates may be disregarded, since  $\text{H}_2\text{S}$  readily converts them into sulphides.

**426.**  $\text{H}_2\text{S}$  often causes a fine *white* precipitate of S, owing to the presence of certain oxidizing substances, such as Cl,  $\text{HNO}_3$ ,  $\text{HClO}_3$ ,  $\text{H}_2\text{CrO}_4$ ,  $\text{H}_2\text{MnO}_4$ ,  $\text{HMnO}_4$ ,  $\text{H}_2\text{SO}_3$ ,  $\text{Fe}_2\text{Cl}_6$ , etc.; this precipitate is known by its perfect whiteness, and by not being separable by standing or filtration, it may be neglected; great care must, however, be taken not to pass over  $\text{SnS}_2$  as S; it resembles S in running through filter-paper, but is distinctly *yellow* in color; it may often be coagulated by being shaken or heated.

**426 a.** Certain changes may be observed whilst  $\text{H}_2\text{S}$  is being passed; they should be carefully noted. Thus Pb and Hg, during precipitation by  $\text{H}_2\text{S}$ , show characteristic colors (149, 144); a solution colored reddish-yellow by  $\text{H}_2\text{CrO}_4$  becomes green; and a solution colored green by  $\text{H}_2\text{MnO}_4$  or purple by  $\text{HMnO}_4$  becomes colorless.

**427.** If on boiling the filtrate or solution after passing  $\text{H}_2\text{S}$  a yellow precipitate forms, this shows the presence of an arsenic or a stannic compound; in this case pass  $\text{H}_2\text{S}$  into the boiling-hot liquid as long as it causes any further precipitate, filter off and add this precipitate to any precipitate already obtained by  $\text{H}_2\text{S}$  to be examined by Table II.

**428.** Other substances may be present in this residue besides  $\text{SiO}_2$ , but  $\text{SiO}_2$  is distinguished from them all by readily dissolving in hot  $\text{KHO}$  solution.

The other substances which may remain undissolved by  $\text{HCl}$  are  $\text{Al}_2\text{O}_3$ ,  $\text{Fe}_2\text{O}_3$ ,  $\text{Cr}_2\text{O}_3$ , rendered difficultly soluble by the ignition to which the residue has been subjected; they are dissolved by long-continued heating with strong  $\text{HCl}$ , and the solution is added to the other solution which is to be examined for Group III A.

$\text{BaSO}_4$  and  $\text{SrSO}_4$  may also remain undissolved; they are detected by heating some of the residue strongly for a short time in the inner blowpipe flame upon a loop of platinum wire, then moistening the loop with a drop of strong  $\text{HCl}$  and holding it in the outer part of the Bunsen flame; crimson-red flashes, appearing red through the indigo-prism, prove the presence of Sr; a yellowish-green flame coloration shows Ba.

A less ready method of detecting Ba and Sr is to dry the insoluble residue, fuse it on platinum foil with three or four times as much fusion mixture; allow it to cool, then boil with water until the mass is disintegrated, filter, wash the residue well upon the filter, and examine it by Table IV (438) for Ba and Sr only.

**429.** Mn and Zn are very liable to be precipitated with  $\text{Fe}_2\text{H}_6$ ,  $\text{Al}_2\text{H}_6$ ,  $\text{Cr}_2\text{H}_6$ . The precipitation of Mn is partly prevented by keeping the liquid from exposure to the air after adding  $\text{AmHO}$ ; but the further precaution should be taken of dissolving the precipitate produced by  $\text{AmHO}$  in  $\text{HCl}$  and reprecipitating it by  $\text{AmHO}$ , then filtering and adding the filtrate to the original filtrate from the Group III A precipitate: this treatment of the precipitate should be repeated several times, if small quantities of Zn or Mn have to be looked for in presence of Al or Fe; or better still, Groups III A and III B may be precipitated together (459) and the precipitate examined by Table III c (464, 465).

**430.** The filtrate or solution, after adding  $\text{AmCl}$  and  $\text{AmHO}$ , may be colored *blue* by Ni, or *reddish-violet* by  $\text{Cr}_2\text{H}_6$ , which is dissolved in the excess of  $\text{AmHO}$ .

The  $\text{Cr}_2\text{H}_6$  must be removed before proceeding to examine for the three remaining groups; it is readily separated as a pale-green flocculent precipitate by boiling the liquid for a short time in a porcelain dish, adding a little more  $\text{AmHO}$  if the liquid does not smell of  $\text{NH}_3$  after being boiled for a short time; this precipitate is filtered off, and the filtrate, which is colorless if the Cr has been entirely removed by boiling the liquid sufficiently long, is further examined for Groups III B, IV, and V, by the General Table.

The coloration due to Ni differs from that caused by  $\text{Cr}_2\text{H}_6$  in not being thus removed by boiling the liquid.

**431.** A *brown* or *dark-colored* filtrate shows that Ni is present,\* some of the  $\text{NiS}$  precipitated by  $\text{Am}_2\text{S}$  having been dissolved in the excess of  $\text{Am}_2\text{S}$  and giving the filtrate its dark color. If this coloration is noticed the filtrate should be poured into a porcelain dish and boiled, occasionally adding water if necessary, until on removing the lamp for a short time the black  $\text{NiS}$  settles, leaving a colorless liquid. The liquid is then run through a small filter, the colorless filtrate examined by the General Table for Groups IV and V, and the black precipitate tested for Ni by fusing a portion of the filter-paper stained by the  $\text{NiS}$  in a colorless borax bead; if a bead is obtained which when heated in the outer flame is *violet* or *brown* whilst hot and *yellow* when cold, and *which becomes gray and opaque in the inner flame*, Ni is certainly present, and need not be further tested for in Table III B.

**432.** Part of the precipitate produced by  $\text{Am}_2\text{CO}_3$  often adheres firmly to the inside of the vessel in which it has been produced; in this case rinse the tube out several times with distilled water and then dissolve the precipitate by pouring in some hot  $\text{H}\bar{\text{A}}$ , causing it to run over the inside of the tube, and add this solution to the  $\text{H}\bar{\text{A}}$  solution of the  $\text{Am}_2\text{CO}_3$  precipitate made in Table IV.

\* If Groups III A and III B have been precipitated together by  $\text{AmCl}$ ,  $\text{AmHO}$ , and  $\text{Am}_2\text{S}$  added in succession (459) the brown coloration of the filtrate may also be due to  $\text{Cr}_2\text{H}_6$ : on boiling the filtrate colored by  $\text{Cr}_2\text{H}_6$  in a porcelain dish for several minutes, *pale-green*  $\text{Cr}_2\text{H}_6$  is precipitated, which cannot be mistaken for *black*  $\text{NiS}$ .

## 433. TABLE I.—SILVER GROUP.

The precipitate produced by HCl may consist of  $\text{PbCl}_2$ ,  $\text{AgCl}$ , and  $\text{Hg}'_2\text{Cl}_2$ ; the precipitate on the filter is washed five or six times with boiling water, the first portions of the washings being kept apart :

<p>The first portions of the washings may contain <math>\text{PbCl}_2</math> in solution, which if present in any quantity will be deposited in brilliant crystals on cooling. Test for, or confirm, the presence of Pb, by adding to the washings <math>\text{K}_2\text{CrO}_4</math>; a yellow precipitate. readily soluble in <math>\text{KHO}</math>:</p> <p><i>Presence of Pb.</i></p>	<p><i>Residue</i> : pour upon the precipitate on the filter some hot <math>\text{AmHO}</math>; when this has run through, heat it again, and pour it once more upon the filter :</p> <table border="1" data-bbox="384 500 835 745"> <tr> <td data-bbox="384 500 570 745"> <p><i>Filtrate</i> : add <math>\text{HNO}_3</math> until the liquid is acid: white precipitate, uniting into curdy particles when shaken or boiled:</p> <p><i>Presence of Ag.</i></p> </td><td data-bbox="570 500 835 745"> <p><i>Residue</i> on the filter is black; dry at a gentle heat (best in a steam bath), scrape the precipitate off the filter, mix with some dry <math>\text{Na}_2\text{CO}_3</math>, and heat in a small tube closed at one end, observing the precautions in the note to par. 147. Small globules of Hg form on the sides of the tube, often appearing only when the inside of the tube is rubbed or examined with a lens:</p> <p><i>Presence of Hg' (Mercurousum).</i></p> </td></tr> </table>	<p><i>Filtrate</i> : add <math>\text{HNO}_3</math> until the liquid is acid: white precipitate, uniting into curdy particles when shaken or boiled:</p> <p><i>Presence of Ag.</i></p>	<p><i>Residue</i> on the filter is black; dry at a gentle heat (best in a steam bath), scrape the precipitate off the filter, mix with some dry <math>\text{Na}_2\text{CO}_3</math>, and heat in a small tube closed at one end, observing the precautions in the note to par. 147. Small globules of Hg form on the sides of the tube, often appearing only when the inside of the tube is rubbed or examined with a lens:</p> <p><i>Presence of Hg' (Mercurousum).</i></p>
<p><i>Filtrate</i> : add <math>\text{HNO}_3</math> until the liquid is acid: white precipitate, uniting into curdy particles when shaken or boiled:</p> <p><i>Presence of Ag.</i></p>	<p><i>Residue</i> on the filter is black; dry at a gentle heat (best in a steam bath), scrape the precipitate off the filter, mix with some dry <math>\text{Na}_2\text{CO}_3</math>, and heat in a small tube closed at one end, observing the precautions in the note to par. 147. Small globules of Hg form on the sides of the tube, often appearing only when the inside of the tube is rubbed or examined with a lens:</p> <p><i>Presence of Hg' (Mercurousum).</i></p>		

434. TABLE G.—EXAMINATION OF PRECIPITATE FORMED ON BOILING THE HCl SOLUTION WITH  $\text{H}_2\text{SO}_4$ . (See par. 424, p. 252.)

The precipitate produced by  $\text{H}_2\text{SO}_4$  may contain Au,  $\text{PbSO}_4$ ,  $\text{BaSO}_4$ , and  $\text{SrSO}_4$ ; the last three substances are perfectly white, and Au, if present, is therefore usually seen by coloring the precipitate brown or black; the Au is also usually visible during precipitation by  $\text{H}_2\text{SO}_4$ , or after the white sulphates have subsided, since it colors the liquid blue by transmitted, and reddish by reflected light.

Ba and Sr can often be at once detected (if present) by taking a little of the moist precipitate on a loop of platinum wire, heating in the inner blowpipe flame for some time, moistening with HCl, and examining the flame coloration; but a more reliable method of proceeding is the following:

Rinse the precipitate into a small porcelain dish, using as little water as possible; dissolve in the liquid a small crystal of  $\text{H}_2\text{T}$ , or add a few drops of strong  $\text{H}\bar{\text{A}}$ ; then add  $\text{AmHO}$ , until it is just in excess, and boil for a short time; let stand and decant through a filter:

<p><i>Solution</i> may contain <math>\text{PbSO}_4</math>; add <math>\text{H}\bar{\text{A}}</math> in excess, then <math>\text{K}_2\text{CrO}_4</math>; a yellow precipitate soluble in <math>\text{KHO}</math>:</p> <p><i>Presence of Pb.</i></p>	<p><i>Residue</i> : pour upon the residue in the dish a little HCl and a few drops of <math>\text{HNO}_3</math> heat to boiling, let stand, decant into a porcelain dish:</p> <table border="1" data-bbox="358 1279 835 1406"> <tr> <td data-bbox="358 1279 598 1406"> <p><i>Solution</i> : boil until the smell of Cl is removed, add several drops of fresh <math>\text{FeSO}_4</math> solution; blue coloration and reddish precipitate:</p> <p><i>Presence of Au.</i></p> </td><td data-bbox="598 1279 835 1406"> <p><i>Residue</i> : dry by gently heating the dish, fuse with fusion mixture on platinum foil, and examine the mass for Ba and Sr as directed in the last par. of (428).</p> </td></tr> </table>	<p><i>Solution</i> : boil until the smell of Cl is removed, add several drops of fresh <math>\text{FeSO}_4</math> solution; blue coloration and reddish precipitate:</p> <p><i>Presence of Au.</i></p>	<p><i>Residue</i> : dry by gently heating the dish, fuse with fusion mixture on platinum foil, and examine the mass for Ba and Sr as directed in the last par. of (428).</p>
<p><i>Solution</i> : boil until the smell of Cl is removed, add several drops of fresh <math>\text{FeSO}_4</math> solution; blue coloration and reddish precipitate:</p> <p><i>Presence of Au.</i></p>	<p><i>Residue</i> : dry by gently heating the dish, fuse with fusion mixture on platinum foil, and examine the mass for Ba and Sr as directed in the last par. of (428).</p>		

## 435. TABLE II.—COPPER AND ARSENIC GROUPS.

The precipitate produced by  $H_2S$  in the  $HCl$  solution, after having been well washed with hot water until the last few drops of the washing water, collected in a test-tube, give no precipitate or only a very slight milkiness on addition of  $AgNO_3$ , is removed from the filter and boiled with  $KHO$  or a small quantity of  $Am_2S_3^*$  and filtered:

## 435 a. GROUP II A.—COPPER GROUP.

<p><i>The Residue</i> may contain <math>HgS</math>, <math>PbS</math>, <math>Bi_2S_3</math>, <math>CuS</math>, <math>CdS</math>. After having washed the precipitate and allowed it to drain, remove it from the filter into a porcelain dish (26 d), pour upon it strong <math>HNO_3</math> sufficient to cover it completely, and heat gently as long as any red fumes come off, adding more strong <math>HNO_3</math> if necessary to prevent evaporation to dryness. Boil off nearly all the acid, add a little water and dilute <math>H_2SO_4</math>, let stand for some time, stirring occasionally, and filter:</p>			
<p><i>Residue</i> may contain <math>HgS</math> (black), <math>PbSO_4</math> (white), (note 1). Remove the precipitate into a porcelain dish, using as little water as possible; add a little <math>H_2A</math>, then <math>AmHO</math> in excess, boil and filter:</p>	<p><i>Filtrate</i> may contain <math>Bi</math>, <math>Cu</math>, <math>Cd</math>. Add <math>AmHO</math> in excess, boil and filter: [Note.—A blue coloration shows presence of <math>Cu</math>.]</p>		
<p><i>Residue</i>: Dry on the filter at a gentle heat (best in the steam oven). Scrape the dried residue off the filter (note 2), and heat it, mixed with dry <math>Na_2O_2</math>, in a small ignition-tube; globules of <math>Hg</math> form on the sides of the tube, which become visible when rubbed or when looked for by a lens:  <i>Presence of <math>Hg</math></i><sup>1</sup> (as Mercuricum).</p>	<p><i>Filtrate</i>: Add <math>H_2A</math> in excess, then <math>K_2CrO_4</math>, a yellow precipitate:  <i>Presence of</i></p>	<p><i>Precipitate</i> (not easily seen in dark-blue solution): wash with <math>H_2O</math>, water, dissolve off the filter by pouring upon it a few drops of boiling dilute <math>HCl</math>; pour this solution into a large quantity of cold distilled water, a milkiness appearing at once or after still some time, shows <math>Cu</math>.  <i>Presence of <math>Bi</math></i>. [Note.—A precipitate produced by <math>AmHO</math> must always be tested as above. <math>Bi</math>, since <math>Pb</math> and <math>Ag</math> might possibly appear here as white precipitates, and be mistaken for <math>Bi</math>.]</p>	<p><i>Filtrate</i>: Add <math>HCl</math> until the liquid is acid, then pass <math>H_2S</math> to saturation, filter and wash quickly, keeping the filter covered with precipitate; remove the precipitate to a small porcelain dish, pour upon it some dilute <math>H_2SO_4</math>, boil and filter on a covered filter (note 3):</p>
	<p><i>Residue</i>: Dissolve by pouring a little boiling <math>HNO_3</math> upon the filter; add to the solution <math>AmHO</math> in excess, then <math>H_2A</math> in excess, and stand for some time, stirring occasionally, and filter:  <i>Presence of <math>Cu</math></i>.</p>	<p><i>Filtrate</i>: Add much water and pass <math>H_2S</math> for some time (or add much <math>H_2O</math>, water), a yellow precipitate shows:  <i>Presence of <math>Cd</math></i>. [Note.—If this precipitate is dark-colored, <math>CuS</math> is present (see note 3); filter it off, boil it once more with <math>H_2SO_4</math>, filter and test the filtrate by <math>H_2S</math>, as above for <math>Cd</math>.]</p>	

\* If traces of  $Cu$  have to be tested for,  $Na_2S$  should be substituted for  $Am_2S_3$ , since  $CuS$  is somewhat soluble in  $Am_2S_3$ .



## GROUP II B.—ARSENIC GROUP.

The filtrate may contain  $\text{As}_2\text{S}_3$ ,  $\text{Sb}_2\text{S}_3$ ,  $\text{SnS}_2$  ( $\text{SnS}$ ) [ $\text{Au}_2\text{S}_3$ ,  $\text{PtS}_2$ ]. Add to it  $\text{HCl}$  until it is acid; the above sulphides if present will be reprecipitated; filter.

[*Note*.—A mere milkiness due to precipitation of  $\text{S}$  may be disregarded. If  $\text{KHO}$  or  $\text{NaHO}$  have been used above to dissolve these sulphides,  $\text{H}_2\text{S}$  must be passed into the acid filtrate, and any further precipitate thus obtained must be added to the precipitate obtained on acidifying; this precaution is unnecessary if  $\text{Au}_2\text{S}_3$  or  $\text{Na}_2\text{S}$  has been employed.]

The following method of examining the precipitate may then be employed (see note 4, below):

435 b. Remove the precipitate from the filter, using as little water as possible, into a porcelain dish; add a little strong  $\text{HCl}$  and heat, occasionally dropping in a crystal of  $\text{KClO}_3$  until the precipitate dissolves entirely, or leaves only a slight residue of sulphur. Heat this solution just short of boiling (note 5, until it ceases to smell of  $\text{Cl}$  (note 6), then cool it and pour it (through a filter, if necessary) into a small flask (181) containing pieces of pure  $\text{Zn}$ ; if the gas does not come off readily, add a little strong  $\text{HCl}$ , and cause the gas, after passing through some  $\text{PbAc}_2$  solution (see caution 181), to bubble through some  $\text{AgNO}_3$  solution contained in a broad test-tube. As soon as the evolution of gas slackens, drop in more  $\text{Zn}$ , unless some remains undissolved in the flask; and as soon as the gas ceases to come off (some  $\text{Zn}$  still remaining undissolved), the test-tube containing the  $\text{AgNO}_3$  solution is removed; if any black precipitate has formed in this solution, it must be tested as below; if the liquid has remained clear and colorless, it may be rejected, since  $\text{As}$  and  $\text{Sb}$  are absent.

Residue in the flask may consist of  $\text{Sn}$  ( $\text{Pt}$ ,  $\text{Au}$ ), and excess of  $\text{Zn}$ ; remove any deposit of  $\text{Sn}$  by shaking the pieces of  $\text{Zn}$  well with the liquid, and quickly pouring off into a dish, decant the liquid and wash the  $\text{Sn}$  by decantation in the dish; boil the residue with a little strong  $\text{HCl}$  in a test-tube, then quickly dilute, decant (note 7), and add  $\text{HgCl}_2$  to the solution; a white precipitate (becoming black with  $\text{AmHO}$ ) shows:

*Presence of Sn* (note 8).

To the liquid in the test-tube add several drops of  $\text{AgNO}_3$  solution; filter:

*Precipitate*: Wash well with boiling water until  $\text{HCl}$  gives no milkiness with the last few drops of the washing-water, then pour upon the filter a little boiling dilute  $\text{H}_2\text{I}$  solution; boil it and pour it again through the filter; repeat this several times, then add to the solution  $\text{HCl}$  and pass  $\text{H}_2\text{S}$ ; an orange-red precipitate shows:

*Presence of Sb.*

*Note*.—This precipitate is often only seen on settling from the yellow liquid.

*Filtrate*: Dilute several drops of  $\text{AmHO}$  largely in a test-tube, and add this liquid drop by drop to the filtrate; a light-yellow precipitate, which settles into flocks on shaking the liquid, shows:

*Presence of As* (note 8).

*Note 1*.—A white residue does not necessarily show the absence of  $\text{Hg}$ .

*Note 2*.—If the amount of the residue is too small to be removed; the portion of the paper containing it may be cut up and mixed with  $\text{Na}_2\text{CO}_3$ . For the precautions requisite in performing this test see par. (147).

*Note 3*.—Unless this precipitate is filtered in a covered funnel, washed quickly, and examined at once,  $\text{CuS}$  is liable to be oxidized and to pass into the filtrate as  $\text{CuSO}_4$ ; it then discolors the  $\text{CdS}$  precipitate, or may be entirely overlooked if present in small quantity.

*Note 4*.—The detection of  $\text{As}$ ,  $\text{Sb}$ ,  $\text{Sn}$  by the method given in

(435b) is very trustworthy and delicate (208); it is excelled in these respects by the method in (210). For beginners, however, the process (209) is recommended by its greater simplicity.

*Note 5.*—The solution must not be heated to boiling, else  $\text{SbCl}_5$  or  $\text{SnCl}_4$  might be volatilized if present.

*Note 6.*—In this solution As and Sn will be present as stannicum and arsenicum, and in this state are not well suited for detection; the following modification makes their detection much more rapid and certain, especially when they are present only in small quantity. After warming until the liquid ceases to smell of Cl, add  $\text{H}_2\text{SO}_4$  drop by drop, until the liquid after being warmed for a short time smells of  $\text{SO}_2$ ; then heat it short of boiling until the smell of  $\text{SO}_2$  is removed, and pour (through a filter, if necessary,) into the H apparatus. Great care must be taken to remove the  $\text{SO}_2$  entirely, else Sb and As may remain as sulphides in the H apparatus.

*Note 7.*—A residue here will contain Au or Pt if they are present; if they are to be tested for refer to (473 and 474).

*Note 8.*—The precipitation by  $\text{H}_2\text{S}$  in the General Table will often show in what condition As or Sn were present (427); further distinctive tests may be tried on the original solution by (187–190) and (201–203).

#### 436. TABLE III A.—IRON GROUP.

The precipitate produced by boiling, after addition of  $\text{AmCl}$  and of  $\text{AmHO}$  in excess, may contain  $\text{Al}_2\text{Ho}_3$ ,  $\text{Cr}_2\text{Ho}_3$ ,  $\text{Fe}_2\text{Ho}_3$ ; the color of the precipitate will usually show whether it contains any quantity of  $\text{Fe}_2\text{Ho}_3$  (reddish-brown), or  $\text{Cr}_2\text{Ho}_3$  (pale-green), since  $\text{Al}_2\text{Ho}_3$  is colorless.

Dissolve the precipitate by pouring upon the filter a little boiling dilute  $\text{HCl}$ , add to the solution pure  $\text{NaHO}$  in excess, boil for some time and filter:

<p>1. <i>Filtrate</i> may contain <math>Al_2H_6</math> dissolved in excess of <math>NaHO</math>. Add <math>HCl</math> gradually to the liquid, shaking meanwhile with a small piece of blue litmus-paper; just as the liquid becomes neutral, a colorless gelatinous precipitate will form, should <math>Al</math> be present. If acid is added in excess, this precipitate dissolves, but reappears on adding <math>AmHO</math> in excess. The appearance of this precipitate shows:</p> <p><i>Presence of Al.</i></p> <p><i>Note.</i>—Caustic soda and potash is very liable to contain <math>Al_2H_6</math>; the analyst must ascertain that the <math>Al</math> does not come from this source.</p>	<p>2. <i>Precipitate</i> may contain <math>Cr_2H_6</math> and <math>Fe_2H_6</math>. Dry upon the filter, carefully avoiding charring the paper; when sufficiently dried the substance will have shrunk considerably into small, hard, dark-colored pieces; detach these from the filter and fuse them with a little solid <math>Na_2CO_3</math> and <math>KNO_3</math> upon a piece of platinum foil with turned-up edges, and either supported upon a pipeclay triangle or held in the Bunsen flame by crucible-tongs. Continue the fusion by heating the under surface of the foil in the blowpipe flame as long as any frothing occurs; then allow to cool; a yellow mass indicates the presence of <math>Cr</math> (note 1). Boil the platinum foil in a small porcelain dish with water until the substance is entirely dissolved or only a dark-brown powder (<math>Fe_2O_3</math>) remains undissolved; take out the foil, allow the powder to settle, and decant the liquid through a filter:</p>		
	<table border="1"> <tr> <td data-bbox="383 552 600 758"> <p>3. <i>Residue</i> in porcelain dish; dissolve by pouring in a little <math>HCl</math> and boiling, add some water, then a few drops of <math>KCyS</math> solution; a blood-red coloration shows:</p> <p><i>Presence of Fe</i> (note 2).</p> </td><td data-bbox="600 552 832 758"> <p>4. <i>Filtrate</i> will be yellow if <math>Cr</math> is present (note 1 a). Add <math>HA</math> gradually until it is in excess (shown by the color becoming reddish), boil for several minutes, then add <math>PbA_2</math> solution; a yellow precipitate shows:</p> <p><i>Presence of Cr</i> (note 3).</p> </td></tr> </table>	<p>3. <i>Residue</i> in porcelain dish; dissolve by pouring in a little <math>HCl</math> and boiling, add some water, then a few drops of <math>KCyS</math> solution; a blood-red coloration shows:</p> <p><i>Presence of Fe</i> (note 2).</p>	<p>4. <i>Filtrate</i> will be yellow if <math>Cr</math> is present (note 1 a). Add <math>HA</math> gradually until it is in excess (shown by the color becoming reddish), boil for several minutes, then add <math>PbA_2</math> solution; a yellow precipitate shows:</p> <p><i>Presence of Cr</i> (note 3).</p>
<p>3. <i>Residue</i> in porcelain dish; dissolve by pouring in a little <math>HCl</math> and boiling, add some water, then a few drops of <math>KCyS</math> solution; a blood-red coloration shows:</p> <p><i>Presence of Fe</i> (note 2).</p>	<p>4. <i>Filtrate</i> will be yellow if <math>Cr</math> is present (note 1 a). Add <math>HA</math> gradually until it is in excess (shown by the color becoming reddish), boil for several minutes, then add <math>PbA_2</math> solution; a yellow precipitate shows:</p> <p><i>Presence of Cr</i> (note 3).</p>		

*Note 1.*—If  $Mn$  was present in the original substance, it is often precipitated more or less completely with the  $Fe$ , and will be detected by imparting to the fused mass a bluish-green color; this should be noted, as the  $Mn$  may possibly be entirely precipitated in this group, and will then not be detected in the next group.

*Note 1 a*—The green color, due to the presence of  $Mn$ , conceals the pale yellow color due to  $Cr$ ; but on dissolving the cool mass in water and boiling with  $HA$ , the manganese coloration is destroyed; and, on filtering from any dark-colored precipitate, the yellow color due to the  $Cr$  becomes evident.

*Note 2.*— $Fe$  is always detected here in the ferric condition, since even if originally present as ferrous, it will have been oxidized by boiling with  $HNO_3$  as directed in the General Table. To ascertain in which condition the  $Fe$  was present, boil some of the original substance with  $HCl$ , and divide the  $HCl$  solution (filtered if necessary) into two parts: to one portion add  $K_2Fe_2Cy_{12}$ , a dark-blue precipitate shows *presence of Fe'' (Ferrous)*: to another portion add  $KCyS$ , a blood-red color shows *presence of Fe''' (Ferric)*.

*Note 3.*— $Cr$  may have been present as a chromic salt ( $Cr$  replacing  $H$  in an acid) or a chromate ( $Cr$  being united with  $O$  to form an acid radicle). Ascertain in which state it was present by boiling some of the original substance with  $Na_2CO_3$  solution and filtering; a yellow filtrate, which on being acidified with  $HA$  gives a yellow precipitate with  $PbA_2$ , shows that  $Cr$  was present as a

*chromate*; Cr previously combined with an acid as a chromic salt, remains undissolved as green  $\text{Cr}_2\text{H}_2\text{O}_6$ , and does not color the filtrate: It is found by 2 (in the above Table).

---

#### 437. TABLE III B.—ZINC GROUP.

The precipitate produced by  $\text{Am}_2\text{S}$  may contain  $\text{ZnS}$ ,  $\text{MnS}$ ,  $\text{NiS}$ ,  $\text{CoS}$ . The color of the precipitate will usually show whether it contains any quantity of  $\text{NiS}$  or  $\text{CoS}$  which are black, or consists only of  $\text{MnS}$  (pink) or  $\text{ZnS}$  (white).

*Note.*—The presence of Ni, indicated by the filtrate, after boiling with  $\text{AmHO}$  in the General Table, being blue in color, and also by a dark-brown filtrate being obtained after boiling with  $\text{Am}_2\text{S}$ , will usually have been proved to a certainty by the examination in the borax bead of the black  $\text{NiS}$ , precipitated by boiling the dark-brown filtrate from  $\text{Am}_2\text{S}$  (431); if Ni has been thus detected its further detection in the  $\text{Am}_2\text{S}$  precipitate, as directed below, will be unnecessary.

Remove the precipitate from the filter with cold dilute  $\text{HCl}$  in the way described in par. 26 c; take out the paper and allow the liquid to stand, occasionally stirring it well. The precipitate will either dissolve leaving only white sulphur, in which case the milky liquid may be at once examined by 2 (below), or a black residue will be left which must be filtered off and examined by 1 (below), the filtrate being examined by 2:

<p>1. <i>Residue</i> (black) may contain <math>\text{NiS}</math>, <math>\text{CoS}</math>.  <i>Examination by the borax bead.</i>          Fuse a small quantity in a clear borax bead in the extreme tip of the outer blow-pipe flame, note its color; then fuse the bead for some time in the inner blow-pipe flame, and again observe its color. Any one of the following observations may be made:</p>	<p><i>Blue bead</i> in both flames shows:  <i>Presence of Co.</i>          Co need not be further tested for, but if any doubt exists as to the presence of Ni the rest of the residue must be examined as below.</p> <p><i>Brown or yellow bead</i>, when cold after fusing in the outer flame, which becomes <i>gray or opaque</i> in the inner flame, shows <i>Presence of Ni, Absence of Co.</i>          Unless more traces of Co have to be tested for, no further examination need be made.</p>	<p><i>Further examination of the residue.</i> — Rinse the residue from the filter into a porcelain dish, using as little water as possible: pour in a little strong <math>\text{HCl}</math> and boil for some time, adding occasionally a small crystal of <math>\text{KClO}_3</math> until the black residue is entirely dissolved or only a small quantity of dark sulphur remains; then boil down nearly to dryness (a blue liquid shows presence of <math>\text{Co}</math>), dilute with a little water, filter (if necessary) into a boiling-tube and pour in <math>\text{K}_2\text{Cr}_2\text{O}_7</math> solution slowly until the precipitate formed at first is just redissolved, boil briskly for several minutes, and add much strong <math>\text{Na}_2\text{CO}_3</math> (or make decidedly alkaline with <math>\text{NaOH}</math> and add <math>\text{Br}</math>-water); heat nearly to boiling, and allow the liquid to stand at least ten minutes, filter. (Note 1, below):</p>	<p>3. <i>Precipitate</i> (black): wash and confirm the presence of <math>\text{Ni}</math> by fusing some of the precipitate (or the paper stained with the precipitate) in a borax bead in the outer and inner blow-pipe flames: a bead <i>yellow</i> when cold, becoming <i>black</i> in the inner flame shows:  <i>Presence of Ni.</i></p> <p>4. <i>Filtrate</i> must be warmed with more <math>\text{Na}_2\text{CO}_3</math> (or <math>\text{Br}</math>-water), and filtered from any additional precipitate which may form, then evaporated to dryness, and strong <math>\text{HNO}_3</math> poured upon the residue as long as any frothing is caused; this is then evaporated just to dryness, the residue dissolved in water, excess of <math>\text{KHO}</math> added to it, and any precipitate filtered off and fused into a borax bead (taking the paper stained by the precipitate if the precipitate is not easily removed): a blue bead shows:  <i>Presence of Co.</i></p>
<p>2. <i>Solution or filtrate</i> may contain <math>\text{Zn}</math>, <math>\text{Mn}</math>.          Boil in a porcelain dish until it ceases to smell of <math>\text{H}_2\text{S}</math>, then drop in a small crystal of <math>\text{KClO}_3</math> and boil for several minutes. After cooling the liquid add pure <math>\text{NaOH}</math> until after stirring the liquid it turns red litmus-paper blue, stir well and filter:</p>	<p>5. <i>Precipitate is white at first, but rapidly darkens</i> in the air. Fuse a portion of it with solid <math>\text{Na}_2\text{O}_2</math> and <math>\text{KNO}_3</math> on platinum; a bluish-green mass is obtained on cooling, showing:  <i>Presence of Mn.</i>  <i>Note.</i> — <math>\text{Mn}</math> may have been originally present, either replacing <math>\text{H}</math> in an acid, or combined with <math>\text{O}</math> as an acid radicle. Its presence in an acid radicle is indicated by the color of the original solution, since in the green, of manganate, it is green, and as permanganate it is purple; these colors disappear on boiling the <math>\text{HCl}</math> solution for some time, or on passing <math>\text{H}_2\text{S}</math>.</p>	<p>6. <i>Filtrate</i>, pass <math>\text{H}_2\text{S}</math>, a white precipitate forms, often somewhat discolored, showing:  <i>Presence of Zn.</i>  <i>Note.</i> — From a dilute solution of <math>\text{ZnH}_2\text{O}_2</math> in <math>\text{NaOH}</math> the <math>\text{ZnH}_2\text{O}_2</math> is precipitated on boiling; hence after adding <math>\text{NaOH}</math>, as directed above, the liquid must not be boiled, else <math>\text{Zn}</math> might be precipitated with the <math>\text{MnH}_2\text{O}_2</math> and escape detection. But by diluting and boiling the above filtrate it is usually possible to cause any <math>\text{ZnH}_2\text{O}_2</math> it holds in solution to precipitate (especially if the alkalinity of the liquid be somewhat reduced by cautious addition of <math>\text{HCl}</math> in quantity insufficient to make the liquid acid). If this precipitate be separated by pouring the liquid through a double filter and then be dissolved off the filter and the sides of the boiling-tube by a little boiling <math>\text{H}_2\text{A}</math>, <math>\text{ZnS}</math> may be precipitated perfectly white from this solution, by <math>\text{H}_2\text{S}</math> showing:  <i>Presence of Zn.</i></p>	<p><i>Note 1.</i> — If this precipitate adheres to the sides of the boiling-tube, it is dissolved off after the liquid has been emptied out) by pouring in a little boiling <math>\text{HCl}</math>; from this solution the <math>\text{Ni}</math> is precipitated by addition of <math>\text{KHO}</math>, and the precipitate filtered and tested by the borax bead from <math>\text{Ni}</math>.</p>

## 438. TABLE IV.—BARIUM GROUP.

The precipitate produced by  $\text{Am}_2\text{CO}_3$  may contain  $\text{BaCO}_3$ ,  $\text{SrCO}_3$ ,  $\text{CaCO}_3$ ; it is to be well washed with boiling water. Before dissolving the precipitate off the filter much may usually be learnt by an examination of the flame coloration (see 1 below) yielded by the  $\text{HCl}$  solution. The results thus obtained are afterwards confirmed in the wet way (see 2 below).

1. *Examination by Flame Coloration.*—Take a small quantity of the precipitate out of the filter upon the end of a glass rod, and dissolve it by moving the end of the rod about in several drops of  $\text{HCl}$  on a watch-glass; dip into this solution a loop of platinum wire which imparts no color to the flame, then hold the loop for some time in the Bunsen flame, repeating the process if the coloration is not satisfactorily observed at first.  $\text{Ca}$  will impart to the flame a *yellowish-red* color,  $\text{Sr}$  a *crimson-red*,  $\text{Ba}$  a *yellowish-green*. The red colorations yielded by  $\text{Ca}$  and  $\text{Sr}$  are distinguished by viewing the flame through the indigo-prism; the  $\text{Ca}$  coloration then appears *dingy-green*, whilst the  $\text{Sr}$  coloration still appears *crimson*;  $\text{Ba}$  is usually found by its coloration remaining after the others have disappeared. When all three metals are present together their colorations are apt to interfere with one another; but if at any time a red color is visible through the indigo-prism,  $\text{Sr}$  is certainly present; if a green coloration is visible without using the prism, the presence of  $\text{Ba}$  is proved.

2. *Examination in the Wet Way.*—The rest of the precipitate is dissolved off the filter by pouring upon it a little boiling  $\text{H}\bar{\text{A}}$ , again boiling the liquid, and pouring it once more through the filter if the precipitate is not entirely dissolved. Test whether the solution is acid, and if it is not add  $\text{H}\bar{\text{A}}$  whilst stirring until the liquid is acid, then pour off a small portion (*A*) of the acid solution into a test-tube, and put by the larger portion (*B*). To the portion (*A*), after it has been cooled by immersing the lower part of the tube in cold water or holding it in a stream from the tap, add several drops

of  $\text{CaSO}_4$  solution; if an immediate precipitate forms refer to I (below); if no immediate precipitate forms heat the liquid to boiling, and if a precipitate then forms refer to II, if no precipitate forms even after several minutes refer to III.

<p>I. An immediate precipitate is produced by <math>\text{CaSO}_4</math> in the cold, showing: <i>Presence of Ba.</i>          Portion (B) must then be tested for Sr and Ca which may also be present: examine it as directed below:          Add <math>\text{K}_2\text{CrO}_4</math> solution until the color of the liquid is reddish-yellow; heat and filter through a double filter-paper, pouring the filtrate through the same filter repeatedly if necessary until the liquid is quite clear; then add to the liquid, which must be orange-red in color, <math>\text{AmHO}</math> until the color changes to pale yellow, then add <math>\text{Am}_2\text{CO}_3</math> solution:</p>		
<p>No precipitate forms, showing:          Absence of Sr and Ca.</p>	<p>A precipitate forms, showing Sr, Ca, or both of them, to be present. Add to the liquid <math>\text{Am}_2\text{CO}_3</math> in excess, filter, reject the filtrate and dissolve the precipitate off the filter in as little boiling <math>\text{H}\bar{\text{A}}</math> as possible; pour off a small part (A') of this solution, reserving the larger portion (B'). To (A') add <math>\text{CaSO}_4</math> solution and boil:</p>	
	<p>No precipitate forms:          Absence of Sr.          Examine portion (B') for Ca as directed at (C) in III.</p>	<p>A precipitate forms:          Presence of Sr.          Examine portion (B') for Ca as directed at (C), II.</p>

II. A precipitate is not formed at once on the addition of  $\text{CaSO}_4$ , but appears on boiling the liquid, showing:

*Absence of Ba and Presence of Sr.*

C. Portion (B) is then examined for Ca as directed below: Add dilute  $\text{H}_2\text{SO}_4$  in excess, boil and filter, reject the precipitate; add to the filtrate several drops more  $\text{H}_2\text{SO}_4$  and boil; if this causes any precipitate boil and filter, and again test the filtrate by addition of  $\text{H}_2\text{SO}_4$  and boiling: repeat this process if necessary. To the clear filtrate, which gives no further precipitate on addition of  $\text{H}_2\text{SO}_4$  and boiling, add gradually  $\text{AmHO}$  until after mixing the liquid by thorough stirring or shaking it turns red litmus-paper blue, then add  $\text{Am}_2\text{C}_2\text{O}_4$  solution and warm gently; a white precipitate, often appearing only after a time, shows:

*Presence of Ca.*  
 (See note below.)

III. No precipitate is produced by  $\text{CaSO}_4$ , even on boiling, showing:

*Absence of Ba and Sr and Presence of Ca.*

C. Confirm the presence of Ca by adding to portion (B) excess of  $\text{AmHO}$  (35 a), then  $\text{Am}_2\text{C}_2\text{O}_4$  solution; a white precipitate forms, showing:

*Presence of Ca.*

(See note below.)

*Note.*—If mere traces of Ca have to be tested for, the filter-paper used in the analysis must first be freed from any traces of Ca it may contain by wetting it with dilute  $\text{HCl}$ , and then thoroughly washing it with distilled water, as described in the note to par. 23 a (p. 59).

#### 439. TABLE V.—POTASSIUM GROUP.

The filtrate, after addition to the original solution of all the group-reagents in succession, may still contain  $\text{Mg}$ ,  $\text{K}$ ,  $\text{Na}$ , and  $\text{NH}_4$ . Since, however,  $\text{NH}_4$  salts have been added as group-reagents, it is useless to test for  $\text{NH}_4$  in this filtrate;  $\text{NH}_4$  must be detected by boiling a portion of the original substance with  $\text{KHO}$  solution, and ascertaining whether any  $\text{NH}_3$  gas is evolved, which would be recognized by its smell or its action on moistened red litmus-paper. It is, however, usually unnecessary to test specially here for  $\text{NH}_4$ , since it will have been already tested for in the preliminary examination (390). Proceed to examine for Group V as directed below.

Evaporate the filtrate from Group IV to dryness in a porcelain dish, scrape out the solid residue and heat it to redness upon a piece of platinum foil in the Bunsen flame as long as any white fumes are seen to be given off on removing the foil for an instant from the flame (Note 1). All  $\text{Am}$  compounds are thus entirely removed. If any residue remains (Note 2) it is to be dissolved by boiling the foil in a test-tube with a small quantity of water to which several drops of dilute  $\text{HCl}$  have been added. Divide this solution into two equal parts:



<p><b>EXAMINATION FOR Mg.</b></p> <p>Before testing for Mg in one portion of this solution it is necessary to remove from it any traces of Ba, Sr, or Ca which it may possibly contain, and which might else be mistaken for Mg.</p> <p>Add, therefore, several drops of <math>\text{H}_2\text{SO}_4</math> to the liquid, boil for a short time and let stand; then, whether a precipitate has formed or not, add a little <math>\text{AmCl}</math> and <math>\text{AmHO}</math> in excess, then several drops of <math>\text{Am}_2(\text{C}_2\text{O}_4)_3</math>, and warm gently; if any precipitate has formed proceed to filter at once as quickly as possible and test the filtrate once more by adding <math>\text{H}_2\text{SO}_4</math>, then <math>\text{AmCl}</math>, <math>\text{AmHO}</math> in excess, and <math>\text{Am}_2(\text{C}_2\text{O}_4)_3</math>, and warming as directed above; and if any further precipitate forms, filter and again test the filtrate. To the liquid in which the addition of <math>\text{H}_2\text{SO}_4</math>, <math>\text{AmCl}</math>, <math>\text{AmHO}</math> in excess, and <math>\text{Am}_2(\text{C}_2\text{O}_4)_3</math> causes no precipitate, add <math>\text{Na}_2\text{HPO}_4</math>, and if no precipitate forms at once warm gently, shake or stir the liquid violently and let it stand for some time: a white, crystalline precipitate shows:</p> <p style="text-align: center;"><i>Presence of Mg.</i></p>	<p><b>EXAMINATION FOR K AND Na.</b></p> <p>In the other portion proceed to test for K and Na as directed below. The presence of Mg in no way interferes with the detection of K and Na, and hence if it has been detected its removal is unnecessary.</p> <p><i>Flame coloration.</i>—Dip into the solution a loop of platinum-wire which has been proved when held in a Bunsen flame not to impart any color to it. Hold the loop in the Bunsen flame; one of the following results will be observed:</p> <table border="1" data-bbox="481 495 833 706"> <tr> <td data-bbox="481 503 699 698"> <p><i>A bright yellow flame coloration, indicating the presence of Na (note 3). Examine this coloration through the indigo-prism; it appears red, indicating presence of K: if no red color is visible K is probably absent or present only in very minute quantity.</i></p> </td><td data-bbox="709 503 833 698"> <p><i>A pale violet coloration, appearing red or purple through the indigo-prism, shows:</i></p> <p style="text-align: center;"><i>Presence of K, absence of Na.</i></p> </td></tr> </table> <p>It is usual to confirm the results of the examination of the flame coloration by pouring the remainder of the solution upon a watch-glass, adding to it several drops of <math>\text{PtCl}_4</math> (note 4) and stirring well for some time; the formation of a yellow precipitate shows:</p> <p style="text-align: center;"><i>Presence of K (note 5).</i></p>	<p><i>A bright yellow flame coloration, indicating the presence of Na (note 3). Examine this coloration through the indigo-prism; it appears red, indicating presence of K: if no red color is visible K is probably absent or present only in very minute quantity.</i></p>	<p><i>A pale violet coloration, appearing red or purple through the indigo-prism, shows:</i></p> <p style="text-align: center;"><i>Presence of K, absence of Na.</i></p>
<p><i>A bright yellow flame coloration, indicating the presence of Na (note 3). Examine this coloration through the indigo-prism; it appears red, indicating presence of K: if no red color is visible K is probably absent or present only in very minute quantity.</i></p>	<p><i>A pale violet coloration, appearing red or purple through the indigo-prism, shows:</i></p> <p style="text-align: center;"><i>Presence of K, absence of Na.</i></p>		

*Note 1.*—If the residue is small in amount it may be ignited in the dish; but this is not to be generally recommended, as the porcelain dish is liable to be cracked by the heat, and it is also difficult to get entirely rid of the  $\text{NH}_4$  salts by heating only in porcelain.

*Note 2.*—It is not safe to place much reliance upon an examination of the foil for the detection of a small quantity of residue, but it may usually be detected by its producing a crackling noise whilst the foil is cooling immediately after its removal from the flame. Should there be any doubt, the foil must be boiled with water and a few drops of  $\text{HCl}$ , and the solution examined for Mg, K, and Na, as directed above.

*Note 3.*—A yellow coloration, more or less intense, will almost always be obtained here, since nearly all substances and reagents contain small quantities of Na; hence the student must note the intensity of the coloration, and judge from it whether the quantity of Na is small or large; he must then enter accordingly, either "*Presence of Na,*" or "*Presence of trace of Na.*"

*Note 4.*—If iodine is present,  $\text{PtCl}_4$  will produce an intense red coloration; hence if iodine is suspected to be present, before add-

ing  $\text{PtCl}_4$ , the above solution should be evaporated to dryness with a little strong  $\text{HNO}_3$ , the residue dissolved in a few drops of dilute  $\text{HCl}$  and tested with  $\text{PtCl}_4$  for  $\text{K}$ . This is of course unnecessary if, before precipitating Group III in the General Table, the solution has been evaporated to dryness with  $\text{HNO}_3$ . In case of uncertainty, add  $\text{PtCl}_4$  to a drop only of the above solution; and if iodine is found to be present proceed as directed above.

*Note 5* — Mere traces of  $\text{K}$  and  $\text{Na}$  may be detected by adding  $\text{PtCl}_4$ , and evaporating the liquid to dryness in a porcelain dish upon a water-bath (504); then pouring absolute alcohol into the dish and stirring; any yellow residue shows *presence of K*, the solution giving the pure  $\text{Na}$  coloration if  $\text{Na}$  is present. By filtering off the yellow residue and washing it with absolute alcohol, it will yield the pure  $\text{K}$  flame coloration.

#### EXAMINATION FOR ACID RADICLES.

**440.** Several acid radicles will probably have been already detected for certain, if present, by the preliminary examination. Such are  $\text{CO}_3$ ,  $\text{SO}_3$ ,  $\text{S}$ ,  $\text{NO}_2$ ,  $\text{ClO}$ ,  $\text{NO}_3$ ,  $\text{ClO}_3$ ,  $\bar{\text{A}}$ : their reactions are so characteristic that, unless interfered with by the presence of other substances, their presence or absence will have been proved to a certainty: in case any uncertainty is felt, the analyst should refer to the reactions given in Section IV for the particular acid radicle in question, and select a reaction which will be decisive.

The examination for metals also frequently yields proof of the presence of certain acid radicles. Thus on passing  $\text{H}_2\text{S}$  into the  $\text{HCl}$  solution:

- |  |  |
|--|--|
| A reddish-yellow solution, becoming green and depositing white sulphur, shows:   | <i>Presence of <math>\text{CrO}_4</math></i>     |
| A green solution, becoming colorless,  | <i>Presence of <math>(\text{MnO}_4)''</math></i> |
| A purple solution, " "   | <i>Presence of <math>(\text{MnO}_4)'</math></i>  |
| A yellow precipitate of $\text{As}_2\text{S}_3$ , appearing only when the liquid is boiled, shows:                             | <i>Presence of <math>\text{AsO}_4</math></i>     |
| An insoluble residue of $\text{SiO}_2$ , after evaporating to dryness the filtrate from the $\text{H}_2\text{S}$ group, shows: | <i>Presence of <math>\text{SiO}_2</math></i>     |

And a yellow precipitate obtained with  $\text{AmHMoO}_4$  before precipitating Group III, shows: *Presence of  $\text{PO}_4$ .*

The presence of any of these acid radicles thus detected will require no further confirmation.

**441.** Of the acid radicles which remain to be tested for, some can be detected by the plan drawn out below (442-445): others are best detected by special tests made on the original substance (446-454).

Much time and trouble may usually be saved by considering what acid radicles can possibly be present. A careful preliminary examination will usually have limited this number considerably, and it may be further reduced by the knowledge already obtained as to the solubility of the substance under analysis, and the metals it contains. A reference to the Table of Solubilities (455), and the explanatory paragraphs, 456, 457, and 458, will show how this knowledge is applied.

#### GENERAL EXAMINATION FOR CERTAIN ACID RADICLES.

**442.** Before employing the liquid tests it is advisable to separate from the substance any metals other than K, Na, and Am, which it may contain, since some of these are liable to be precipitated by the reagents added for the detection of the acid radicles. If alkali metals alone are present this separation is unnecessary, since they are not precipitated by any of the reagents.

This separation of the metals may usually be effected by boiling a portion of the substance with  $\text{Na}_2\text{CO}_3$  solution, which must be added as long as it causes any precipitate. Filter from the precipitate, and divide the clear filtrate into five equal portions. Reserve one of these portions in case of accident, and a second for the tests for organic acid radicles, and acidify the others whilst they are hot by addition of  $\text{HCl}$ ,  $\text{HNO}_3$ , and  $\text{H}\bar{\text{A}}$  respectively: examine these portions as directed below, using a separate part for each test:

443. I. Portion acidified with HCl.		Present.	
On addition of $\text{BaCl}_2$ solution, a white precipitate insoluble on boiling (444, 1), . . . . . }		$(\text{SO}_4)''$ .	
On addition of $\text{BaCl}_2$ solution, a semi-transparent precipitate insoluble on boiling (444, 2), . . . . . }		$(\text{SiF}_6)''$ .	
On addition of $\text{AmCl}$ and $\text{Am}_2\text{CO}_3$ , a semi-transparent precipitate, . . . . . }		$(\text{SiO}_2)''$ .	
On addition of $\text{FeSO}_4$ solution, a dark-blue precipitate, . . . . . }		$(\text{Fe}_3\text{Cy}_{15})^{\text{vi}}$ .	
On addition of $\text{Fe}_2\text{Cl}_6$ solution, a dark-blue precipitate: } $\text{FeSO}_4$ yielding a light-blue precipitate, . . . . . }		$(\text{FeCy}_6)^{\text{iv}}$ .	
On addition of $\text{Fe}_2\text{Cl}_6$ solution, a red coloration destroyed by $\text{HgCl}_2$ , . . . . . }		$(\text{CyS})$ .	
II. Portion acidified with $\text{HNO}_3$ . Refer to (444, 3).	Present.	III. Portion acidified with $\text{HA}$ .	Present.
On addition of $\text{AgNO}_3$ , a <i>pure white</i> precipitate, easily soluble in $\text{AmHO}$ (444, 1), . . . }	$\text{Cl}'$	On addition of $\text{PbA}_2$ solution a yellow precipitate, . . . }	$(\text{CrO}_4)''$
On addition of $\text{AgNO}_3$ , a <i>light-yellow</i> precipitate, with difficulty soluble in $\text{AmHO}$ , . . . }	$\text{Br}'$	On addition of $\text{CaCl}_2$ solution a white gelatinous precipitate, . . . }	$\text{F}''$ (probably).
On addition of $\text{AgNO}_3$ , a <i>yellow</i> precipitate, almost insoluble in $\text{AmHO}$ , . . . . . }	$\text{I}'$	On addition of $\text{CaCl}_2$ a white pulverulent precipitate, . . . . . }	$(\text{C}_2\text{O}_4)''$ (probably).
Refer to (445).		On addition of $\text{Fe}_2\text{Cl}_6$ a yellowish-white precipitate (449), . . . }	$(\text{PO}_4)'''$ , or $(\text{AsO}_4)'''$

## NOTES ON THE PRECEDING TABLE.

444. 1. Unless the  $\text{Na}_2\text{CO}_3$  solution used in preparing the solution for these tests was pure  $(\text{SO}_4)''$  and  $(\text{Cl})'$ , if detected, may have been present only as impurities in the  $\text{Na}_2\text{CO}_3$ ; portions of the original substance should then be tested by (446) and (447).

2. The presence of  $(\text{SiF}_6)''$  should be confirmed by other tests, such as adding  $\text{KCl}$  (300) and evolving  $\text{HF}$  by strong  $\text{H}_2\text{SO}_4$  (301); the  $\text{HF}$  is most readily evolved from the  $\text{BaSiF}_6$  precipitate (301 a).

3. Should  $\text{AgNO}_3$  yield a black precipitate, this proves the presence of a sulphide, or possibly a theiosulphate; add  $\text{HNO}_3$ , and boil; the black  $\text{Ag}_2\text{S}$  will thus be decomposed, leaving a milky liquid

\*  $\text{F}$  will be readily detected in this precipitate, or better in the original substance by (452);  $(\text{C}_2\text{O}_4)''$  by (453).

## 268 SPECIAL TESTS FOR ACID RADICLES. [445-449.

in which any other precipitate is readily seen after being coagulated by heating or shaking.

**445.** It must be remembered that  $(\text{Cy})'$ ,  $(\text{FeCy}_6)^{\text{iv}}$ ,  $(\text{Fe}_2\text{Cy}_{12})^{\text{vi}}$ , and  $(\text{CyS})$  are also precipitated by  $\text{AgNO}_3$ , and therefore if these acid radicles have been already found, a precipitate produced by  $\text{AgNO}_3$  does not prove the presence of  $(\text{Cl})'$ ,  $(\text{Br})'$ , or  $(\text{I})'$ , which acid radicles must be especially examined for, as directed below.

If chloride, bromide, and iodide have all to be tested for, a portion of the  $\text{Na}_2\text{CO}_3$  solution must be examined by (273); or the precipitate obtained by  $\text{AgNO}_3$  (see 443, II) may be tested by (273 a) for Cl, Br, I.

If only bromide and iodide have to be tested for, use par. 274.

### SPECIAL TESTS FOR ACID RADICLES TO BE MADE ON THE ORIGINAL SUBSTANCE.

**446. Sulphate.**—A portion of the original substance is boiled with  $\text{HCl}$ , the liquid decanted (or filtered if not clear), and  $\text{BaCl}_2$  added to it: a white precipitate shows *presence of  $\text{SO}_4$* .

**447. Chloride.**—A portion of the original substance is warmed with  $\text{HNO}_3$ , the solution decanted (or filtered if necessary), and  $\text{AgNO}_3$  solution added to it, a *perfectly white* precipitate, easily dissolved by warm  $\text{AmHO}$ , shows *presence of chloride*.

**448. Cyanide** if present will have been detected by its special test and by the smell of bitter almonds, which is given off by the substance, after adding  $\text{H}_2\text{SO}_4$  (411). Add to some of the substance  $\text{Am}_2\text{S}$  in excess and boil; filter from any dark-colored precipitate, boil the filtrate for a short time, acidify with  $\text{HCl}$  and add  $\text{Fe}_2\text{Cl}_6$ ; a red coloration, not destroyed by heat, but removed by addition of  $\text{HgCl}_2$ , shows *presence of Cy* [possibly as  $(\text{FeCy}_6)^{\text{iv}}$ ,  $(\text{Fe}_2\text{Cy}_{12})^{\text{vi}}$ , or  $(\text{CyS})$ , if these acid radicles have been found (443, 1)].

**449. Arsenate.**—This acid radicle cannot be present unless As was detected during the examination for metals; the presence of  $(\text{AsO}_4)'''$  is there rendered probable by the precipitation of yellow  $\text{As}_2\text{S}_3$  occurring only when the liquid saturated with  $\text{H}_2\text{S}$  is boiled. If As has been found amongst the metals, proceed to test for  $(\text{AsO}_4)'''$  by adding to a portion of the  $\text{Na}_2\text{CO}_3$  solution

(442) HCl until it is acid, then AmCl, AmHO in excess and  $\text{MgSO}_4$ ; filter off any precipitate which forms on warming and shaking the liquid, and pour a few drops of  $\text{AgNO}_3$  solution upon the white precipitate on the filter, a change of color to brown shows *presence of*  $(\text{AsO}_4)'''$ .

**450. Phosphate.**—Boil some of the original substance with dilute  $\text{HNO}_3$ , add a little of this clear solution (filtered if necessary) to some AmHMoO<sub>4</sub> solution, shake and stir the liquid well, and if no precipitate forms warm *very gently*, a yellow precipitate shows *presence of*  $(\text{PO}_4)'''$ .

*Note.*—If  $(\text{AsO}_4)'''$  has been detected (449) this test for  $(\text{PO}_4)'''$  is only reliable when the yellow precipitate has been obtained either in the cold or by employing a *very gentle* heat. If any doubt is felt concerning the presence of  $(\text{PO}_4)'''$ , boil some of the substance with strong HCl, and examine for  $(\text{PO}_4)'''$  after having entirely separated the  $(\text{AsO}_4)'''$  by passing  $\text{H}_2\text{S}$  into the boiling HCl solution. A doubtful precipitate with AmHMoO<sub>4</sub> may be proved to contain phosphate by dissolving it in AmHO, adding AmCl and  $\text{MgSO}_4$ , filtering and dropping  $\text{AgNO}_3$  upon the precipitate, if phosphate it turns *yellow*, if arsenate *brown*.

**451. Borate.**—Warm a portion of the substance with a little dilute HCl, dip into the solution a strip of turmeric-paper and dry it in a steam-oven (or at a gentle heat), if the slip is *reddish-brown* and becomes *blue-black* when moistened with AmHO, the *presence of*  $(\text{BO}_3)'''$  is shown.

**452. Fluoride.**—Pour upon a portion of the powdered substance strong  $\text{H}_2\text{SO}_4$ ; and warm the mixture in a leaden or platinum crucible covered with a watch-glass, which has been coated with wax or paraffin, and has then had characters traced through the film with the point of a penknife; the characters are etched upon the glass, showing *the presence of F.*

If  $\text{SiO}_2$  is known to be present, the test must be made by conducting the gas, evolved on heating the substance with strong  $\text{H}_2\text{SO}_4$ , into dilute AmHO, when a deposit of gelatinous  $\text{H}_4\text{SiO}_4$  proves the *presence of F.*

**453. Oxalate.**—The precipitate produced by  $\text{CaCl}_2$  in III (443), or the original substance if it contains no car-

bonate, is mixed on a watch-glass with  $\text{MnO}_2$  free from carbonate and strong  $\text{H}_2\text{SO}_4$ , and gently warmed;  $\text{CO}_2$  is evolved and detected by holding over the bubbles which rise from the mixture a rod moistened with lime-water, or by inverting another watch-glass containing on its under surface a drop of lime-water over the glass containing the mixture.

**453 a. Tartrate.**—The presence of  $\bar{\text{T}}$  will probably have been indicated in the preliminary examination (388 or 412). In order to confirm its presence, any metals of Groups I, II, and III must be separated by  $\text{H}_2\text{S}$  or  $\text{Am}_2\text{S}$  (see last two pars. of 358 b), and  $\text{Ca}\bar{\text{T}}$  is then precipitated from the liquid, made just alkaline with  $\text{AmHO}$  and mixed with a little  $\text{AmCl}$ , by adding excess of  $\text{CaCl}_2$ , shaking well and letting stand for some time. A precipitate may consist of  $\text{Ca}\bar{\text{T}}$ ,  $\text{Ca}_3(\text{PO}_4)_2$ , or  $\text{CaO}$ . By shaking this precipitate with cold  $\text{KHO}$  solution,  $\text{Ca}\bar{\text{T}}$  is dissolved and may be reprecipitated from the solution (or filtrate) by diluting and boiling for some time; the precipitate is then gently warmed, after decanting the liquid, with a little *very dilute*  $\text{AmHO}$  and a crystal of  $\text{AgNO}_3$ , a metallic mirror shows the *presence of  $\bar{\text{T}}$* .

**454. Sulphide.**—Very small quantities of S may be detected by boiling the substance with  $\text{KHO}$  for some time, and adding to the clear solution alkaline  $\text{PbA}$ , solution, when a black precipitate or coloration will appear.

---

The above directions must only be considered as affording examples of how the principal acid radicles are detected. By reference to the individual reactions in Section IV, and to the directions there given for the detection of similar acid radicles when occurring together, this method will serve for the detection of all the commonly occurring acid radicles.

455. TABLE SHOWING SOLUBILITY OF SUBSTANCES IN WATER AND ACIDS.

A blank signifies that the solubility is unknown or unimportant.

	K.	Na.	Am.	Mg.	Ba.	Sr.	Ca.	Fe <sup>+</sup> Fe <sup>+++</sup>	Al.	Cr.	Zn.	Mn.	Ni.	Co.	Hg <sup>+</sup>	Pb.	Bi.	Cu.	Cd.	Sb.	Sn <sup>+</sup> , Sn <sup>+++</sup>	As <sup>+</sup>	Ag.	Hg <sup>+</sup>	
1. Oxide.....	w.	w.	w.				w-a.	a.	a.i.						a.	a.	a.	a.	a.	a.m.	a.i.	w-a.m.	a.	a.	O.
2. Sulphide....	w.	w.	w.	w-a.	w.	w.	w-a.	a.m.			a.	a.	a.m.	a.m.	a.m.	a.p.	a.p.	a.p.	a.	a.m.	a.m.	a.p.	a.p.	a.p.	S.
3. Chloride.....	w.	w.	w.	w.	w.	w.	w.	w.	w.	a.i.	w.	w.	w.	w.	w.	w.i.	w.b.	w.	w.	w.	w.b.	w.	i.	a.i.	Cl.
4. Iodide.....	w.	w.	w.	w.	w.	w.	w.	w.			w.	w.			a.	w-a.m.					w.		i.	a.	I.
5. Sulphate....	w.	w.	w.	w.	w.	w.	w-a-m-i.	w.	w.	w-a.	w.	w.	w.	w.	w.b.	a.i.	w.b.	w.	w.	w.	w.b.		w-a.p.	w.b.a.p.	SO <sub>4</sub> .
6. Nitrate.....	w.	w.	w.	w.	w.	w.	w.	w.	w.	w.	w.	w.	w.	w.	w.b.	w.	w.b.	w.			w.		w.	w.b.	NO <sub>3</sub> .
7. Phosphate..	w.	w.	w.	a.	a.	a.	a.	a.	a.	a.	a.	a.	a.	a.	a.	a.		a.					a.	a.	PO <sub>4</sub> .
8. Carbonate..	w.	w.	w.	a.	a.	a.	a.	a.	a.		a.	a.	a.	a.	a.	a.		a.					a.	a.	CO <sub>3</sub> .
9. Borate.....	w.	w.	w.	a.	a.	a.	a.	a.	a.		a.	a.	a.	a.		a.	a.	a.	w-a.					w.	BO <sub>3</sub> .
10. Arsenite....	w.	w.	w.	a.	a.	a.	a.	a.	a.	a.	a.	a.	a.	a.	a.	a.	a.	a.		a.			a.	a.	AsO <sub>3</sub> .
11. Arsenate....	w.	w.	w.	a.	a.	a.	a.	a.	a.				a.	a.	a.	a.		a.		a.			a.	a.	AsO <sub>4</sub> .
12. Chromate...	w.	w.	w.	w.	w.	w.	a.	a.					a.	a.	a.	a.		a.		a.			a.	a.	CrO <sub>4</sub> .
13. Fluoride....	w.	w.	w.	w.	w.	w.			w.	a.	a.	w.	i.	a.	w-a.	a.i.	a.	w.		a.			a.	a.	F.
14. Oxalate.....	w.	w.	w.	a.	a.	a.	a.	a.				w-a.	a.	a.	a.	a.	a.	a.			w.		a.	a.	C <sub>2</sub> O <sub>4</sub> .

Only the most commonly occurring compounds are contained in this table, the solubility of other substances may be obtained by reference to a treatise on chemistry, or to Storer's *Dictionary of Solubilities*.



**456.** The table is thus arranged: in a horizontal line at the head are placed the more commonly occurring metals, which yield salts or basic oxides; in the vertical column on the left is a list of that portion of the names of the compounds thus formed, which corresponds to the acid radicle; the same are also placed in chemical symbols in a vertical column on the right. The solubility of a compound is denoted by letters:

- w. Signifies soluble in water.
- a. Soluble in acids; the term standing for HCl, HNO<sub>3</sub>, and aqua regia.
- a<sup>m</sup>. Soluble in muriatic (hydrochloric) acid.
- n<sup>a</sup>. Soluble in nitric acid.
- a<sup>mn</sup>. Soluble in a mixture of muriatic and nitric, but not in either separately.
- i. Insoluble in water and acids.
- w.a. Letters thus placed together with a stop between, signify that in different states the substance shows these different solubilities.
- w-a. Letters connected by a hyphen indicate that the substance is only slightly soluble in the first solvent, and may therefore partially fall under the class denoted by the second letter.
- w.b. Decomposed more or less by much water with formation of a basic salt which is not soluble in water but soluble in acid.

**457.** To find the solubility of any compound of one of the metals placed at the top of the table, glance down the vertical column which is headed by this element, the letter indicating the solubility of the compound will be found in a horizontal line with the acid radicle portion of the name. Thus to find the solubility of zinc sulphate, it is only necessary to glance down the vertical column with Zn at its head; on a horizontal line with "sulphate" (SO<sub>4</sub>) stands the letter *w*, showing that zinc sulphate is soluble in water.

**458.** The way in which this table is used after ascertaining the solubility of a substance under analysis, and detecting the metals present in it, may be explained by an example. In a substance *which was entirely soluble in water*, the metals found were K, Ba, Ag. We glance down the columns headed by these three metals, and can thus see at once which acid radicles may be present; all acid radicles might be present combined with K, since all its salts are soluble in water, but the presence of Ba in a substance soluble in water shows that SO<sub>4</sub>, PO<sub>4</sub>, CO<sub>3</sub>, BO<sub>3</sub>, AsO<sub>3</sub>, AsO<sub>4</sub>, CrO<sub>4</sub>, and F, cannot be present, since these acid radicles form compounds with Ba which are insoluble in water; Ag excludes in addition S, Cl, Br, and I; hence amongst the more commonly occurring acid radicles only NO<sub>3</sub> need be tested for.

This example shows how much we may often simplify the examination for acid radicles, when the metals present in a substance and also its solubility are known, by referring to the above table.

REMARKS ON THE PRECIPITATION OF  
GROUP III.

459. In the General Table (419) it is assumed that in the absence of  $(\text{PO}_4)'''$  the two Sub-groups III A and III B can be separated from one another by adding first  $\text{AmCl}$  and then excess of  $\text{AmHO}$  to the solution; Group III A alone being thus precipitated, and Group III B being afterwards precipitated by adding  $\text{Am}_2\text{S}$  to the filtrate. Now although  $\text{AmCl}$  entirely prevents the precipitation of Group III B by  $\text{AmHO}$ , if the solution is kept covered from the air; still if the members of Group III A are present at the same time in the solution they are precipitated by  $\text{AmHO}$ , and the presence of  $\text{AmCl}$  will not prevent Mn and Zn from being precipitated with them in smaller or larger quantity, Mn particularly showing a tendency to be precipitated with Fe, and Zn with Cr. Hence if either Mn or Zn, especially the former, be present in small quantity only, it may be entirely precipitated in Group III A; this is not a serious matter in the case of Mn, since it is readily detected in the ordinary examination of the precipitate by Table III A by yielding a *green* mass on fusion with  $\text{Na}_2\text{CO}_3$  and  $\text{KNO}_3$ ; but Zn may thus be entirely passed over, since if precipitated with  $\text{Cr}_2\text{H}_2\text{O}_4$  its presence would certainly not be detected in Table III A. The method of dissolving the precipitate several times in  $\text{HCl}$  and reprecipitating with  $\text{AmHO}$ , to some extent meets this difficulty; but it is preferable whenever a precipitate is produced on adding  $\text{AmCl}$  and  $\text{AmHO}$ , and traces of Mn and more particularly of Zn have to be tested for, to precipitate Groups III A and III B together, by adding in succession  $\text{AmCl}$ ,  $\text{AmHO}$ , and  $\text{Am}_2\text{S}$ , and boiling. The precipitate is then examined by Table III C (464, 465) for Groups III A and III B, the filtrate being examined, as directed in the General Table for Groups IV and V. It must, however, be understood that if no precipitate is produced by  $\text{AmCl}$  and  $\text{AmHO}$ ,  $\text{Am}_2\text{S}$  may be added, and the precipitate (if any) examined by Table III B, since in the absence

of Group III A the members of Group III B are not precipitated by AmHO in the presence of AmCl.

460. *The presence of  $(PO_4)'''$  in the HCl solution* which is to be examined for Groups III, IV, and V involves no special procedure if AmHO added after AmCl produces no precipitate, since the phosphates of Groups III and IV and of Mg must be absent. If, however, on adding AmCl and AmHO a precipitate is formed, the directions given at the head of Table III D (466) must be followed, and the precipitate must be examined by that table. The reason for this departure from the ordinary course of analysis is that, whereas Al, Cr, Ba, Sr, Ca, and Mg, if present as phosphates, are completely precipitated by AmHO, the phosphates of Ni, Co, Mn, Zn, and Fe, are only partially precipitated by AmHO; they are, however, entirely precipitated by  $Am_2S$ . The filtrate from  $Am_2S$  is then examined as directed in the General Table for Groups IV and V.

The principles on which the method drawn out in Table III D is founded are:

1. The insolubility of the phosphates of Al, Fe, and Cr in  $H\bar{A}$  in the presence of alkaline acetates, the other portions of the precipitate being soluble.

2. The separation of all the  $(PO_4)'''$  which is in the  $H\bar{A}$  solution combined with Ba, Sr, Ca, or Mg, by  $Fe_2Cl_6$  in an  $H\bar{A}$  solution.

The further separation of Al, Fe, and Cr phosphates is somewhat complicated by the fact that  $AlPO_4$  is only decomposed by fusion mixture when  $SiO_2$  is also present, and this  $SiO_2$  has to be removed after the fusion by methods which will be intelligible on reference to (290) and (291).

461. The reason why, in obtaining the precipitate for Table III D, the precipitates produced by AmHO and  $Am_2S$  are filtered and washed separately, is that phosphates of Fe, Zn, Mn, Ni, and Co are converted by  $Am_2S$  into sulphides, forming at the same time ammonium phosphate which remains in solution, and this would precipitate Ba, Sr, Ca, Mg as phosphates from the solu-

tion, even if they were not originally present in that condition. Now this besides complicating the process of analysis, would also render it impossible to state whether Ba, Sr, Ca, and Mg were originally present as phosphates or not. Hence the phosphates are first precipitated by  $\text{AmCl}$  and  $\text{AmHO}$ , then any members of Group III B are precipitated from the filtrate by  $\text{Am}_2\text{S}$ . The two precipitates are mixed and treated with  $\text{Am}_2\text{S}$ , which will dissolve away the  $(\text{PO}_4)'''$  from Fe, Zn, Mn, Ni, Co phosphates if present, leaving the other phosphates undecomposed; hence, if on filtering and testing the filtrate with  $\text{AmCl}$ ,  $\text{AmHO}$ , and  $\text{MgSO}_4$  a white crystalline precipitate is obtained, this indicates the presence of  $(\text{PO}_4)'''$  and indirectly establishes the presence in the original precipitate of phosphate of some one or more of the metals Fe, Zn, Mn, Ni, Co.

462. Oxalates, borates, fluorides, and silicates of Ba, Sr, Ca, Mg, would likewise be precipitated by  $\text{AmHO}$  in Group III A; but by the evaporation of the  $\text{HCl}$  solution after passing  $\text{H}_2\text{S}$ ,  $\text{H}_3\text{BO}_3$ , and  $\text{HF}$  are usually volatilized, and  $\text{H}_2\text{SiO}_3$  becomes insoluble; oxalates are also decomposed by a gentle ignition after evaporation.

\* \* The best course to be pursued in precipitating and examining Group III, will be found in par. 463.

#### 463. RULES FOR THE PRECIPITATION AND EXAMINATION OF GROUPS III A AND III B.

*Note.*—Refer to pars. 459-462 for explanation of these rules.

Hence the following rules may be laid down for precipitating and detecting members of Group III A and III B in the  $\text{HCl}$  solution, a small portion of which has been tested for  $(\text{PO}_4)'''$  by  $\text{AmHMoO}_4$ :

I. *If the addition of  $\text{AmHO}$  after  $\text{AmCl}$  causes no precipitate*,  $\text{Am}_2\text{S}$  is at once added, the liquid boiled, and the precipitate, if any, examined for members of Group

III B by Table III B (437); this course is pursued whether  $(PO_4)'''$  is present or absent.

II. *If the addition of AmHO after AmCl causes a precipitate and  $(PO_4)'''$  is not present*, the plan of precipitation depends upon the object of the analysis.

If small quantities of Zn need not be examined for, the method given in the General Table (419) may be followed.

If traces of Zn have to be tested for in the solution, add AmCl, then AmHO and Am<sub>2</sub>S in excess, boil, filter, and examine the precipitate by Table III c (464), or III c<sub>2</sub> (465).

*Note.*—The method described in Table III c<sub>1</sub> gives reliable results, and is to be used for very careful analysis; the method in Table III c<sub>2</sub> is, however, much more simple, and is sufficiently accurate for general use.

III. *If the addition of AmHO after AmCl causes a precipitate and  $(PO_4)'''$  is present*, the method of precipitation to be adopted will be found in par. 466, together with the table for the examination of the group precipitate.

#### 464. TABLE III c<sub>1</sub>.—FOR THE SEPARATION OF GROUPS III A AND III B BY BaCO<sub>3</sub>.

The precipitate may contain Fe, Al, Cr, Zn, Mn, Ni, Co. Rinse it off the filter into a porcelain dish, using as little water as possible, add some strong HCl and boil, adding at intervals a small crystal of KClO<sub>3</sub>, until all is dissolved but a small quantity of yellow sulphur. Evaporate very nearly to dryness, dilute with a little water, and pour (through a filter, if necessary) into a small flask. Cool, pour in a small quantity of BaCO<sub>3</sub> suspended in water, cork the flask tightly and shake well; repeat the addition of BaCO<sub>3</sub> and agitate until the precipitate is distinctly whitened by the excess of BaCO<sub>3</sub>; then shake well and allow the flask to stand by corked for at least fifteen minutes, occasionally shaking it vigorously. Let the precipitate subside, filter: wash

the precipitate with a little cold water allowing the washings to run through into the filtrate; then wash thoroughly, rejecting the washing water:

<p>1. <i>Precipitate</i> may contain Fe, Al, Cr, and <math>\text{BaCO}_3</math>: dissolve by pouring a little boiling <math>\text{HCl}</math> upon the filter, returning the boiling acid to the filter if necessary to dissolve any undissolved precipitate; remove Ba from the boiling solution by adding dilute <math>\text{H}_2\text{SO}_4</math> gradually, until after allowing the precipitate to settle, a few additional drops of acid cause no further precipitate (Note 1): filter, add pure <math>\text{NaHO}</math> in excess to the filtrate, boil and filter:</p>		<p>2. <i>Filtrate</i> may contain Zn, Mn, Ni, Co, and <math>\text{BaCl}_2</math>. Remove Ba by adding to the boiling liquid dilute <math>\text{H}_2\text{SO}_4</math> gradually, until the last few drops produce no further precipitate in the clear liquid from which the <math>\text{BaSO}_4</math> has been allowed to settle; filter; add pure <math>\text{NaHO}</math> in excess to the cold filtrate, stir well and filter:</p>	
<p><i>Precipitate</i>: examine for Fe and Cr by column 2, Table III A (436).</p>	<p><i>Filtrate</i>: examine for Al by column 1, Table III A (436).</p>	<p><i>Precipitate</i> may contain Mn, Ni, Co: rinse off the filter into a porcelain dish with as little water as possible, add some strong <math>\text{HCl}</math> and boil; evaporate nearly to dryness, add a little water and dissolve some solid <math>\text{NaA}</math> in the liquid, pass <math>\text{H}_2\text{S}</math> to saturation, filter:</p>	<p><i>Filtrate</i>: pass <math>\text{H}_2\text{S}</math>, a white precipitate indicates:</p> <p><i>Presence of Zn.</i></p>
<p><i>Note 1.</i>—The separation of Ba may be neglected here, the precipitate being at once boiled with <math>\text{NaHO}</math>; in this case much white <math>\text{BaCO}_3</math> will remain with the <math>\text{Fe}_2\text{O}_3</math> after fusion to separate Cr; it will however dissolve with the Fe and cause no complication.</p>		<p><i>Precipitate</i>: examine for Ni and Co according to Table III B, column 1 (437).</p>	<p><i>Filtrate</i>: add <math>\text{AmHO}</math> in excess; a flesh-colored precipitate shows:</p> <p><i>Presence of Mn.</i></p>

465. TABLE III C<sub>2</sub>.—FOR EXAMINATION OF GROUPS III A AND III B WHEN MIXED.

Remove the precipitate produced by  $\text{AmCl}$ ,  $\text{AmHO}$  and  $\text{AmS}$  from the filter by opening the filter out inside a porcelain dish, pouring in dilute  $\text{HCl}$ , and gently stirring and shaking the liquid: take out the filter-paper, stir well, and if any *black* residue is left, filter (see Note 1):

<p><i>Residue</i> If black may con- tain <math>\text{NiS}</math> and <math>\text{CoS}</math>: examine it as di- luted in Table III B, col- umn 1 (457).</p>	<p><i>Filtrate</i> may contain <math>\text{Al}</math>, <math>\text{Fe}</math>, <math>\text{Cr}</math>, <math>\text{Zn}</math>, <math>\text{Mn}</math>. Carefully note its color, which is best seen by pouring it into a white porcelain dish, and according to the result of this observation, examine the liquid by column I and II.</p>
<p>I. <i>The liquid is perfectly colorless</i>: <i>Absence of Cr</i>. Boil the liquid in the dish for a few minutes until it is no longer smells of <math>\text{H}_2\text{S}</math>, then add a small crystal <math>\text{KClO}_3</math> and boil down to a small bulk: allow to cool, add <math>\text{NaHO}</math> in excess, stir well and filter:</p>	<p><i>Filtrate</i> may contain <math>\text{Zn}</math>, <math>\text{Al}</math>. Divide into two equal parts (Note 2). Into one portion pass <math>\text{H}_2\text{S}</math>: a white precipitate forms at once: <i>Presence of Zn</i>. To the other portion add dilute <math>\text{HCl}</math> gradually whilst stirring: if no precipitate forms, add <math>\text{HCl}</math> in excess, and <math>\text{AmHO}</math> in excess, and heat. A colorless flocculent precipitate: <i>Presence of Al</i>.</p>
<p><i>Precipitate</i> may contain <math>\text{Fe}</math>, <math>\text{Mn}</math>. Dry and use it on platinum foil with <math>\text{Na}_2\text{CO}_3</math> and <math>\text{KNO}_3</math>: a blue-green mass shows: <i>Presence of Mn</i>.</p>	<p><i>Precipitate</i> may contain <math>\text{Fe}</math>, <math>\text{Al}</math>, <math>\text{Cr}</math>: examine it by column 1 Table III C<sub>1</sub> (464). <i>Filtrate</i> may contain <math>\text{Zn}</math>, <math>\text{Mn}</math>: boil and whilst boiling add dilute <math>\text{H}_2\text{SO}_4</math> gradually until the last few drops no further precipitate, filter from <math>\text{FeSO}_4</math>; cool, add pure <math>\text{NaSHO}</math> in excess, stir well and filter:</p>
<p><i>Filtrate</i> may contain <math>\text{Zn}</math>, <math>\text{Mn}</math>: boil and whilst boiling add dilute <math>\text{H}_2\text{SO}_4</math> gradually until the last few drops no further precipitate, filter from <math>\text{FeSO}_4</math>; cool, add pure <math>\text{NaSHO}</math> in excess, stir well and filter: <i>Precipitate</i>: dry and fuse with <math>\text{Na}_2\text{CO}_3</math> and <math>\text{KNO}_3</math> on platinum foil; blue-green mass: <i>Presence of Mn</i>.</p>	<p><i>Filtrate</i>: pass <math>\text{H}_2\text{S}</math>; white precipitate: <i>Presence of Zn</i>.</p>

*Note 1.*—A mere milkiness, due to the separation of sulphur, shows the absence of Ni and Co, and does not render filtration necessary.

*Note 2.*—Al and Zn may also be detected without dividing the filtrate by adding to it HCl gradually, until it becomes acid; then AmHO until alkaline, a colorless flocculent precipitate shows *Presence of Al*: this is filtered off and H<sub>2</sub>S passed into the filtrate, a white precipitate shows *Presence of Zn*.

TABLE III D FOR EXAMINATION OF PHOSPHATES IN GROUP III.

466. If (PO<sub>4</sub>)''' is found in the HCl solution in the General Table (419) after precipitating Groups I and II, and a precipitate forms on addition of AmCl and AmHO, these reagents are added, the latter in slight excess, to the HCl solution; it is then gently heated, filtered quickly, and washed well with hot water. To the filtrate Am<sub>2</sub>S is added in excess and the solution boiled, then filtered, and the filtrate examined for Groups IV and V (419) and (431). Transfer the two precipitates yielded by AmHO and Am<sub>2</sub>S to a porcelain dish, and stir well with a little Am<sub>2</sub>S, filter off and wash well. Examine the precipitate by the following table:

<i>Filtrate:</i> Add Fe <sub>2</sub> Cl <sub>6</sub> (note 2) drop by drop as long as a precipitate forms (large excess must be carefully avoided), and until the liquid, after being well stirred or shaken, remains reddish. Warm gently for some time, filter whilst hot, and wash with hot water:		
<i>Filtrate:</i> add AmCl, AmHO (note 3), and Am <sub>2</sub> S: filter:		<i>Precipitate</i> containing FePO <sub>4</sub> may be neglected.
<i>Filtrate:</i> add Am <sub>2</sub> CO <sub>3</sub> : filter:	<i>Precipitate:</i> examine by Table III c (464, 465) for Zn, Mn, Ni, Co, also for Al and Cr. Test a portion of the original solution (or solution of the substance in HCl) for Fe'' and Fe''' by note 2 (436).	
<i>Precipitate</i> may contain Ba(O <sub>2</sub> ), Sr(O <sub>2</sub> ), CaCl <sub>2</sub> . Examine this precipitate by itself by Table IV (438). Ba, Sr or Ca, if found in this precipitate, were present in the solution as phosphates.	<i>Filtrate</i> may contain Mg; add Na <sub>2</sub> HPO <sub>4</sub> , warm and shake well. White crystalline precipitate:  <i>Presence of Mg</i> (as phosphate).	

*Note*—The filtrate should be examined for (PO<sub>4</sub>)''' by adding MgSO<sub>4</sub>; if this is present Fe, Zn, Mn, Ni, Co, one or more were present as phosphates.



The precipitate produced by  $\text{AmCl}$ ,  $\text{AmHO}$ , and  $\text{Am}_2\text{S}$  may contain Ba, Sr, Ca, Mg as phosphates; Fe, Zn, Mn, Ni, Co as sulphides; and Al, Cr as hydrates. Remove it from the filter, and heat it with dilute  $\text{HCl}$  in a porcelain dish. If this does not dissolve it, drop in several small crystals of  $\text{KClO}_3$ , and evaporate very nearly to dryness. Filter off S, if necessary (note 1); then add a solution of  $\text{H}\bar{\text{A}}$  and  $\text{Na}\bar{\text{A}}$  (No. 84) (526), as long as any precipitate is produced. Heat gently, and filter while hot:

<p><i>Precipitate</i> may consist of <math>\text{FePO}_4</math>, <math>\text{AlPO}_4</math>, <math>\text{CrPO}_4</math> (note 4). Dry the precipitate on the filter, and fuse it for a short time with finely powdered <math>\text{SiO}_2</math> and fusion-mixture, adding a little <math>\text{KNO}_3</math> in a platinum crucible or on platinum foil. When cold, dissolve by boiling with a little distilled water, add <math>\text{Am}_2\text{CO}_3</math>, allow to stand for a short time, stirring occasionally; let the precipitate subside, filter:</p>		
<p><i>Filtrate</i>: add <math>\text{H}\bar{\text{A}}</math> until the solution is acid, boil for a short time, then add <math>\text{Pb}\bar{\text{A}}</math>: yellow precipitate of <math>\text{PbCrO}_4</math>.  <i>Presence of Cr</i>, (as phosphate).  <i>Note</i>.—A white precipitate on addition of <math>\text{Pb}\bar{\text{A}}</math> may be disregarded.</p>	<p><i>Precipitate</i>: may contain <math>\text{H}_2\text{SiO}_4</math>, Fe and Al as silicates, and <math>\text{Fe}_2\text{H}_2\text{O}_6</math>. Acidify with <math>\text{HCl}</math>, evaporate to dryness, and heat the dry residue gently. Warm with a few drops of strong <math>\text{HCl}</math>, add hot water, and filter:</p>	
	<p><i>Filtrate</i>: add pure <math>\text{NaHO}</math> in excess, boil and filter:</p>	<p><i>Residue</i> is <math>\text{SiO}_2</math> and may be neglected.</p>
	<p><i>Filtrate</i>: acidify with <math>\text{HCl}</math>, and add <math>\text{AmHO}</math> in excess, boil: white, gelatinous precipitate:  <i>Presence of Al</i>, (as phosphate).</p>	<p><i>Precipitate</i> is brown <math>\text{Fe}_2\text{H}_2\text{O}_6</math>: dissolve by boiling with <math>\text{HCl}</math>, and add <math>\text{KCys}</math>: blood-red coloration:  <i>Presence of Fe</i>, (as phosphate).</p>

*Note 1*.—Add several drops of this  $\text{HCl}$  solution to some  $\text{AmHMoO}_4$  solution in a test-tube and warm gently; if no yellow precipitate forms, even after standing for a time ( $\text{PO}_4'''$  is absent; this proves that only Ni, Co, Mn, Zn, and Fe can be present in the solution, which may therefore be at once examined by Table III c (464, 465), or as directed in Note 3 below.

*Note 2*.—A few drops of  $\text{Fe}_2\text{Cl}_6$  should be added to a small part only of the filtrate and heated; if it causes no precipitate ( $\text{PO}_4'''$ , and therefore Ba, Sr, Ca, and Mg, are absent, and to the other portion  $\text{AmCl}$ ,  $\text{AmHO}$ , and  $\text{Am}_2\text{S}$  may be at once added, and the precipitate examined by Table III c (464, 465), or the solution may be examined as directed in Note 3.

*Note 3*.—Since this filtrate has to be tested for Groups III A and III B, it may be examined either by adding  $\text{AmCl}$ ,  $\text{AmHO}$ , and  $\text{Am}_2\text{S}$  together, and analyzing the precipitate by Table

III c, or by precipitating Groups III A and III B, separately by  $\text{AmCl}$  and  $\text{AmHO}$ , and then  $\text{Am}_2\text{S}$ , and examining the precipitates by Tables III A and III B (436, 437).

127.54  
96

*Note 4*—Since  $\text{CrPO}_4$  is rarely present, this precipitate may generally be tested by boiling with excess of  $\text{NaHO}$ , and filtering;  $\text{AlPO}_4$  will be in solution, and is detected by acidifying with  $\text{HCl}$  and adding  $\text{AmHO}$  in excess, which gives a gelatinous precipitate;  $\text{Fe}_2\text{H}_6$  remains as a precipitate, and is detected by dissolving in  $\text{HCl}$  and adding  $\text{KC}_2\text{S}$ .

#### Bb. THE SUBSTANCE POSSESSES METALLIC LUSTRE, AND IS PROBABLY A METAL OR ALLOY.

After making the Preliminary Examination in (467–469), either of the two methods given below (470, 471) may be adopted.

The first method is by far the most ready and certain for the examination of a metal or alloy of unknown composition, since all metals, even when present in small quantity only, may thus be detected.

In making the solution by the second method,  $\text{Sn}$ ,  $\text{Sb}$ ,  $\text{Au}$ , and  $\text{Pt}$  are left undissolved, all other metals passing into solution; it is not, however, to be recommended as a *general* process for examination of metals and alloys, since the following complications are apt to occur: If  $\text{As}$  is present with  $\text{Sn}$ , a part or the whole of the  $\text{As}$  may remain in the residue;  $\text{Pt}$ , if present with a sufficient quantity of  $\text{Ag}$ , may pass entirely into solution; and  $\text{Sb}$  will always partly dissolve. The process, however, is useful in cases where an alloy, known to contain a large proportion of  $\text{Sn}$  or  $\text{Sb}$ , has to be examined for other metals, which are thus dissolved away at once from the bulk of the  $\text{Sn}$  or  $\text{Sb}$ : but care must be taken that  $\text{As}$  is not passed over in the presence of  $\text{Sn}$ .

#### PRELIMINARY EXAMINATION.

**467.** Note its color, and smell, if any, when rubbed with the hand; also whether it is crystalline or not. Note also the hardness of the metal, by seeing if it can be scratched or cut by a steel knife; also whether on being struck smartly with a hammer it breaks to powder

(brittle), or flattens out (malleable). Then try the following experiments :

Experiment.	Observation.	Inference.
468. I.—Heat a small portion on charcoal in the inner blowpipe flame.	The results obtained by (403) and (404) in the Preliminary Table may be here obtained.	The inference will be found stated in (403) and (404).
469. II.—Heat a portion in a small tube closed at one end.	A metallic sublimate forms. It consists of small liquid globules.	Presence of Hg, Cd, As, two latter detected by (468).
<i>Note.</i> —As and S can frequently only be detected as arsenate and sulphate by fusion in the outer flame with $\text{KNO}_3$ .	A yellow sublimate melting to reddish-yellow drops, and the substance heated <i>slowly</i> in a tube open at both ends evolves $\text{SO}_2$ . [See (368), III. (417).]	Presence of Hg.  Presence of S from a sulphide.

#### PROCESSES FOR SOLUTION AND EXAMINATION OF A METAL OR ALLOY.

**470. METHOD I.**—Pour some rather dilute HCl upon the powdered metal in a small flask covered with a watch-glass with its concave surface upwards, and heat for some time just short of boiling; if the metal dissolves readily, continue heating until the metal is completely dissolved, and examine the HCl solution according to the General Table (419).

Frequently HCl alone does not effect complete solution, two or three drops of strong  $\text{HNO}_3$  should then be poured in, and more  $\text{HNO}_3$  and HCl added occasionally when the action ceases, or when red fumes are no longer given off. When the metal has entirely disappeared, add a little more strong HCl, and boil as long as any Cl or reddish fumes are given off; then dilute with a little water, heat to boiling, and cool; filter if there is any white residue :

*Residue*, if crystalline, is probably  $\text{PbCl}_2$ , and will be found to dissolve entirely if washed with sufficient boiling water.

In this solution the presence of Pb is confirmed by adding  $\text{K}_2\text{CrO}_4$ , which gives a yellow precipitate soluble in excess of  $\text{KHO}$ :

*Presence of Pb.*

If any residue is left after washing with boiling water, it is probably  $\text{AgCl}$ ; confirm the presence of Ag by pouring upon the residue hot  $\text{AmHO}$ , it dissolves completely, but is reprecipitated on adding  $\text{HNO}_3$  in excess:

*Presence of Ag.*

*Filtrate* (note 1): dilute largely with  $\text{H}_2\text{O}$  (see note 2), and whether this causes a precipitate or not, pass  $\text{H}_2\text{S}$  to saturation into the solution; examine any precipitate thus produced by Table II (435), and proceed to examine the filtrate for Groups III, IV, and V, as directed in the General Table (419).

*Note 1.*—If Au and Pt may be present, they must be tested for in the  $\text{H}_2\text{S}$  precipitate, according to the directions in (472–474).

*Note 2.*—A white precipitate appearing on dilution, is due to the presence of Bi, Sb, or Sn.

**471. METHOD II.**—Pour upon the finely divided metal some strong  $\text{HNO}_3$ ,\* and heat in a small flask covered with a watch-glass as long as any red fumes appear. One of two results will occur:

I.	II.		
<p><i>The substance dissolves completely with or without addition of water.</i> Absence of Pt, Au, Sb, Sn (note 1). Examine the solution, after boiling nearly to dryness, and diluting with water (note 2), by the General Table (419).</p>	<p><i>A Residue is left:</i> add some hot water and boil, then filter and wash the residue on the filter well with boiling water (note 2). The residue may present the following appearances:</p> <table border="1" data-bbox="386 1036 831 1294"> <tr> <td data-bbox="386 1036 683 1294"> <p>1. <i>Entirely metallic or black powder:</i> probably Pt or Au. Dissolve by heating in a small flask, covered with a watch-glass, with a little <math>\text{HCl}</math> to which a few drops of <math>\text{HNO}_3</math> have been added. When completely dissolved add more <math>\text{HCl}</math>, and boil down in an evaporating basin nearly to dryness, examine the solution for Au and Pt by (474) commencing with the addition of <math>\text{KCl}</math>, and using only the left-hand side of the table, since Sn will be absent.</p> </td><td data-bbox="683 1036 831 1294"> <p>2. <i>White powder</i> may contain Sn, Sb, As, possibly also Pt and Au concealed in it (note 3).</p> </td></tr> </table>	<p>1. <i>Entirely metallic or black powder:</i> probably Pt or Au. Dissolve by heating in a small flask, covered with a watch-glass, with a little <math>\text{HCl}</math> to which a few drops of <math>\text{HNO}_3</math> have been added. When completely dissolved add more <math>\text{HCl}</math>, and boil down in an evaporating basin nearly to dryness, examine the solution for Au and Pt by (474) commencing with the addition of <math>\text{KCl}</math>, and using only the left-hand side of the table, since Sn will be absent.</p>	<p>2. <i>White powder</i> may contain Sn, Sb, As, possibly also Pt and Au concealed in it (note 3).</p>
<p>1. <i>Entirely metallic or black powder:</i> probably Pt or Au. Dissolve by heating in a small flask, covered with a watch-glass, with a little <math>\text{HCl}</math> to which a few drops of <math>\text{HNO}_3</math> have been added. When completely dissolved add more <math>\text{HCl}</math>, and boil down in an evaporating basin nearly to dryness, examine the solution for Au and Pt by (474) commencing with the addition of <math>\text{KCl}</math>, and using only the left-hand side of the table, since Sn will be absent.</p>	<p>2. <i>White powder</i> may contain Sn, Sb, As, possibly also Pt and Au concealed in it (note 3).</p>		

\* If Hg is found in the preliminary examination and small quantities of Ag have to be tested for, the Hg should be expelled by heating the substance strongly in a porcelain crucible before dissolving in acid, as  $\text{Hg}(\text{NO}_3)_2$  tends to prevent the precipitation of Ag by  $\text{HCl}$  in Group I.

*Note 1.* The solution is liable to contain small quantities of these metals, which must always be tested for in a careful analysis.

*Note 2.*— $\text{BiOCl}$  will often precipitate on dilution, but the precipitate will disappear on adding  $\text{HCl}$  and boiling, being thus easily distinguished from the Group I precipitate.

*Note 3.*—If much residue is obtained a small quantity of it may be heated in a test-tube with  $\text{HCl}$  adding  $\text{KClO}_3$ , if it dissolves entirely dissolve the whole of the residue, then examine the solution by Table II, commencing at (435 b): if it refuses to dissolve, proceed with the rest of the residue as directed below. If the residue is small in quantity, examine it at once as directed below.

*Examination of the Non-metallic Residue insoluble in  $\text{HNO}_3$ .*—Dry the residue on the filter at a gentle heat, mix it thoroughly with about an equal quantity of powdered  $\text{Na}_2\text{CO}_3$  and  $\text{NaNO}_3$ , and fuse the mixture in a porcelain crucible for some time; then pour the melted substance out into a porcelain dish; allow it to cool, and pour upon it cold water; let it stand for some time. After crushing the mass with a pestle and stirring it occasionally, filter, and wash the residue on the filter with dilute alcohol, throwing away the washings:

*Residue* may contain Sn, Sb, Pt, Au. Place it in a small porcelain dish, pour in a little  $\text{HCl}$  and heat, then add water; whether the residue has dissolved or not place in the liquid a slip of Pt foil and drop upon it a piece of pure Zn; H gas is given off; wait until this ceases, taking care that there is some Zn left undissolved, and adding more Zn if the first piece has entirely dissolved. Now remove the platinum strip; if it is stained black *Sb is present*.

The residue in the dish may consist of Sn, Au, and Pt; remove the Zn, rinsing off any substance adhering to it into the dish, stir the liquid in the dish well, then pour off the liquid, carefully leaving the residue; pour in water, stir well and again pour off carefully. Boil the residue for some time with strong  $\text{HCl}$  in a test-tube, dilute, decant, add  $\text{HgCl}_2$ , a white precipitate forms:—*Presence of Sn.*

*Residue:* dissolve by warming with  $\text{HCl}$  and  $\text{HNO}_3$ , and examine the solution for Au and Pt by (474), using only the left-hand portion.

*Filtrate* may contain  $(\text{AsO}_4)'''$ : add  $\text{HNO}_3$  until the solution is acid and boil, evaporating the liquid in a dish if very bulky. Pour in  $\text{AgNO}_3$  as long as it gives any precipitate, and add gradually  $\text{AmHO}$  diluted with 10 or 12 times its bulk of water, a brown precipitate shows:

*Presence of As.*

## SEPARATION AND DETECTION OF Au AND Pt.

**472.** Au and Pt will, in the ordinary course of analysis, be entirely precipitated as sulphides in the Second Group, if  $H_2S$  is passed for some time into the hot  $HCl$  solution; and since these sulphides are soluble in  $KHO$  and in  $Am_2S$ , on examining the  $H_2S$  precipitate by Table II (435), the Au and Pt will pass into the filtrate when the precipitate is boiled with  $KHO$  or  $Am_2S$ . No modification of the process described in Table II is required for the detection of these two metals until the examination of the residue in the hydrogen flask is commenced. The Au and Pt will be present in this residue, associated with Sn if it be present, and usually also with at least a part of the Sb (if present), since Zn and Pt if in contact in an acid liquid cause the separation of metallic Sb (193). Accordingly when Au and Pt have to be tested for, the ordinary course of analysis is to be followed until the residue in the hydrogen flask is obtained; and this is examined as directed below (473).

**473.** After washing the residue left in the hydrogen flask by decantation in a porcelain dish and removing any excess of Zn, boil it with a little strong  $HCl$  in a test-tube for several minutes, allow the residue to settle, and decant the liquid:

<p><b>474. Residues.*</b> Pour upon the residue in the dish a little <math>HCl</math>, and several drops of <math>HNO_3</math>, and boil gently very nearly to dryness; add some <math>KCl</math> solution, and evaporate once more very nearly to dryness. Pour some absolute alcohol into the cool dish and stir well for a time, allow the precipitate to settle and decant the liquid, wash the precipitate by stirring it with a little more alcohol, and decant the liquid when the precipitate has settled:</p>	<p><b>Solution:</b> Add <math>HgCl_2</math>, a white precipitate, which does not appear at once if only traces of Sn are present:</p>
<p>Precipitate will consist of yellow <math>K_2PtCl_6</math> and excess of <math>KCl</math>, dissolve it in a little boiling water, pour it into a white dish, add several drops of <math>HCl</math>, then <math>SnCl_2</math>: an orange-red coloration confirms the <i>Presence of Pt.</i></p>	<p><i>Presence of Sn.</i></p>

\* If Sb has not been detected already in the precipitate produced by the gases in  $AgNO_3$  solution (435 b), this residue should be examined for Sb. Boil it once more for some time with strong  $HCl$  to remove all Sn, wash the residue well by decantation and boil it in the dish with  $H_2T$  adding a few drops of  $HNO_3$ , decant and test the liquid for Sb by adding  $HCl$  and passing  $H_2S$ : the residue is then examined as above (474).

### EXAMINATION OF SUBSTANCES INSOLUBLE IN WATER AND ACIDS.

**475.** *An insoluble substance may consist of one or more of the following substances, all of which are white except  $\text{Fe}_2\text{O}_3$ ,  $\text{Cr}_2\text{O}_3$ ,  $\text{FeCr}_2\text{O}_4$ ,  $\text{PbCrO}_4$ , S, C, and  $\text{AgCl}$  which has been exposed to light.*

*Note.*—Substances embraced in brackets ( ) may possibly be present, those in square brackets [ ] *improbably*, because they are soluble in water.

1. $\text{BaSO}_4$ ,	Insoluble in water and acids.
2. $\text{SrSO}_4$ ,	Insoluble “ “
3. $[\text{CaSO}_4]$ , }	{ Not perfectly insoluble in water, soluble in hot HCl : hence it should pass into the acid solution.
4. $(\text{PbSO}_4)$ , }	
5. $\text{PbCrO}_4$ ,	Insoluble after being strongly heated.
6. $[\text{PbCl}_2]$ ,	{ Soluble in boiling water, and should therefore have been removed if the residue was well washed with boiling water.
7. $\text{AgCl}$ ,	
	{ This may have been originally present as such, or may have been derived from the use of HCl in making the solution, or by the action of aqua regia on the insoluble substances $\text{AgBr}$ , $\text{AgI}$ , $\text{AgCy}$ , $\text{Ag}_6\text{Fe}_2\text{Cy}_{12}$ , $\text{Ag}_4\text{FeCy}_8$ .
8. $\text{SiO}_2$ ,	
9. $(\text{Al}_2\text{O}_3)$ , }	Either uncombined, or as a silicate.
10. $(\text{Fe}_2\text{O}_3)$ , }	
11. $(\text{Cr}_2\text{O}_3)$ , }	Insoluble after being strongly ignited, but these are usually dissolved by long boiling with strong HCl.
12. $\text{FeCr}_2\text{O}_4$ ,	
13. $(\text{SnO}_2)$ ,	Chrome iron ore, native.
14. $\text{Sb}_2\text{O}_3, \text{Sb}_2\text{O}_4$	Native or ignited.
15. $\text{CaF}_2$ ,	{ Also a few other fluorides, and some metaphosphates and arsenates.
16. S,	
17. C,	
	{ Yellow, slowly soluble in strong $\text{HNO}_3$ , giving red fumes, and yielding $\text{H}_2\text{SO}_4$ . Black, and quite insoluble.

*Note.*—If sufficient of the substance is at disposal, the preliminary examination (476) may be made on a portion of it. In case the quantity of substance is small, however, the whole of the substance must be employed for the examination by fusion (477).

### 476. PRELIMINARY EXAMINATION.

*Note.*—The substance must be in the state of dry powder. Make

a careful examination of it with a pocket lens. Expts. I and II need only be made if the substance is light in color.

Experiment.	Observation.	Inference.
I. Observe whether the substance darkens when allowed to stand in the light for some time.	The color changes to violet or black.	Presence of AgCl.
II. Pour a little $\text{Am}_2\text{S}$ upon a portion of the substance on a watch-glass.	The substance blackens; pass on to III. The substance does not blacken; pass on to V, omitting III and IV, since Pb and Ag must be absent. Brownish precipitate.	Presence of Pb or Ag.
III. Heat some of the substance with a little water and a small piece of KCy, filter off (keeping the residue), to the filtrate add $\text{Am}_2\text{S}$ .		Presence of AgCl. <i>Confirmatory.</i> By warming the substance with $\text{AmHO}$ , filtering, and adding excess of $\text{HNO}_3$ to the filtrate, a white precipitate will form, which, on being shaken well or heated, coagulates into flocks.
IV. Wash the residue from Exp. III well upon the filter: a. <i>It is white</i> ; drop $\text{Am}_2\text{S}$ upon it. b. <i>It is dark-colored</i> ; pour in a little $\text{H}_2\text{A}$ , and boil after adding $\text{AmHO}$ in excess; filter, to the filtrate add $\text{H}_2\text{A}$ in excess and $\text{K}_2\text{CrO}_4$ .	The residue blackens.	Presence of $\text{PbSO}_4$ (or $\text{PbCl}_2$ ).
V. Take up some of the substance upon a moistened loop of platinum wire; heat it for a short time in the inner blow-pipe flame, then moisten with a drop of strong $\text{HCl}$ , and hold in the outer part of a Bunsen flame.	Yellow precipitate, soluble in $\text{KHO}$ . A <i>reddish-yellow</i> coloration, <i>dusky-green</i> when seen through the indigo-prism. A <i>crimson-red</i> coloration appearing <i>deep-red</i> through the indigo-prism. A <i>yellowish-green</i> coloration. [ <i>Note.</i> —If all these are present the colorations may often be seen in succession, Ca first, then Sr, and Ba last and very permanent.] Yellow sublimate forms on the sides of tube. When strongly heated the substance smoulders, and ultimately burns away.	Presence of $\text{PbSO}_4$ (or $\text{PbCl}_2$ ).  Presence of Ca Presence of Sr Presence of Ba } as sulphate.
VI. Heat in a small test-tube or ignition-tube, then on a piece of porcelain or broken glass, or platinum foil.		Presence of S.
VII. Heat with strong $\text{H}_2\text{SO}_4$ in a platinum crucible or leaden cup covered with a watch-glass (296). Or if a silicate is present examine by passing the gas into $\text{AmHO}$ (297).	The glass is etched.	Presence of C.
VIII. Fuse some of the substance in a bead of $\text{NaAmHPO}_4$ first in the outer then in the inner blow-pipe flame.	Gelatinous precipitate obtained in the $\text{AmHO}$ .  Particles are seen floating undissolved in the melted bead. <i>Green-colored</i> bead. <i>Reddish-brown</i> bead colorless when cold, and becoming <i>greenish</i> in the inner flame.	Presence of F. Presence of F.  Presence of $\text{SiO}_2$ . Presence of Cr.  Presence of Fe.



## ACTUAL EXAMINATION OF INSOLUBLE SUBSTANCES.

**477.** Mix the finely powdered substance\* with three or four times its weight of fusion-mixture, and heat the mixture in a small covered porcelain crucible,† until it melts; keep it in fusion for at least ten minutes. Allow the crucible to cool, then pour some water into it, and allow to stand or boil until the solid mass is loosened from the crucible; boil this in a porcelain dish with distilled water, crushing the mass by pressure with a pestle, if it does not quickly fall to pieces. Allow the residue to settle, pour off the solution through a filter, and boil the residue with a little more water; pour off through the same filter, adding this filtrate to the former one.

For the examination of the undissolved residue see (478), for the filtrate (479).

**478. *Residue on the filter:*** wash well with boiling water, then make a hole in the bottom of the filter, wash the residue through into a test-tube by pouring upon it a little boiling dilute  $\text{HNO}_3$  (see Note 1, foot of table), and boil; the residue dissolves entirely if the fusion has been continued sufficiently long (see Note 2). Filter if necessary, and add to this filtrate or solution  $\text{HCl}$ , filter:

---

\* Which, if it is found by Exp. VI, to contain free S or C, must first be heated strongly for some time in a porcelain crucible.

† If Pb and Ag have been proved to be absent by the preliminary tests, or if they have been first removed by boiling the substance with  $\text{KCy}$  solution, then with  $\text{H}\bar{\text{A}}$  and excess of  $\text{AmHO}$  and washing well, a platinum crucible may be used. The use of a platinum crucible is preferable, since after fusing in porcelain small quantities of  $\text{Al}_2\text{O}_3$  and  $\text{SiO}_2$  will always be found, being derived from the action of the  $\text{Na}_2\text{CO}_3$  on the porcelain.

<p><i>Precipitate</i> may contain <math>\text{AgCl}</math>, <math>\text{PbCl}_2</math>. Examine by Table I (433).</p>	<p><i>Filtrate</i>, which must give no more precipitate with <math>\text{HCl}</math>, is boiled in a porcelain dish as long as any <math>\text{Cl}</math> is given off, diluted with cold water, saturated with <math>\text{H}_2\text{S}</math>, and filtered:</p>		
<p><i>Precipitate</i> may contain <math>\text{PbS}</math>, <math>\text{SnS}_2</math>, <math>\text{Sb}_2\text{S}_3</math>. Examine by Table II (435).</p>	<p><i>Filtrate</i>, which must give no further precipitate when <math>\text{H}_2\text{S}</math> is again passed through it, is boiled in a porcelain dish until it no longer smells of <math>\text{H}_2\text{S}</math>; a little strong <math>\text{HNO}_3</math> is then added, and the liquid boiled again for several minutes (note 3); <math>\text{AmCl}</math> is then added and the liquid boiled; then <math>\text{AmHO}</math> is added in excess:</p>		
<p><i>Precipitate</i> may contain <math>\text{Fe}_2\text{H}_2\text{O}_4</math>, <math>\text{Al}_2\text{H}_2\text{O}_4</math>, <math>\text{Cr}_2\text{H}_2\text{O}_4</math>. Examine by Table III A (436).</p>	<p>To filtrate, add <math>\text{Am}_2\text{S}</math>: [this will rarely cause any precipitate, but a precipitate here is to be examined by Table III B (437)]; then to the solution or filtrate add <math>\text{Am}_2\text{CO}_3</math>; filter:</p>		
	<table border="0"> <tr> <td data-bbox="505 657 657 761"> <p><i>Precipitate</i> may contain <math>\text{BaCO}_3</math>, <math>\text{SrCO}_3</math>, <math>\text{CaCO}_3</math>. Examine by Table IV (438).</p> </td><td data-bbox="657 657 840 761"> <p><i>Filtrate</i> may be examined for <math>\text{Mg}</math> according to Table V (439), but this is only necessary if <math>\text{SiO}_2</math> is present.</p> </td></tr> </table>	<p><i>Precipitate</i> may contain <math>\text{BaCO}_3</math>, <math>\text{SrCO}_3</math>, <math>\text{CaCO}_3</math>. Examine by Table IV (438).</p>	<p><i>Filtrate</i> may be examined for <math>\text{Mg}</math> according to Table V (439), but this is only necessary if <math>\text{SiO}_2</math> is present.</p>
<p><i>Precipitate</i> may contain <math>\text{BaCO}_3</math>, <math>\text{SrCO}_3</math>, <math>\text{CaCO}_3</math>. Examine by Table IV (438).</p>	<p><i>Filtrate</i> may be examined for <math>\text{Mg}</math> according to Table V (439), but this is only necessary if <math>\text{SiO}_2</math> is present.</p>		

*Note 1.*—If  $\text{Ag}$  and  $\text{Pb}$  are known to be absent by the preliminary experiments,  $\text{HCl}$  may be used for dissolving the residue and is preferable; if effervescence is produced by these acids the presence of an alkaline earth is proved.

*Note 2.*—A residue here may consist of chrome iron ore; this may be dissolved by heating with  $\text{HNO}_3$  and  $\text{KClO}_4$ , and the solution tested by the General Table, when  $\text{Cr}$  and  $\text{Fe}$  will be found.

*Note 3.*—It is best to evaporate this solution quite to dryness, using a water-bath towards the end if the substance spirits. This is always necessary if  $\text{SiO}_2$  is present in the substance, since it is sometimes obtained in the acid solution, and is by this means at once separated.

**479. Aqueous Solution of the Fused Mass.**—Divide into two parts, A and B.

A. Add  $\text{HCl}$  until the liquid is distinctly acid, and evaporate in a porcelain dish to dryness, finishing the process on a water-bath if necessary to avoid spirting; then continue to heat the dish gently until the residue is quite dry; pour in a little strong  $\text{HCl}$  and warm, dilute and heat again; an insoluble residue shows presence of  $\text{SiO}_2$ .

Filter, and examine the filtrate by the General Table (419) more particularly for  $\text{Al}$ , but other bases should

also be tested for, which being soluble in alkalis pass into this solution, such as Cr, Mn, Zn, Sn.

B. Test separate portions for the following acid radicles as described below:

1.  $(Cl)'$ : Acidify with  $HNO_3$  and add  $AgNO_3$ , white precipitate easily soluble in  $AmHO$ .
2.  $(SO_4)'''$ : Acidify with  $HCl$  and add  $BaCl_2$ , white precipitate insoluble on boiling.

*Note.*—Unless the fusion mixture employed was free from  $NaCl$  and  $Na_2SO_4$ , tests 1 or 2 are worthless.

3.  $(CrO_4)''$ : Acidify with  $H\bar{A}$  and add  $Pb\bar{A}_2$ , yellow precipitate; a chromate is seen also by the yellow color of the solution. Cr thus detected may have been present acting as a metal or as a constituent of an acid radicle.
4.  $(PO_4)'''$ : Acidify with  $HNO_3$ , add a few drops to  $AmHMoO_4$  solution, and warm gently. A yellow precipitate forms, often only after a time, or on gently heating. If As has been found on passing  $H_2S$  into the hot  $HCl$  solution  $(PO_4)'''$  must be tested for in a portion of the filtrate after boiling off  $H_2S$ , else the above yellow precipitate may be due to  $(AsO_4)'''$ .
5.  $(F)$ : Add  $HCl$  in excess to a part of the solution, stir well, and let stand until the  $CO_2$  has escaped, then add  $AmHO$  in excess, then  $CaCl_2$ , as long as it produces any precipitate, and let stand for a time; filter off, dry the precipitate and examine it for F by pouring strong  $H_2SO_4$  upon it in a platinum crucible covered with a watch-glass (296).

#### ANALYSIS OF SILICATES.

**480.** The presence of silica is shown by Exp. VIII in the preliminary examination of insoluble substances (476); when it has been found it becomes necessary to examine for all metals, since many silicates which are soluble when alone, become insoluble when mixed or combined with insoluble silicates.

The ordinary analytical course requires but slight modification to adapt it to the examination of silicates. Many silicates are entirely decomposed by heating with strong HCl for some time just short of boiling; if this decomposition has been effected, only a colorless residue of silicic acid will remain, which when filtered off is completely dissolved by heating it with  $\text{Na}_2\text{CO}_3$  solution.

If the silicate is not completely decomposed by hot strong HCl, some of it must be fused with fusion mixture as directed in (477), and the cool residue boiled with water; the solution and residue are then examined by (478) and (479), remembering however that all metals may be present.

481. Since Na and K cannot be tested for in the solution obtained after fusion, a separate portion must be examined for K and Na by one of the following processes:

I. Evaporate the powdered substance several times in a platinum dish or crucible either with hydrofluoric acid and subsequently with strong  $\text{H}_2\text{SO}_4$ , or with five times its weight of calcium fluoride mixed into a paste with strong  $\text{H}_2\text{SO}_4$ , in either case heating finally until no more white fumes are evolved. Boil the cool residue with water, add  $\text{BaCl}_2$  solution as long as it causes any precipitate, then add  $\text{AmHO}$  in excess and  $\text{Am}_2\text{CO}_3$  as long as it causes any precipitate, filter and examine the filtrate for K and Na as directed under Table V (439).

II. Mix the powdered substance intimately with four times its weight of barium hydrate, and heat strongly in a platinum crucible for about half an hour; dissolve the cool mass in dilute HCl, add  $\text{AmHO}$  in excess and then  $\text{Am}_2\text{CO}_3$  in excess, filter, rejecting the precipitate; evaporate the filtrate to dryness, ignite the residue, then boil it with water; add a little HCl to the solution and test for K and Na as under Table V (439).

---

#### 482. ANALYSIS OF SUBSTANCES CONTAINING CYANOGEN.

If a small quantity of the substance to be analyzed, when examined by (304) or (306), is found to contain

cyanogen, the usual course of analysis must be somewhat modified, since the presence of cyanogen would produce confusing results. The cyanogen may be present as a cyanide, sulphocyanide, ferrocyanide, ferricyanide, or cobalticyanide, rarely as a manganocyanide or chromicyanide. Since the method to be adopted when a cyanide or sulphocyanide only is present, is much more simple than that rendered necessary by the presence of the other cyanogen compounds above mentioned, it is advisable first to ascertain in what form the cyanogen occurs, by the following preliminary experiments made on a small portion of the substance; according to the results yielded by these experiments the further examination of the substance is made by method I (484), or by Method II (485).

#### PRELIMINARY EXAMINATION.

**483.** Boil a portion of the substance for several minutes with  $\text{KHO}$  solution, then add some  $\text{Na}_2\text{CO}_3$  solution as long as it causes any precipitate, and boil again for several minutes; filter, make the cold filtrate just acid with  $\text{HCl}$ , filter if necessary, and to separate portions add:

Reagent added.	Result.	Inference.
1. $\text{FeSO}_4$ solution . .	{ Blue precipitate . White precipitate.	Presence of $(\text{FeCy}_6)^{\text{iv}}$ $(\text{FeCy}_6)^{\text{vi}}$ . Probable presence of $(\text{CoCy}_6)^{\text{vi}}$ .
2. $\text{Fe}_2\text{Cl}_6$ solution . .	{ Blue precipitate . Blood-red coloration	Presence of $(\text{FeCy}_6)^{\text{iv}}$ . Presence of $(\text{CyS})$ .
3. $\text{ZnSO}_4$ solution as long as it causes any precipitate. <i>Confy.</i> —Filter off the precipitate, add $\text{ZnSO}_4$ solution to the filtrate, and if it causes any precipitate, pour again through the filter, repeating this process until no further precipitate is produced by $\text{ZnSO}_4$ . Fuse some of the precipitate in a clear borax bead . .	{ Light-brown precipitate . White precipitate .  A blue bead is produced }	{ Presence of $(\text{FeCy}_6)^{\text{vi}}$ . Presence of $(\text{FeCy}_6)^{\text{iv}}$ $(\text{CoCy}_6)^{\text{vi}}$ .  Presence of $(\text{CoCy}_6)^{\text{vi}}$ .

If these preliminary tests show the presence of  $(\text{FeCy}_6)^{\text{iv}}$ ,  $(\text{FeCy}_6)^{\text{v}}$ ,  $(\text{CoCy}_6)^{\text{vi}}$ , one or more, examine the remainder of the substance by Method II (485). If the absence of these acid radicles is indicated by not obtaining the results in the 2d column which indicate their presence proceed according to Method I (484).

If none of the above acid radicles are found in the preliminary experiments, the only modification requisite in the ordinary analytical course is to expel  $\text{HCy}$  by boiling after adding  $\text{HCl}$  in preparing the solution (383), or in testing for Group I (419).

**484. METHOD I.**—If  $(\text{CyS})'$  is present, boil the portion of the substance which is to be examined for metals with strong  $\text{HNO}_3$  for some time, dilute a little and boil again. If any residue remains decant the solution and proceed to endeavor to dissolve the residue as directed at the end of (420), any undissolved residue being then examined according to (475 *et seq.*). The  $\text{HNO}_3$  solution is evaporated nearly to dryness and examined by (419).

Another portion of the substance is examined for acid radicles in the usual way.

**485. METHOD II.**—(Fresenius.) Boil the substance with water, filter, and wash the residue with boiling water, adding the washings to the filtrate:

<p><b>Filtrate:</b> Examine this for metals (more especially the alkalies) and for acid radicals according to the directions given for analyzing a liquid (380). The acid radical should be first tested for in a portion of the solution and if <math>(FeCy)_3</math> and <math>(FeCy)_2</math> are found, the solution must be evaporated to dryness with <math>HNO_3</math>, and the residue strongly heated (see General Table) after filtering from the <math>H_2S</math> precipitate, in order to destroy these cyanogen radicals.</p>	<p><b>Residue:</b> Boil with <math>KHO</math> solution for several minutes, then add some <math>Na_2CO_3</math> solution as long as it causes any precipitate in the solution or the filtrate, and boil again; filter and wash the residue: then add more <math>KHO</math>,* heat and filter:</p>	<p><b>Filtrate:</b> Pass <math>H_2S</math>, and if it causes any precipitate continue to pass the gas until the liquid is saturated, then add more <math>KHO</math>,* heat and filter:</p>	<p><b>Residue:</b> Dissolve and examine for metals in the usual manner (385). Note.—Fe, Co, Mn, Cr, if found in this residue, were not present as <math>(Fe_2Cy)_3</math>, <math>(Fe_2Cy)_2</math>, <math>(Co_2Cy)_3</math>, <math>(Co_2Cy)_2</math>.</p>
<p><b>Precipitate:</b> Wash well with boiling water, and boil the precipitate with strong <math>HNO_3</math>, a black residue of <math>HgS</math> may remain, filter this off after diluting the acid, and confirm the presence of <math>Hg</math> in the precipitate by heating it with <math>Na_2CO_3</math> in a bulb-tube (147). The filtrate (or solution if <math>HgS</math> is absent) is evaporated to dryness, the residue is then dissolved in a little hot strong <math>HCl</math>, the solution is much diluted, saturated with <math>H_2S</math>, and filtered:</p>	<p><b>Filtrate:</b> Add dilute <math>HNO_3</math> gradually until the liquid is just acid, pass <math>H_2S</math> to saturation and filter:</p>	<p><b>Filtrate:</b> Divide into two parts, <math>\alpha</math>, <math>\beta</math>:</p> <p><math>\alpha</math>. Examine for acid radicals in the usual way, testing for <math>(Co_2Cy)_3</math> by adding excess of <math>ZnSO_4</math>, filtering and trying whether the precipitate gives a blue bead with borax.</p>	<p><math>\beta</math>. Evaporate to dryness and fuse the residue. When cold boil it with water and filter:—</p>
<p><b>Precipitate, Ex-</b>amine for Pb in excess, and Cu by Table II (485 a).</p>	<p><b>Filtrate:</b> Add <math>AmCl</math>, <math>AmHO</math> in excess, and <math>Am_2S</math>, and examine for Zn, Mn, Ni, Co, etc., by Table III <math>C_3</math> (465).</p>	<p><b>Residue:</b> Dissolve in <math>HCl</math>, and test for Al, Fe, Mn, Co (465); the last three if found were present as cyanogen acid radicals.</p>	<p><b>Filtrate:</b> Acidify a portion if yellow with <math>H_2A</math>, and add <math>PbA_2</math>, a yellow precipitate shows <math>(CrO_4)^{--}</math>, the Cr having been present as cyanogen acid radical.</p>
			<p>Test another part for Al by adding <math>HCl</math> in excess, then <math>AmHO</math> in excess.</p>

\* If  $H_2S$  causes a precipitate to form,  $NaHS$  (or  $KHS$ ) may be added drop by drop until it no longer gives any precipitate in the liquid or in the filtrate; this may be done instead of passing  $H_2S$  to saturation and then adding  $KHO$ .

**486.** The advantage of employing this somewhat complicated and tedious method of Fresenius is that it usually enables the analyst to ascertain whether Fe, Co, (Mn, Cr), when present, form a part of a complex cyanogen acid radicle or simply act as metals combined with acid radicles. For if Fe, Co, Mn, Cr, are found in the residue left after boiling with KHO and  $\text{Na}_2\text{CO}_3$ , they were present in the latter form, if found in the alkaline filtrate they were present as cyanogen acid radicles; if they were detected as cyanogen acid radicles in the aqueous solution they were entirely present in that solution in that form, although they will also be detected in the examination for metallic acid radicles; if they were detected only as metallic radicles and not as cyanogen acid radicles, they were present as metallic radicles in the aqueous solution.

**487.** The following two processes have been proposed for removing cyanogen from a substance before analysis, after which the substance may be examined for metals in the usual way (382 *et seq.*); but these methods, though much more easy and rapid, are less perfect than the preceding ones, since Fe, Co, Mn, Cr, if detected, may have been present either as cyanogen acid radicles, or as metals united with an acid radicle, or in both forms.

1. Pour upon the powdered substance strong  $\text{H}_2\text{SO}_4$  in a porcelain crucible, evaporate to dryness and ignite the residue strongly; when cold dissolve it by heating with a little strong HCl, adding water and heating again. (Rose.)

2. "Cyanogen compounds are best destroyed by fusion in a porcelain crucible with 3 or 4 times their weight of a mixture of 3 parts of  $\text{Am}_2\text{SO}_4$  and 1 part of  $\text{AmNO}_3$ ." The residue left in the crucible is dissolved and examined for metals in the usual way.



### EXAMPLE SHOWING HOW TO ENTER THE RESULTS OF ANALYSIS OF A COMPLEX SUBSTANCE.

The substance given for analysis consisted of a powder containing pink, blue, white, and black particles. It smelt faintly of ammonia.

#### 488. PRELIMINARY EXAMINATION FOR METALS.

Experiment.	Observation.	Inference.
<b>EXPERIMENT I.</b> —Heated in a small dry test-tube.	Water given off which turned red litmus-paper blue. Strong smell of $\text{NH}_3$ gas. Substance blackened, no smell of burning.	Pres. of $\text{NH}_4$ . Pres. of $\text{NH}_3$ . Prob. pres. of Co, Cu, and abs. of T and A.
<i>Confs.</i> —Held in the upper part of the tube a glass rod, with a drop of lime-water hanging on its end.	Slight white sublimate. The lime-water became milky. Brown nitrous fumes evolved, recognized by their smell. Cl gas evolved; found by smell and bleaching litmus.	Pres. of As, $\text{NH}_4$ , or Hg. $\text{Co}_2$ evolved. Pres. of nitrate.
<i>Confs.</i> —Boiled a portion of the substance with KHO solution.	$\text{NH}_3$ gas was evolved; recognized by its smell, and by giving white fumes with strong HCl.	Pres. of Cl.
<i>Confs.</i> —Heated strongly another portion of the dried substance with $\text{Na}_2\text{CO}_3$ in a bulb-tube.	No mirror formed.	Pres. of $\text{NH}_4$ . Abs. of Hg. and As.
<b>EXPERIMENT II.</b> —Heated a portion of the substance on a loop of platinum wire in the Bunsen flame, moistened with HCl, and heated again in the flame. Heated for some time in the tip of the inner blow-pipe flame, moistened with HCl, and again held in the Bunsen flame.	<i>Bright-yellow</i> flame. The flame appeared <i>purple</i> through the indigo-prism.  <i>Crimson</i> color appearing <i>intense red</i> through the indigo-prism. <i>Bright green</i> color with <i>blue</i> core.	Pres. of Na.  Pres. of K.  Pres. of Sr. Pres. of Cu.

Experiment.	Observation.	Inference.
EXPERIMENT.III.—Heated a portion of the substance on charcoal in the inner blowpipe flame.	The greater part of the substance fused readily, and was absorbed by the charcoal. Deflagration occurred.	Pres. of alkali salt. Pres. of chlorate, nitrate.
Confs.—Fused in a clear borax bead in the outer and inner blowpipe flames.	A red metallic residue remained. In the outer flame a bead <i>green</i> whilst hot; <i>blue</i> when cold. In the inner flame <i>red</i> and nearly opaque.	Pres. of Cu. Pres. of Cu.
Confs.—Fused on platinum foil with $\text{Na}_2\text{CO}_3$ and $\text{KNO}_3$ .	No <i>bluish-green</i> or <i>yellow</i> mass on cooling.	Abs. of Mn and Cr.
Confs.—Fused on charcoal in the inner blowpipe flame with $\text{Na}_2\text{CO}_3$ .	Red metallic residue. A portion, placed on a silver coin and moistened, gave no black stain.	Pres. of Cu. Abs. of S.

### 489. PRELIMINARY EXAMINATION FOR ACID RADICLES.

Experiment.	Observation.	Inference.
EXPERIMENT I.—Added dilute HCl without heating.	A colorless gas was evolved, which was free from smell, and turned a drop of lime-water milky.	Pres. of carbonate. Abs. of sulphite, hypochlorite, etc.
Heated to boiling.	Cl was evolved, recognized by its smell, and by bleaching moist litmus-paper.	Pres. of nitrate, chlorate, or some other oxidizing substance.
EXPERIMENT II. — Added strong $\text{H}_2\text{SO}_4$ .	A bright yellow chlorous gas evolved, which crackled when warmed.	Pres. of chlorate.
Heated strongly, cooled, and rinsed out.	The tube when dry was seen not to be etched.	Abs. of fluoride.
Confs.—Dropped in copper turnings and heated.	Reddish-brown fumes evolved.	Pres. of nitrate.
Confs.—Boiled a portion of the substance with water, added strong $\text{H}_2\text{SO}_4$ , cooled, and poured in $\text{FeSO}_4$ solution carefully.	A brown ring formed on the surface of the acid.	Pres. of nitrate.

**490. EXAMINATION FOR METALS IN THE WET WAY.**

Boiled a portion of the substance with  $\text{Fe}_2\text{Cl}_6$ ,  $\text{FeSO}_4$ , and  $\text{KHO}$ , added  $\text{HCl}$  in excess, no blue pp.—*Absence of Cy.*

Proceeded to dissolve the substance and examine the solution by the General Table. (See pp. 230, 250.)

Boiled a portion of the substance with water; as it did not completely dissolve, the sol<sup>n</sup> was decanted, and a little dilute  $\text{HCl}$  poured upon the undissolved residue: effervescence occurred, and on boiling  $\text{Cl}$  was smelt. Added a little strong  $\text{HCl}$ , boiled as long as any smell of  $\text{Cl}$  was perceived; the substance was completely dissolved. Cooled this sol<sup>n</sup> and mixed it with the water sol<sup>n</sup> which had been proved previously to give no pp. on addition of a little dilute  $\text{HNO}_3$ :

No pp. was formed. <i>Abs. of Group I.</i> <i>Hg', Ag, and</i> <i>prob. Pb.</i>	Diluted with water and passed $\text{H}_2\text{S}$ until the liquid smelt strongly of the gas, filtered:		
	<i>A black pp.</i> <i>Exam'd by</i> <i>Table II.</i>	<i>Filtrate</i> which gave no further pp. with $\text{H}_2\text{S}$ was pink ( <i>prob. pres. of Co.</i> ). Boiled until it no longer smelt of $\text{H}_2\text{S}$ , added some strong $\text{HNO}_3$ and boiled to dryness (the color of the sol <sup>n</sup> changed to light-brown on addition of $\text{HNO}_3$ , <i>prob. pres. of Fe</i> ), warmed the residue with $\text{HCl}$ , it dissolved completely ( <i>Abs. of <math>\text{H}_2\text{SiO}_4</math></i> ). To a small portion of the $\text{HCl}$ sol <sup>n</sup> added $\text{AmHMoO}_4$ and warmed: no pp.: <i>Absence of <math>(\text{HO})'''</math></i> . To the rem <sup>n</sup> added $\text{AmCl}$ then excess of $\text{AmHO}$ , and boiled; filtered:	
		<i>A brown pp.</i> <i>Exam'd by</i> <i>Table IIIA.</i>	<i>Filtrate</i> was again pink ( <i>presence of Co.</i> ): added much $\text{Am}_2\text{S}$ and boiled, filtered:
			<i>A black pp.</i> <i>Exam'd by</i> <i>Table III B.</i>
		<i>A white pp.</i> <i>Exam'd by</i> <i>Table IV.</i>	<i>Filtrate</i> was yellow ( <i>Absence of Ni</i> ): added $\text{Am}_2\text{O}_3$ , filtered:  <i>Filtrate.</i> <i>Exam'd by</i> <i>Table V.</i>

491. EXAM<sup>n</sup>. OF PP. IN GROUP II.—Removed from the filter into a porcelain dish and boiled with KHO, filtered :

<i>Filtrate:</i> Acidified with HCl, a white milky liquid only: <i>Abs. of Group II B.</i>	<i>Pp.</i> removed from filter by a knife-blade into a porcelain dish and boiled with strong HNO <sub>3</sub> as long as any red fumes came off, added dilute H <sub>2</sub> SO <sub>4</sub> and stirred well:		
	<i>No pp.:</i> <i>Abs. of Hg and Pb.</i>		
	Added to sol <sup>n</sup> . excess of AmHO:		
	<i>No. pp.:</i> <i>Abs. of Bi.</i>	Acidified the sol <sup>n</sup> . with HCl and saturated with H <sub>2</sub> S; filtered off the black pp. rapidly and boiled it with dilute H <sub>2</sub> SO <sub>4</sub> :	
		<i>Pp.:</i> dissolved in a little boiling dilute HNO <sub>3</sub> added AmHO in excess, then excess of H <sub>2</sub> A, then K <sub>2</sub> FeC <sub>2</sub> O <sub>6</sub> , chocolate-red pp.: <i>Pres. of Cu.</i>	<i>Filtrate:</i> diluted much and passed H <sub>2</sub> S, no pp.: <i>Abs. of Cu.</i>

#### EXAM<sup>n</sup>. OF PP. IN GROUP III A.

Dissolved in a little boiling dilute HCl, added pure NaHO in excess, boiled and filtered :

<i>Filtrate:</i> added HCl gradually till the liquid became acid (no pp.), then AmHO in excess: no pp.: <i>Abs. of Al.</i>	<i>Pp.:</i> dried and fused on platinum foil with Na <sub>2</sub> CO <sub>3</sub> and KNO <sub>3</sub> , boiled the colorless ( <i>abs. of Cr</i> ) mass when cold with water: decanted from the undissolved residue:	
	<i>Residue</i> in the dish dissolved by boiling with a little HCl, added several drops of KCyS: a blood-red coloration: <i>Pres. of Fe.</i>	<i>Sol<sup>n</sup>.</i> being colorless proved <i>abs. of Cr</i> . Acidified with H <sub>2</sub> A, boiled, and added PbA <sub>2</sub> , no yellow pp.: <i>Abs. of Cr.</i>

#### EXAM<sup>n</sup>. OF PP. IN GROUP III B.

Rinsed the pp. off the filter with some cold dilute HCl, stirred well, filtered :

<i>Pp.</i> was black and had already been proved to contain no Ni: fused a portion into a clear borax bead: bead blue in both flames: <i>Pres. of Co.</i>	<i>Filtrate:</i> boiled until it no longer smelt of H <sub>2</sub> S, added a crystal of KClO <sub>3</sub> , boiled until the smell of Cl ceased, added pure NaHO in excess:	
	<i>No pp.</i> <i>Abs. of Mn.</i>	Passed H <sub>2</sub> S into the sol <sup>n</sup> . no pp.: <i>Abs. of Zn.</i>

EXAM<sup>n</sup>. OF PP. IN GROUP IV.

1. *Exam<sup>n</sup>. by Flame-col<sup>n</sup>.*—Dissolved a small quantity of the pp. in a few drops of HCl upon a watch-glass, dipped a loop of platinum wire into the sol<sup>n</sup>. and held it in the Bunsen flame: a crimson-red color was imparted to the flame, which appeared intense red through the indigo-prism: *Pres. of Sr.* This coloration was followed by a yellowish-green, very persistent, col<sup>n</sup>.; *Presence of Ba.*

2. *Exam<sup>n</sup>. in the Wet Way.*—Dissolved the rest of the pp. in as little boiling H $\bar{A}$  as possible; to a small part of the solution, perfectly cool, added CaSO<sub>4</sub> sol<sup>n</sup>.; an immediate pp. formed: *Pres. of Ba.*

To the remainder of the H $\bar{A}$  sol<sup>n</sup>. (proved to be acid to litmus) added K<sub>2</sub>CrO<sub>4</sub> until the liquid appeared yellow, warmed and poured through a double filter:

Pp., which contained all the Ba present, was re-jected.	To the clear filtrate (which was orange red in color) added AmHO until the color changed to light yellow, then added Am <sub>2</sub> CO <sub>3</sub> in excess and filtered:	
	Filtrate was rejected.	Pp.: dissolved in as little boiling H $\bar{A}$ as possible; added to a small portion of the sol <sup>n</sup> . CaSO <sub>4</sub> sol <sup>n</sup> . and boiled, a pp. formed, showing <i>pres. of Sr.</i> To the remainder of the H $\bar{A}$ solution added H <sub>2</sub> SO <sub>4</sub> , boiled, filtered, and to the filtrate (which gave no pp. after being boiled with more H <sub>2</sub> SO <sub>4</sub> ) added AmHO in excess and Am <sub>2</sub> C <sub>2</sub> O <sub>4</sub> , no pp.: <i>Abs. of Ca.</i>

EXAM<sup>n</sup>. OF FILTRATE FROM GENERAL TABLE FOR GROUP V.

Evaporated the filtrate to dryness in a porcelain dish, scraped out the residue upon platinum foil, and ignited strongly until white fumes ceased to appear: dissolved the residue off the foil by boiling with water to which several drops of HCl had been added, divided the sol<sup>n</sup>. into two unequal parts:

<p>To the larger portion added several drops of <math>H_2SO_4</math>, then <math>AmHO</math> in excess and several drops of <math>Am_2C_2O_4</math>, on heating a slight pp. formed; filtered, and added to the clear filtrate <math>Na_2HPO_4</math>, a white crystalline pp.:  <i>Pres. of Mg.</i></p>	<p>Into the smaller portion dipped a loop of platinum-wire, and held it in the Bunsen flame, an intense yellow color:  <i>Pres. of Na,</i>          appearing violet through the indigo-prism:  <i>Pres. of K.</i>          Confirmed the pres. of K by stirring a fresh portion of the soln. with <math>PtCl_4</math> on a watch-glass, a yellow pp.:  <i>Pres. of K.</i></p>
--	--

Metals found: Cu, Fe, Co, Ba, Sr, Mg, Na, K,  $NH_4$ .

#### 492. EXAM<sup>n</sup>. FOR ACID RADICLES.

The following acid radicles have already been detected:  $(CO_3)''$ ,  $(NO_3)'$ ,  $(ClO_3)'$ .

By reference to the table of solubilities (455), under the columns corresponding to the above metals, the only salt insoluble in acids is found to be  $BaSO_4$ ; since Ba is present, and the substance is entirely soluble in acids  $(SO_4)$  is absent.

The following acid radicles have also been proved to be absent in the general examination for metals:

$(CrO_4)''$ , by the HCl solution not becoming green on passing  $H_2S$ .

$(AsO_3)'''$  and  $(AsO_4)'''$ , by no pp. forming in Group II B.

$(SiO_3)''$  and  $(SiF_6)''$ , by no residue insol. in HCl remaining on evaporating to dryness with HCl for Group III A.

$(PO_4)'''$ , by testing with  $AmHMoO_4$  in the HCl sol<sup>n</sup> for Group III A.

The organic acid radicles  $\bar{A}$  and  $\bar{T}$  are absent, since there was no smell of burning on heating the substance.

The only commonly occurring acid radicles remaining to be specially tested for are therefore Cl, Br, I  $(C_2O_4)''$ , and  $(BO_3)'''$ ; and of these Br and I are probably absent,

since no violet fumes of I or brown fumes of Br were evolved with strong  $\text{H}_2\text{SO}_4$ .

Boiled a portion of the substance with pure  $\text{Na}_2\text{CO}_3$  sol<sup>n</sup>., and filtered ; acidified portions of the filtrate with—

$\text{HNO}_3$ .	$\text{H}\bar{\text{A}}$ .
<p><i>Added <math>\text{AgNO}_3</math>, a perfectly white pp., easily sol. in <math>\text{AmHO}</math> :</i></p> <p><i>Pres. of Cl.</i></p>	<p><i>Added <math>\text{CaSO}_4</math>, no pp. :</i></p> <p><i>Abs. of <math>(\text{C}_2\text{O}_4)''</math>.</i></p>
<p>Moistened a piece of turmeric-paper with <math>\text{HCl}</math> sol. of the substance, and dried at <math>100^\circ</math> ; no brown stain produced :</p> <p><i>Abs. of <math>(\text{BO}_3)'''</math>.</i></p>	

493. Found  $\left\{ \begin{array}{l} \text{Metallic radicles : Cu, Fe, Co, Ba, Sr,} \\ \text{Mg, Na, K, NH}_4. \\ \text{Acid radicles : CO}_3, \text{NO}_3, \text{ClO}_3, \text{Cl.} \end{array} \right.$

## SECTION VII.

**APPARATUS, REAGENTS, ETC., REQUIRED FOR  
THE FOREGOING COURSE.****LIST OF APPARATUS REQUIRED BY EACH  
STUDENT.**

**494.** The following list is so arranged that :

Division I contains all apparatus which is required only for the experiments given in Section I.

Division II is apparatus required both for Section I and for the Sections on Analysis.

Division III is apparatus used only in analysis.

Hence a student intending only to perform the experiments in Section I will require the apparatus in Divisions I and II. If he is omitting this portion, and intends only to work through the Sections on Analysis, he must be furnished with the apparatus in Divisions II and III, and will not require that in Division I.

*Note.*—Apparatus contained in large brackets [ ] is useful, but not indispensable.

---

**DIVISION I.** (See also the note at the end of this list.)

- 1 wide-mouthed, stoppered, white-glass gas jar ; about 24 oz. capacity, and 2 inches inside across the neck, whose upper edge should be ground so as to be closed with a ground-glass plate.
- 1 small flanged cylinder, 6 inches in height,  $1\frac{1}{2}$  inches internal diameter, ground at the edge.
- 1 deflagrating spoon, brass cap  $2\frac{1}{2}$  inches across.
- 1 small two-necked Woulffe's bottle, not less than 4 oz. capacity ; or a small wide-necked bottle of not less capacity.
- 1 small retort, best tubulated with stopper, about 4 to 6 oz. capacity.



## DIVISION II.

- 6 test-tubes, 5 or 6 inches long and from  $\frac{1}{2}$  to  $\frac{3}{4}$  inch in diameter.
- 1 test-tube cleaner.
- 2 lengths of hard glass tubing, rather more than  $\frac{1}{8}$ th inch internal diameter.
- 1 thistle funnel.
- [1 clock-glass, about 4 inches across.]
- 2 round ground-glass plates, 3 inches across.
- 2 porcelain dishes, one  $2\frac{1}{2}$  inches across, one 3 inches across.
- 3 watch-glasses, about 2 inches across.
- 2 small flasks, about 4 oz. capacity.
- Red or black india-rubber tubing;\* 1 piece 18 inches long and  $\frac{5}{8}$ th inch internal diameter, 1 piece 6 inches long and  $\frac{3}{8}$ th inch internal diameter.
- 1 Bunsen burner\* with separate rose-top.
- 1 small pestle and mortar.
- 1 iron tripod stand.
- 1 piece of wire gauze, 5 inches square.
- 1 round file.
- 1 triangular file.
- 1 small set of cork-borers.
- 1 small retort-stand, 12 inches in height, and fitted with 3 brass rings.

## DIVISION III.

- 4 glass funnels, two 2 inches across at the top, two  $2\frac{1}{2}$  inches across at the top.
- 2 tobacco-pipe cleaners, for cleaning glass tubes.
- 2 boiling tubes, 6 or 7 inches long, from  $1\frac{1}{4}$  to  $1\frac{1}{2}$  inches across.
- 1 test-tube stand with at least 12 holes.
- 1 nest of 3 or 4 small-sized beakers.
- 1 length of glass rod.

---

\* The Bunsen burner and gas-tubing may be fixtures on each working bench, and will not then be required for each student.

- 1 small porcelain crucible about  $1\frac{1}{2}$  inch across, and lid.
- 1 strip of platinum foil, 2 inches by 1 inch.
- 2 pieces of platinum wire,\* each  $1\frac{1}{2}$  to 2 inches long.
- 1 Black's blowpipe.
- 1 pipeclay triangle, 2 inches in the side.
- 1 pair of brass crucible tongs.
- 1 large flask for wash-bottle, 18 oz. capacity and from 1 to  $1\frac{1}{2}$  inch across the neck.
- 1 wooden filtering-stand.
- 1 tin filter-dryer.
- 1 small wicker basket for draining test-tubes and other glass and porcelain apparatus after they have been washed.

Little toilet-baskets, 12 inches in length by  $5\frac{1}{2}$  and  $4\frac{1}{2}$  inches high, serve well.

*Note.*—For the experiments with gases the student will also require:

- A pneumatic trough, or earthenware pan as a substitute for it (see Note, p. 20).
- A small piece of candle.
- A wax taper, or slips of wood.
- A few sound corks of such a size as to fit test-tubes or small flasks.

---

*Caution.*—In a laboratory where many students are working, it is best to mark at once each piece of apparatus with initials, number, or with some private mark, so as to be able to recognize it if mislaid. Glass, metal, or porcelain apparatus may be notched with the edge of the three-cornered file, or characters may be scratched upon it with the sharp points made by breaking off the tip of a three-cornered file. Wooden apparatus can be written upon with pen and ink, or scratched with the tip of a penknife.

---

\* Platinum wire containing Ba is not uncommon in commerce; it is known by giving a persistent green color to the Bunsen flame, and is useless for flame colorations.

## APPARATUS FOR GENERAL USE.

**495.** The following apparatus need not be purchased by each student, but should be kept in constant readiness for general use:

**496.** *An Indigo-prism.*—A hollow glass-prism having an extremely small angle at its extremity, with ground-glass stopper fitting water-tight into the neck, is purchased; this is to be very nearly filled with solution of indigo, and the stopper then tied in securely with fine copper wire. The indigo solution is made by diluting a solution of indigo in strong sulphuric acid with water, in a thin glass beaker or a porcelain dish, until it has the required intensity of color;\* the liquid is then allowed to stand for several days to give time for a sediment to form and settle completely, and when quite clear is decanted into the prism. In the thicker part of the prism the blue must be so intense as to arrest the passage of the light from a flame colored yellow with a sodium compound, and to cause the potassium-flame coloration to appear red.

**497.** *Several small Slips of Cobalt Glass.*—This glass is used for the same purpose as the indigo-prism, but the prism is to be preferred more especially for the examination for Sr.

**498.** *Set of Cork-borers.*—These are used for perforating a cork into which a piece of glass tubing has to be inserted. Nests of borers, containing three or four small sizes, will be sufficient.

**499.** *Sulphuretted-hydrogen Apparatus.*—It is highly advisable to employ a large apparatus which will supply the gas to all who are working in the laboratory. The use of a small private apparatus by each student, not only causes unnecessary waste of materials and time, but vitiates the air of the laboratory whilst it is being washed out.


Many forms of apparatus have been devised to supply the gas in a constant stream; probably the follow-

---

\* The addition of the indigo solution to 10 times its measure of water will usually give the required shade.

ing, the simplicity, efficiency, and cheapness of which render it very satisfactory, will be found the most generally useful :

FIG. 39.



*Note.*—The level of the liquids in the vessels is shown by dotted lines; the vessel A when lowered from its stand is also shown in dotted outline. The whole is drawn in section.

**500. *Sulphuretted-hydrogen Apparatus.***—Two large bottles, A and B, with wide necks and tubulures near the bottom, and of at least two quarts capacity, are fitted as shown in section in Figure 39.\* India-rubber stoppers should be used instead of corks, as the latter soon become sodden with acid and rotten, and also lose their elasticity and permit leakage. Into B (after laying it upon its side to prevent breaking the bottom) is placed a quantity of small pebbles or broken glass, sufficient to form a layer about half an inch deep; upon this is dropped the ferrous sulphide in lumps; the cork is then inserted the clamp at E being closed. The glass tube fixed in the lower tubulure of B should be bent downwards, so that its end nearly touches the bottom of the vessel. This

\* This apparatus may be purchased completely fitted from Messrs. Mottershead of Manchester and from most apparatus sellers.

drains the acid out of B to the lowest possible level when A is lowered. Strong commercial hydrochloric acid\* is poured into A, then an equal bulk of water is added, and the liquids are well mixed by shaking the vessel.

When the gas is required, the vessel A is raised by being placed on a stand of suitable height, the acid runs into B, generates the gas by acting on the ferrous sulphide, and by the pressure of the liquid in A this gas is forced through a little water in the washing-bottle C, and thence through a bent tube at D into the liquid which is to be saturated with sulphuretted hydrogen.

When the flow of the gas is to be stopped a small screw-clamp on the india-rubber joint at E is tightened; by proper management of this screw-clamp the stream of gas can be either totally arrested or controlled to any required extent; this clamp must always be so regulated as to permit of as little waste of gas as possible whilst using the apparatus.

During working hours the vessel A should be kept raised, as shown in the figure; when laboratory work ceases it should be lowered to the position shown in dotted outline. The vessel B is always kept slightly raised by being placed on a wooden block or some bricks, so as to encourage the flow of the last portions of acid out of it into A when A is lowered. When the acid is spent, it is easily poured out of A and replaced by fresh; lumps of ferrous sulphide are added as required through the neck of B. The removal of the spent acid and the cleansing of the apparatus should be performed in the open air, or at a sink in a cupboard provided with a good draught.

The neck of A should be kept stopped with a cork carrying a bent funnel whose bend is filled with oil or glycerin; this lessens the escape of gas dissolved in the acid, and tends to prevent the apparatus from causing a

---

\*  $\text{H}_2\text{SO}_4$  does not answer for this purpose as well as  $\text{HCl}$ , since  $\text{FeSO}_4$  is apt to crystallize in the apparatus, impeding or preventing the action of the acid and its flow through the connecting-tube.

smell. If several students require to use the gas at the same time, the broad stopper of B is perforated and fitted with additional tubes and wash-bottles, or a system of distributing pipes or T-pieces connected with the tube from the cork in B may be employed.

An apparatus for private use may be fitted as shown in Fig. 40. The larger bottle contains pieces of ferrous sulphide upon which HCl, diluted with an equal measure of water, is poured through the thistle-funnel: the gas bubbles through a little water in the smaller bottle, and thence into the solution required to be saturated.

FIG. 40.



The sulphuretted hydrogen apparatus should stand in a cupboard with a glass front and small doors: the cupboard must be furnished with a flue in which a strong draught is created by an argand or ring gas-burner, made of fireclay or steatite and not of metal; if the cupboard doors are kept closed as much as possible, and students when using the gas are careful to avoid allowing its unnecessary escape, the atmosphere of the laboratory may be kept tolerably free from this badly smelling gas. The cupboard should have within it a drain into which are emptied the spent acid and washings of the apparatus.

**500 a. Passing Sulphuretted Hydrogen into a Liquid.**— Each student must keep a bent glass tube (D, Figs. 39 and 40), made according to the directions in par. 9: when the gas has to be passed into a liquid, the shorter end of this tube is fitted into the india-rubber joint E, and the other end is passed to the *bottom of the liquid*; on slightly opening the clamp E, a stream of the gas will bubble up through the liquid and may be regulated by the clamp; when the gas has passed for about five minutes the liquid will generally be saturated; if this is the case it will smell of the gas strongly after the air above the liquid has been blown out of the vessel and the vessel has been well shaken. Great care must be taken to close the clamp at E completely when the gas is stopped.

The tube D must be thoroughly cleansed after use, employing a tobacco-pipe cleaner if necessary for its inside.

**501.** *A small Agate Pestle and Mortar* (Fig. 32, p. 72).—This is required for powdering very hard substances, more particularly minerals. The substance must have been already broken into small fragments, and these are then crushed to powder by pressure and “trituration,”\* until a powder is produced in which no particles or grains are felt when it is pressed or rubbed with the pestle or the finger; such a powder is commonly termed an “impalpable powder.”

*Caution.*—On no account must a substance be broken by placing it in the mortar and striking it blows with the pestle, as this is very liable to cause the fracture of the mortar.

**502.** *Several small Leaden Cups or a Platinum Crucible* should be kept in readiness for testing F by paragraph (295) or (296). The leaden cups are easily made by beating out thick sheet-lead into the required shape, the mouth being of such a size as to be readily covered by a watch-glass; the small leaden inkpots often employed in school-desks serve well for this purpose.

**503.** *A Steam-oven.*—This is a small oven of sheet-

FIG. 41.



copper, heated to the temperature of boiling water (100° C.) by water contained in the space between the

---

\* Rubbing round and round under the pestle.

exterior of the oven and a copper casing which surrounds it (Fig. 41): the outer casing often has a circular opening cut in the top, covered by a lid when not in use, which serves as a water-bath (504). The level of water in the jacket is maintained constant by a little contrivance shown in section at *c*: a small reservoir communicating by a lateral tube with the space between the oven and its jacket is constantly replenished by water which drops into it from the supply-tube (*d*), the overflow of this reservoir is a tube rising in its centre whose end terminates inside above the level of the lateral tube. The outlet for steam (*f*) may be turned downwards, so that any water formed by the condensation of the steam drops into the reservoir (*e*), or it may be made to communicate with the worm of the still (506), the escape steam being thus condensed into distilled water. Distilled water should always be supplied to the steam oven, as this prevents the formation of a troublesome incrustation in the interior; and since the above arrangement, which is all that can be desired when tap-water is supplied, would occasion much waste of distilled water, a better plan of supplying the oven is to place a vessel of distilled water beside the oven, and connect them with a siphon, having a bulb or vertical closed tube at its bend, to prevent the air boiled out from the water from stopping the action of the siphon.

The steam-oven is employed to drive off the moisture from solid substances which require to be dried at a gentle heat.

**504. Several Copper Water-baths.**—The water-bath consists of a hemispherical copper-dish with its edges turned over inwards; a series of flat copper rings of gradually diminishing diameters are made to rest upon the edges of the bath and upon one another's edges, so that the opening at the top can be made as small as may be desired. The bath is about two-thirds filled with water, and is heated on a tripod-stand until the water boils, the excess of steam escaping by a small hole made just below the edge of the bath: the flame should be so regulated that the water is kept boiling quietly. A



small-sized saucepan is a homely substitute for the copper water-bath.

Any liquid requiring to be evaporated at a gentle heat is placed in an evaporating basin upon this bath, its temperature during evaporation cannot then exceed the temperature of boiling water ( $100^{\circ}\text{C.}$ ).

The top of the water-oven is frequently made to serve as a water-bath, see (503). If the water-bath requires to be used for any considerable length of time, it may be supplied with water in the same way as the steam-oven (503).

**504 a.** The following cheap and simple device serves the purpose of both water-bath and steam-oven on a small scale. Two circular plates of sheet zinc are obtained,\* about four inches in diameter, so as to cover the larger-sized porcelain dishes used by each student; in one of these a circular hole concentric with the edge of the plate and two and three-quarters inches in diameter is cut, in the other a similar hole is made one and a half inches across. These covers when placed on one of the larger porcelain dishes, nearly filled with water and boiled by a lamp, convert it into a water-bath, a small dish containing the liquid to be evaporated being supported in the hole of the plates; small quantities of liquid may be evaporated or solid substances may be dried by placing them on a watch-glass supported on the smaller perforated plate.

In a large laboratory, however, steam-ovens and water-baths are usually kept constantly heated by steam or boiling water, and are used in common by all students. The steam from a large water-bath may advantageously be condensed to distilled water by connecting the escape-pipe of the bath with a long tube of the still-worm.

**505.** *Several Small Tubulated Flasks.*—The small flask shown in Fig. 38 (p. 149) is very convenient for the reactions described in pars. 260, 264, and 273. It is perhaps the form of apparatus best suited for the test for a chloride described in (260), because the  $\text{K}_2\text{Cr}_2\text{O}_7$  powder is not introduced through the same tube as the  $\text{CrO}_2\text{Cl}_2$ .

---

\* Any working tinman will make them to order.

escapes by, and the risk of the mixture in the flask spiriting into the delivery-tube is rendered very small. Its neck should be closed by an india-rubber or glass stopper.

---

#### DISTILLATION OF WATER.

As has been already shown (Exp. 31, p. 37), water ordinarily contains certain solid substances dissolved in it which render it in a chemical sense impure; such water is therefore unfit to be employed for the processes of solution and washing, since any impurity thus introduced into a substance during analysis would be considered when detected to have been present in the original substance. The quantity of these impurities present in any particular water-supply will depend upon the nature of the soil with which the water has been in contact previous to its collection. The water supplied in some districts will be found when subjected to the test given in [(526) Remark 41] to be almost perfectly pure; carefully collected rain-water will invariably yield no indication of dissolved impurity; the water supplied to the majority of laboratories will, however, be found to be unfit for analytical purposes, until it has been freed from the solid substances dissolved in it by being subjected to the process of distillation.


In most towns distilled water can be purchased, but it is preferable to distil in the laboratory or its immediate neighborhood all the water required for use. For this purpose the steam obtained either from a steam-boiler or from a copper-still heated by a furnace or gas-burner, or from the water-baths and steam-ovens in the laboratory, is condensed by a tin worm-pipe immersed in a tub through which a constant stream of cold water runs.

506. The following constant apparatus has proved most serviceable in the author's laboratory, and may be used wherever gas and water are laid on in a suitable way. Its advantage lies in the fact that the still is constantly supplied with the hottest portion of the water from the condenser through a lateral tube. When the

apparatus has been properly fitted up and the supply adjusted, it requires no attention whatever except lighting and extinguishing the gas under the still, and turning the water supply on and off when distillation is started and stopped; it is advisable occasionally to remove from the interior of the still, and the supply-tube connecting it with the condensing vessel, the deposit which accumulates by evaporation of the water.

The accompanying sketch (Fig. 42) renders a lengthened description of the apparatus unnecessary; it represents the apparatus in section. A copper-still (A) is heated by the flame of a large-sized Bunsen burner, or by a ring gas-burner, the steam passing off from the top

FIG. 42.



of the still through a tin tube bent into a spiral form in the condensing-tub (B) which is made of galvanized iron; from this condenser the distilled water flows into a large stoneware vessel (C), with a tap below from

which the water is drawn when required. A long siphon-tube (D) serves as a gauge. The condensing-tub (B) has three tubes let into the side facing the still; one of these (*d*), placed about an inch above the bottom of the vessel, is the inlet for cold water from a cistern whose water-level must be higher than that maintained in A and B; this level is marked by the single-dotted line, and is kept constant by the overflow-tube *e*, which is connected with a pipe leading to a sink or drain. Just below this overflow-tube is a short tube (*f*), which is on a level with another (*g*), opening into the side of the still; by connecting these two tubes with an india-rubber joint, a supply of warm water from the upper part of the condenser is furnished to the still, keeping the level of water in it constant; a small glass jet drawn out from a piece of glass tubing is fitted into this connecting-tube, and has been found sufficient to prevent a too free circulation of water between the still and condenser. This still is supported on an iron ring, projecting as a bracket from the condenser. By having two taps in the inlet-tube (*d*), or by using two clamps on the india-rubber joint which connects it with the supply-pipe from the cistern, much time is economized, as the flow of water may be regulated once for all by one tap or clamp, the other being used only for stopping and starting the stream. The copper still should be furnished with a lid screwing upon its mouth, for convenience in removing the deposit which collects within, and into this lid should be fastened a brass tube carrying a screw-union by which the tin condensing-pipe may be fastened steam-tight to the still. The wooden stand for the still and storing jar is conveniently fitted with shelves and doors to serve as a cupboard.

For tests of purity see Remark 41 (526).

Such a copper still, 23 inches in circumference below and 8 inches in height, when filled to a height of 4 inches and heated by the largest sized Bunsen burner, yields 2 litres of distilled water per hour.

## RECOVERY OF THE METALS FROM SILVER AND PLATINUM RESIDUES.

When  $\text{AgNO}_3$ , or  $\text{PtCl}_4$  has been added to a solution the liquid is not to be thrown into the sink.

**507. Silver Residues.**—If  $\text{AgNO}_3$  has been added to a liquid it must be emptied into a vessel\* labelled "Silver Residue," containing strong  $\text{HCl}$  which precipitates  $\text{AgCl}$ . When the vessel is full the  $\text{AgCl}$  is allowed to settle, and the liquid decanted from the  $\text{AgCl}$ ; more  $\text{HCl}$  is then poured in, and the vessel is again ready for use. When a sufficiently large quantity of  $\text{AgCl}$  has been formed, let it settle, decant off the liquid, and wash the precipitate well by decantation; pour it upon a filter, and dry upon the filter. One of the two following methods may then be employed to obtain metallic  $\text{Ag}$  from the  $\text{AgCl}$ :

1. Mix the  $\text{AgCl}$  with twice its weight of a mixture in equal proportions by weight of  $\text{Na}_2\text{CO}_3$  and  $\text{K}_2\text{CO}_3$ ; place this in a clay crucible of such a size that the mixture only half fills it, and heat in a furnace until the mass becomes liquid; maintain it in a fused condition for about five or ten minutes, remove the crucible, and tap its bottom several times gently upon a brick, to cause the fused particles of  $\text{Ag}$  to unite. Allow the crucible to cool, break it up, and wash the button of  $\text{Ag}$ , which will be found at the bottom, until it is quite free from adhering salts.

This button is then dissolved by heating it in a flask with pure strong  $\text{HNO}_3$ , previously diluted with about half its measure of water; the solution is evaporated to dryness, the residue dissolved in a little water and once more evaporated to complete dryness. This  $\text{AgNO}_3$  is then dissolved in the right proportion of water to furnish reagent No. 48 (par. 525).

2. A readier method for decomposing the  $\text{AgCl}$  consists in placing it in a dish with a little water acidified

---

\* Most conveniently a large bottle, with a funnel in its neck.

with dilute  $\text{H}_2\text{SO}_4$ , and laying upon the  $\text{AgCl}$  some strips of Zn free from Pb; after some hours the Ag will be separated as a black spongy mass. This is well washed by decantation with hot dilute  $\text{H}_2\text{SO}_4$ , and then washed also by decantation with boiling distilled water till free from the  $\text{H}_2\text{SO}_4$ ; the Ag is dissolved in  $\text{HNO}_3$  as directed in the first process, the solution evaporated to dryness, and the  $\text{AgNO}_3$  used for preparing the reagent. Any residue insoluble in  $\text{HNO}_3$  will be unreduced  $\text{AgCl}$ ; this may be filtered off and added to the silver residues for subsequent reduction.

**508. Platinum Residues.**—Liquids to which  $\text{PtCl}_4$  has been added are poured into a vessel\* labelled "Platinum Residues;" when a sufficient quantity has been obtained the liquid and precipitate are poured into a porcelain dish, evaporated to dryness, and heated strongly for some time; as soon as the dish is cool the residue is boiled with water, which is decanted, and upon the residue some solution of oxalic acid is poured; this is evaporated to dryness and the residue strongly ignited. The residue of metallic Pt is once more washed with boiling water, and is then dissolved by heating it with  $\text{HCl}$  to which one-third its measure of  $\text{HNO}_3$  has been added; the solution is evaporated to dryness, the last part of the process being performed over a water-bath.  $\text{HCl}$  is then poured in, and it is once more evaporated to dryness and heated for some time on the water-bath; this residue when dissolved in water forms the reagent No. 28 (525).

---

\* Conveniently a large glass bottle, with a funnel in its neck.

## DIRECTIONS FOR PREPARATION OF REAGENTS.

*Note.*—The reagents required for chemical analysis are much more readily obtained at the present time than they were in former years. They can usually be purchased in a pure state, and therefore processes of preparation, which formerly of necessity occupied a large portion of Manuals of Practical Chemistry, are here omitted. It will be found usually more economical to purchase pure chemicals than to prepare them.

509. Many reagents are required in a dissolved or diluted state; it is a matter of much importance that these solutions should be made of a proper strength, and that the methods of preparing them should be as rapid and simple as possible.

The strength of the solutions is chosen somewhat arbitrarily; indeed, the strength of a solution may be varied with advantage, according to the different purposes for which the reagent is to be employed. It is found, however, that practically a solution of one strength may usually be made to serve all analytical purposes. The proportions of liquid or solid to be mixed with or dissolved in a certain quantity of water, which are given hereafter, are those (or very nearly those) recommended in standard works on analysis. It is not intended, however, to assert that they are for *all* purposes the best, but merely that they are practically useful.

510. A good and economical store-bottle for liquid reagents is the "Winchester Quart" bottle, a stock of which generally accumulates in a laboratory, being continually introduced filled with strong acids. The following methods of filling these have been found simple and easy of execution.

The process of preparing solutions divides itself naturally into the "dilution of liquids" (511), and the "solution of solids" (512).

---

511. DILUTION OF LIQUIDS.

In the following lists the proportions by measure are stated in which the liquid and distilled water are to be mixed; these preparations are roughly, but sufficiently

accurately, obtained by measuring with a foot-rule the height of the bottle outside from the bottom to the projecting ridge at the shoulder, then dividing this height in the required proportions, and marking the point of division by a line scratched upon the bottle with a three-cornered file. Thus dilute HCl (No. 2, 524) is made by mixing strong HCl with three times its measure of water; supposing, then, the measured height of the "Winchester" to the shoulder to be 8 inches, a file-mark is made 2 inches from the bottom, and on filling the bottle to this mark with strong HCl, then up to the shoulder with distilled water, stoppering and shaking up, a dilute acid of the required strength is obtained. This method serves for numbers 2, 3, 4, 6, and 7 (524). For the preparation of No. 1, see column of remarks in the same paragraph.

---

## 512. SOLUTION OF SOLIDS.

For dissolving solids two methods are here described; the former (514) is useful when the solution has to be rapidly prepared, the latter (515) requires less attention and may be used when the solution is not required at once.

**513. Note.**—Against each reagent prepared from a solid there is placed, in paragraph (524) a number, signifying the number of grammes (see *Weights and Measures*, 560 *et seq.*) to be dissolved in 1 Winchester quart of water, and in all the lists will be found the proportion by weight of solid to water. The former number is obtained from the proportion stated in the next column, by considering that an average-sized "Winchester," filled an inch above the ridge round its shoulder, holds about 2400 c.c., and since each c.c. (see *Weights and Measures*, 560 *et seq.*) of water weighs approximately one gramme, we have at once the weight of water (2400 grammes) contained in the bottle, from knowing its capacity;\* and

---

\* This method may also be employed when smaller quantities of the reagent are to be prepared. Measure the number of c.c. required, count the number as grammes, and calculate the proportion of solid required as above.



the weight of water being known, the weight of solid to be dissolved in it is readily calculated from the proportion given.

Thus the proper strength of  $\text{AmCl}$  solution (No. 5, 524) is 1 of solid  $\text{AmCl}$  dissolved in 8 of water, as shown in the fifth column, or a given weight of water must have dissolved in it  $\frac{1}{8}$ th its weight of  $\text{AmCl}$ ; since a "Winchester" holds 2400 grammes of water, we must then dissolve  $\frac{2400}{8} = 300$  grammes of  $\text{AmCl}$  in it, to obtain a solution of proper strength.

**514.** Weigh out the required quantity of the solid substance, dissolve by heating it with distilled water in a large-sized porcelain dish placed on a tripod-stand over a rose-burner; this solution would often crystallize on cooling, hence it should be at once poured into some cold distilled water contained in a large beaker marked to show the quantity of liquid required to fill the "Winchester." Whilst pouring off the liquid press a wet-glass rod, held vertically, against the lip of the dish, and let the bottom of the dish touch the inside of the top of the beaker. These precautions will, with proper care, prevent the liquid from running down outside the beaker. Dilute this up to the "Winchester" mark with distilled water, stir well, and if turbid let stand covered from dust until clear; then pour off into the store-bottle, carefully leaving the sediment behind. If the liquid is wanted in haste, it may be filtered into the bottle. The coarse brown French filter-paper is useful for this purpose, being strong and porous. It is best to filter the liquid before diluting it to the full extent, as the quantity to be passed through the filter is thus lessened and time economized.

The above method is of general application in the solution of solids.

**515. II.**—The following liquid is advantageous, because the process of solution once started proceeds continuously without further attention. It depends on the fact that the specific gravity of water becomes higher the greater the quantity of solid dissolved in it; hence if the solid is immersed in the water, and supported near the surface, a circulation of the liquid is produced by the constant sinking of the water which has been in

contact with the solid and by dissolving some of it has increased in specific gravity, this being replaced by the comparatively lighter liquid from below. This circulation of water over the solid, gradually dissolves it without the use of heat, and without requiring any attention after the process has once been started.

A glass cylinder (A, Fig. 43) open at both ends, has one end covered with a piece of muslin (shown by a fine dotted line), which is fastened on by an india-rubber ring, embracing its edges; this cylinder, with a muslin bottom, is supported in a large beaker by means of two pieces of glass rod or tube, bent as shown at *c c c*; the cylinder stands at such a height that the muslin bottom is about two inches below the mark made on the beaker to show the Winchester quart measure. Distilled water is then poured into the beaker until it buries the muslin to the depth of about half an inch; the weighed substance, best in small pieces, is placed in the cylinder, and the whole allowed to stand until the solid has been dissolved; with large quantities of substance this will usually require several hours, with smaller quantities about twenty minutes or half an hour.





FIG. 43.

The processes of solution and filtration are performed simultaneously by laying a piece of filter-paper upon the muslin before fastening it on the cylinder; the filter-paper will retard the process, but the coarse gray kind referred to above will do so the least.

Perhaps the most advantageous way of using this process, especially for dissolving large quantities of substance, is to start it overnight; the solution will then be completed by the morning. In this case the filter-paper will be unnecessary, as any fine particles which have passed through the muslin will have settled, and the liquid can be decanted from them.

It must be borne in mind that the solution of many

salts in water, especially such salts as contain water of crystallization, increases the volume of the water; hence the quantity of water placed in the beaker must be less than the Winchester-quart measure, and the liquid is made up to the required volume, if necessary, after solution has been effected.



As a convenient substitute for the glass cylinder, A, with muslin bottom, an earthenware colander may be employed (Fig. 43 a), filtration being effected

when necessary by laying a piece of filter-paper inside the colander before placing in it the fragments of the substance. Should the beaker be able to contain more than the Winchester quart of water, a stoppered bottle of suitable size may be immersed in it to raise the water-level sufficiently to reach the substance.

---

#### PREPARATION OF SOLUTIONS REQUIRED FOR TRYING THE REACTIONS OF METALS AND ACID RADICALS, SECTION IV.

**516.** The solutions required for trying these reactions should be prepared of a strength suited to yield them in a characteristic manner; it is therefore advisable that solutions of the requisite strength should be kept in readiness, since if the student has to dissolve some of the solid substance himself, he is liable not to obtain satisfactory results, either because his solution is made inconveniently strong, or because it is in a too diluted state. The plan of keeping these solutions in stock is also recommended by the consideration that it economizes the student's time, and tends to prevent waste of chemicals.

**517.** In paragraphs (529) and (530) are lists of the solutions which are required for the complete series of reactions in Section IV: the first column gives the number, the second the name, and the third the formula of each substance; in the fourth is stated the number of

grammes which are to be dissolved in one Winchester quart of water or acid; in the fifth column the proportion by weight of solid to water, from which the number in column 4 was derived; and in the sixth column will be found special remarks relating to the preparation of particular solutions. The general methods of preparing solutions have already been given in paragraphs (514) and (515).

Pure solid chemicals are almost invariably sold in the crystalline condition, since they have been as a rule purified by the process of crystallization. The weights given below are therefore weights of the crystalline substance; in a few cases the substance was weighed in the uncrystalline condition—these are marked by an (*a*) placed after the name in the first column, signifying “amorphous.” Directions for labelling the store-bottles of these solutions will be found in paragraphs (522) and (523).

---

#### LIST OF THE REAGENTS.

**518.** Paragraphs (524), (525), and (526), contain lists of the reagents, solid and in solution, which should be kept in readiness in the laboratory; they are dissolved in distilled water unless it is otherwise stated.

The reagents are here divided into two classes. The first (524) being in constant use and comparatively inexpensive, should be contained in small bottles placed above each working bench, or there should be at least one complete set for the use of two students. The second class (525, 526) are less frequently required, or are of a more costly nature, and one set may be placed in an accessible part of the laboratory for the use of many students.

**519.** Each reagent is numbered in the following lists; this number is a convenient means of reference, and also serves to keep the reagent bottles always standing in the same order, which much facilitates finding any bottle when wanted; the student is strongly recommended to keep his bottles arranged in the order indicated by these

numbers. The full name and chemical formula accompany each number, and every label in the laboratory should carry legibly the number, name, and formula of the reagent it contains.

**520.** In the last column of each list are placed remarks as to the preparation and the condition of the reagent. The numbers found in the last column but one express the strength of the solution of the reagent if a liquid, by stating the proportion of the strong liquid or solid (as purchased) to the water with which it is mixed, or in which it is dissolved, the proportion being expressed as a ratio in which the water always stands last; in the case of liquids the numbers express proportion by measure (m.), in the case of solids the proportion by weight (w.). A number is also placed in the fourth column opposite each reagent prepared by dissolving a solid substance, which denotes the weight in grammes of the solid to be dissolved in a Winchester quart bottle of water.

Thus, No. 2 (524), hydrochloric acid, is prepared by mixing strong HCl with 3 times its measure of distilled water. No. 5, ammonium chloride is made by dissolving solid AmCl in 8 times its weight of water, or by dissolving 300 grammes of solid AmCl in one Winchester quart of water. Full descriptions are given in paragraphs (511), (514), and (515), of the methods used for making these solutions.

**521.** The special reagents which stand on shelves above the bench are kept in small bottles, the solids (Nos. 18–21, paragraphs (524) in 4 or 6 oz. wide-mouthed stoppered bottles, which are filled when necessary from the store-jars (stoneware jars with ground clasp-covers),\* the liquids (Nos. 1–17, par. 524) in narrow-necked 4 or 6 oz. bottles supplied from the Winchester quart bottles in which the store solutions are kept. The general reagents (525, 526) should be kept in bottles of not less than 12 oz. capacity.

**522. Labelling.**—Bottles intended to contain strong

---

\* Powdered solids are conveniently removed from large stone jars by means of wooden spoons.

acids or alkalies should have labels etched upon the glass: such bottles are easily purchased. An ordinary paper label, however, protected by paraffin varnish, which is applied according to the directions given below (523), will last for a considerable length of time.

The paper labels required for reagents and store-bottles are most conveniently kept ready gummed in the form of sheets either separate or stitched together into the form of a book. They must be stored in a dry place. They should, as was mentioned in (519), bear in bold legible type at least the full name and formula of the substance, the number, if not already printed, being written upon each.\*

Great care should be bestowed on making the moistened label adhere over its whole surface by gentle pressure with a clean cloth.

**523. Varnishing Labels.**—It is advisable to varnish each label, which is intended to be permanent, as soon as it is perfectly dry; the label is thus protected against being soiled or wetted. The ordinary process consists in first sizing the label by brushing over it a solution of isinglass in water, and then, as soon as this has perfectly dried, applying a thin coat of label varnish; the bottle must not be used until this varnish is quite hard.

A much more simple process of varnishing consists in melting some solid paraffin, which must not be heated so strongly as to cause it to emit vapor, and brushing this rapidly over the dry label with an ordinary varnishing brush; a little practice will soon enable one to apply a uniform film which does not soak into the paper, hardens immediately, and resists the action of water, acids, and alkalies. The chief precaution to be observed is to heat the paraffin as little as possible above its melting-point, as, if it is too hot, it greases the paper.

---

\* A book of labels ("A Series of Chemical Labels for Use in Laboratories") is sold by Messrs. Mottershead, of Manchester, which contains the name of each substance in full, printed in clear type, space being left to insert formula and number.

Messrs. Mottershead also supply gummed sheets of labels, printed specially for use with this book, containing all the numbers, names, and formulæ in the lists in pars. 524, 525, 526, 529, 530, 531, and 532.

## 524.—REAGENTS REQUIRED

Reference number.	Name. [The numbers in brackets refer to remarks correspondingly numbered in the sixth column.]	Formula.	Weight of solid in grammes to be dissolved in 1 "Winchester" of water.
<b>Liquids.</b>			
1.	Sulphuric acid <sup>(1)</sup> .....	H <sub>2</sub> SO <sub>4</sub> .....	—
2.	Hydrochloric acid <sup>(2)</sup> .....	HCl.....	—
3.	Nitric acid <sup>(3)</sup> .....	HNO <sub>3</sub> .....	—
4.	Acetic acid <sup>(4)</sup> .....	HA, or HC <sub>2</sub> H <sub>3</sub> O <sub>2</sub> .....	—
5.	Ammonium chloride <sup>(5)</sup> .....	NH <sub>4</sub> Cl.....	300
6.	Ammonia, or Ammonium hydrate <sup>(6)</sup> .....	NH <sub>3</sub> .HO.....	—
7.	Ammonium sulphide <sup>(7)</sup> .....	(NH <sub>4</sub> ) <sub>2</sub> S.....	—
8.	Ammonium carbonate <sup>(8)</sup> .....	(NH <sub>4</sub> ) <sub>2</sub> CO <sub>3</sub> .....	500
9.	Ammonium oxalate.....	(NH <sub>4</sub> ) <sub>2</sub> C <sub>2</sub> O <sub>4</sub> .2H <sub>2</sub> O.....	100
10.	Potash, or potassium hydrate.....	KHO.....	200
11.	Potassium chromate.....	K <sub>2</sub> CrO <sub>4</sub> .....	200
12.	Potassium ferrocyanide.....	K <sub>4</sub> FeCy <sub>6</sub> .3H <sub>2</sub> O.....	200
13.	Potassium ferricyanide <sup>(9)</sup> .....	K <sub>3</sub> FeCy <sub>6</sub> .....	200
14.	{ Sodium phosphate <sup>(10)</sup> , or Hydric disodic phosphate, or Hydrogen disodium phosphate, }	Na <sub>3</sub> HPO <sub>4</sub> .12H <sub>2</sub> O.....	200
15.	Sodium carbonate <sup>(11)</sup> .....	Na <sub>2</sub> CO <sub>3</sub> .10H <sub>2</sub> O.....	600
16.	Calcium sulphate <sup>(12)</sup> .....	CaSO <sub>4</sub> .....	{ (Saturated solution.) }
17.	Barium chloride.....	BaCl <sub>2</sub> .2H <sub>2</sub> O.....	
<b>Solids.</b>			
18.	Sodium carbonate <sup>(13)</sup> .....	Na <sub>2</sub> CO <sub>3</sub> .....	
19.	Borax <sup>(14)</sup> .....	Na <sub>2</sub> B <sub>4</sub> O <sub>7</sub> .....	
20.	Potassium chlorate <sup>(15)</sup> .....	KClO <sub>3</sub> .....	
21.	Test-papers <sup>(16)</sup> .....		

\* Hydrogen sodium ammonium phosphate (NaAmHPO<sub>4</sub>)<sub>4</sub> solution serves better for given the above for NaHPO<sub>4</sub>.

## FOR EACH BENCH.

Proportion of solid by weight (w.) and liquid by measure (m.) to water.	Remarks. [The small numbers refer to corresponding numbers in the second column.]
<p>1 : 5 m.  1 : 3 "  1 : 3 "  2 : 1 "  1 : 8 w.  1 : 3 m.  1 : 3 "  1 : 5 w.  1 : 24 "  1 : 12 "  1 : 12 "  1 : 12 "  1 : 12 "  1 : 12 "  1 : 12 "  1 : 4 "  —  1 : 12 "</p>	<p>1. The strong pure <math>H_2SO_4</math> must be poured in the proper proportion into the water contained in a large thin beaker or an earthen jar, constantly stirring the water all the time: the right proportions are obtained by measurement of the height of the vessel (511). The hot liquid is cooled by immersing the vessel in cold water, and is then poured into the store-bottle. If the acid is pure, no white precipitate (<math>PbSO_4</math>) forms on dilution.  2. Must be colorless, and give after dilution no precipitate with either <math>BaCl_2</math> or <math>H_2S</math>.  3. Must give no precipitate, after dilution with <math>BaCl_2</math> or <math>AgNO_3</math> added to separate portions.  4. Must give no precipitate with <math>BaCl_2</math>.  5. Must give no precipitate or coloration with <math>Am_2S_3</math>.  6. Must give only a very slight precipitate with lime-water, no coloration with <math>Am_2S_3</math>, and separate portions acidified with <math>HNO_3</math> must give no precipitates with <math>BaCl_2</math> and <math>AgNO_3</math>. The "Liquor ammoniæ fortissima" is purchased of 880° specific gravity.  7. Must be yellow, and give with acids <math>H_2S</math> and a white precipitate of S: it must give no precipitate with solutions of Ca or Mg salts.  8. The solid <math>Am_2CO_3</math> is dissolved by being heated with water in a porcelain dish (514): in diluting, one-fourth of the "Winchester" must be filled with strong <math>AmHO</math>.  9. Must give no blue precipitate with <math>Fe_2Cl_3</math> solution.  10. Must yield no precipitate with <math>AmHO</math>.  11. The crystallized salt.  12. See (527).  13. Dry, finely-powdered solid. It must, after having been dissolved in excess of <math>HNO_3</math>, give no precipitate with <math>BaCl_2</math> or <math>AgNO_3</math>, or <math>AmHMoO_4</math> (277), and if evaporated with excess of <math>HCl</math> must leave no residue insoluble in dilute <math>HCl</math> (290 a).  14. The borax is best dried by heating in a platinum or porcelain dish until after melting it has again become solid: this solid mass is then finely powdered in a mortar and kept in a stoppered bottle.  15. In small crystals.  16. Red and blue litmus-papers and turmeric paper in small strips.</p>

precipitation of Mg than does  $Na_2HPO_4$  solution: it is prepared of the strength



## 525.—REAGENTS FOR GENERAL USE,

*Note.*—Except in large laboratories it will be unnecessary to keep which they are contained in the laboratory, may be made up to the proportion stated below in column 4.

Reference number.	Name. [The small numbers in brackets refer to remarks correspondingly numbered in the fifth column.]	Formula.
25.	Sulphuric acid <sup>(1)</sup> .....	H <sub>2</sub> SO <sub>4</sub> .....
26.	Hydrochloric acid <sup>(2)*</sup> .....	HCl.....
27.	Nitric acid <sup>(3)*</sup> .....	HNO <sub>3</sub> .....
28.	†Platinum chloride.....	PtCl <sub>4</sub> .....
29.	Alcohol, Rectified spirit <sup>(4)</sup> .....	C <sub>2</sub> H <sub>5</sub> O.....
30.	Tartaric acid <sup>(5)</sup> .....	$\left\{ \begin{array}{l} \text{H}_2\text{T} \\ \text{H}_2\text{C}_4\text{H}_4\text{O}_6 \end{array} \right\}$ .....
31.	Acid sodium tartrate, Hydric sodic tartrate <sup>(6)</sup> .....	$\left\{ \begin{array}{l} \text{NaHT.H}_2\text{O} \\ \text{NaHC}_4\text{H}_4\text{O}_6.\text{H}_2\text{O} \end{array} \right\}$ .....
32.	Slaked lime <sup>(7)</sup> .....	Ca(OH) <sub>2</sub> .....
33.	Acetic acid <sup>(8)*</sup> .....	$\left\{ \begin{array}{l} \text{HA} \\ \text{HC}_2\text{H}_3\text{O}_2 \end{array} \right\}$ .....
34.	Cobalt nitrate <sup>(9)</sup> .....	Co(NO <sub>3</sub> ) <sub>2</sub> .6H <sub>2</sub> O.....
35.	Hydrofluosillic acid <sup>(10)</sup> .....	H <sub>2</sub> SiF <sub>6</sub> .....
36.	$\left\{ \begin{array}{l} \text{Sulphuretted hydrogen solution,} \\ \text{Hydrogen sulphide,} \end{array} \right\}$ <sup>(11)</sup> .....	H <sub>2</sub> S solution.
37.	Oxalic acid.....	$\left\{ \begin{array}{l} \text{H}_2\text{O}_2.2\text{H}_2\text{O} \\ \text{H}_2\text{C}_2\text{O}_4.2\text{H}_2\text{O} \end{array} \right\}$ .....
38.	Potassium sulphocyanide.....	KCyS.....
39.	Potassium nitrate.....	KNO <sub>3</sub> .....
40.	Lead acetate <sup>(12)</sup> .....	$\left\{ \begin{array}{l} \text{PbA}_2.3\text{H}_2\text{O} \\ \text{Pb}(\text{C}_2\text{H}_3\text{O}_2)_2.3\text{H}_2\text{O} \end{array} \right\}$ .....
41.	Sodium acetate.....	$\left\{ \begin{array}{l} \text{NaA}.3\text{H}_2\text{O} \\ \text{NaC}_2\text{H}_3\text{O}_2.3\text{H}_2\text{O} \end{array} \right\}$ .....
42.	Potassium cyanide <sup>(13)</sup> .....	$\left\{ \begin{array}{l} \text{KCv} \\ \text{KCN} \end{array} \right\}$ .....
43.	Sodium hypochlorite <sup>(14)</sup> .....	$\left\{ \begin{array}{l} \text{Na}_2\text{Cl}_2\text{O} \\ \text{NaClO.NaCl} \end{array} \right\}$ .....
44.	Stannous chloride <sup>(15)</sup> .....	SnCl <sub>2</sub> .2H <sub>2</sub> O.....
45.	Copper <sup>(16)</sup> .....	Cu.....
46.	Zinc <sup>(17)</sup> .....	Zn.....
47.	Steel <sup>(18)</sup> .....	Fe.....
48.	Silver nitrate.....	AgNO <sub>3</sub> .....
49.	Sulphurous acid.....	H <sub>2</sub> SO <sub>3</sub> .....
50.	Copper sulphate.....	CuSO <sub>4</sub> .5H <sub>2</sub> O.....
51.	Magnesium sulphate.....	MgSO <sub>4</sub> .7H <sub>2</sub> O.....
52.	Mercuric chloride.....	HgCl <sub>2</sub> .....
53.	Gold chloride.....	AuCl <sub>3</sub> .....

\* See last column on the preceding page.

† Must be used in very small quantity only, being an expensive reagent.

## FOR THE DETECTION OF METALS.

these reagents in store in quantity. The 12 oz. or 14 oz. bottles in right strength as soon as they are empty, by calculation (518) from

Proportion by weight of solid to water.	Remarks. [The numbers in this column refer to those in brackets in the second column.]
Strong pure.	17. Must be colorless, and form no brown ring when poured below some $\text{FeSO}_4$ solution (247).
" "	18. Strong rectified spirit (methylated), which must leave no residue on evaporation.
1 : 30.	19. The solution should be mixed immediately after preparation with several drops of carbolic acid to prevent a vegetable growth forming in it.
Strong.	20. Dissolve 10 grammes of $\text{H}_2\text{T}$ in 100 c.c.† of water, divide this solution into equal parts, exactly neutralize (35) one part by heating it nearly to boiling, and stirring in solid $\text{Na}_2\text{CO}_3$ in powder; add to this the other part, cool and dilute to 150 c.c. A few drops of carbolic acid should be added to the solution.
1 : 10.	21. Pieces of freshly-burnt lime are placed on a plate, and water is poured upon them until they begin to appear moist on their surface; the superfluous water is then drained off, and as soon as the lime has crumbled to powder, the powder is placed in a broad-mouthed stoppered bottle.
—	22. The bottle should be closed with a loosely-fitting india-rubber stopper, perforated, and with a glass tube passed through it and dipping into the solution; when a drop of the solution is required, the upper end of the glass tube is closed by the finger, and a drop is delivered from the opposite end by slightly relaxing the pressure of the finger.
Solid.	23. The solution must give no precipitate with $\text{Sr}(\text{NO}_3)_2$ solution. For the preparation of it see (528, 4).
Strong.	24. For preparation of this solution see (528, 1). It must not darken on adding $\text{AmHO}$ .
1 : 12.	25. A little $\text{H}_2\text{A}$ must be added to this solution to make it clear.
—	26. Solution of $\text{KCx}$ is made (1 : 12); it decomposes so readily by being kept in water, that it is best made immediately before use by heating a small piece of solid $\text{KCx}$ with distilled water.
1 : 12.	27. The strongest "Liquor Sodæ Chlorinatæ" of commerce.
1 : 100.	28. Crystals of $\text{SnCl}_2$ should be dissolved by heating them with water containing some $\text{HCl}$ ; the solution should be kept in a well-stoppered bottle containing pieces of granulated or sheet tin.
Solid.	29. In strips cut from copper-sheet.
Solid.	30. In strips or rods, or granulated; it must be proved to be free from $\text{As}$ by (181, 182, or 183).
—	31. Common knitting-kneedles broken into short lengths, and kept in a bottle containing pieces of quicklime to prevent rusting.
1 : 12.	
—	
—	
1 : 20.	
(See 528, 2).	
1 : 12.	
1 : 12.	
1 : 20.	
1 : 30.	

† c.c. is the contraction for cubic centimetre. See weights and measures (par. 560).

## 526.—REAGENTS FOR GENERAL USE,

Refer to note under

Reference number.	Name. [The small numbers in brackets refer to corresponding numbers in the fifth column.]	Formula.
57.	Lime-water.....	$\text{Ca}(\text{HO})_2$ .....
58.	Lead acetate in potash <sup>(2)</sup> .....	$\text{PbA}_2 + \text{KHO}$ .....
59.	Potassium dichromate.....	$\text{K}_2\text{Cr}_2\text{O}_7$ .....
60.	Chlorine-water.....	$\text{Cl}$ solution.....
61.	Ferric chloride <sup>(2)</sup> .....	$\text{Fe}_2\text{Cl}_3$ .....
62.	Ferrous sulphate <sup>(2)</sup> .....	$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ .....
63.	Potassium iodide.....	$\text{KI}$ .....
64.	Starch <sup>(1)</sup> .....	—
65.	Indigo solution <sup>(2)</sup> .....	—
66.	Manganese dioxide <sup>(2)</sup> .....	$\text{MnO}_2$ .....
67.	Potassium dichromate <sup>(2)</sup> .....	$\text{K}_2\text{Cr}_2\text{O}_7$ .....
68.	Ether (methylated).....	$(\text{C}_2\text{H}_5)_2\text{O}$ .....
69.	Carbon disulphide.....	$(\text{S})$ .....
70.	Potassium nitrite <sup>(2)</sup> .....	$\text{KNO}_2$ .....
71.	Nitrogen tetroxide solution.....	$\text{N}_2\text{O}_4$ solution.....
72.	Ammonium molybdate <sup>(2)</sup> .....	$(\text{NH}_4)_2\text{HMoO}_4$ .....
73.	Calcium fluoride.....	$\text{CaF}_2$ .....
74.	Potassium disulphate.....	$\text{KHSO}_4$ .....
75.	{ Microcosmic salt, or Hydric ammonic sodic phosphate } <sup>(40 a)</sup> .....	$\text{NaAmHPO}_4 \cdot 4\text{H}_2\text{O}$ .....
76.	{ Wax, or paraffin.....	—
77.	Calcium chloride.....	$\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$ .....
78.	Potassium chloride.....	$\text{KCl}$ .....
79.	Marble.....	$\text{CaCO}_3$ .....
80.*	Distilled water <sup>(1)</sup> .....	$\text{H}_2\text{O}$ .....
81.	Pure sodium hydrate <sup>(2)</sup> , pure soda.....	$\text{NaHO}$ .....
82.	Fusion mixture <sup>(2)</sup> .....	$\text{Na}_2\text{CO}_3 + \text{K}_2\text{CO}_3$ .....
83.	Barium carbonate <sup>(2)</sup> .....	$\text{BaO}$ .....
84.	Solution of sodium acetate in dilute acetic acid, <sup>(4)</sup> .....	$\text{NaA} + \text{HA}$ .....
85.	Sodium nitrate <sup>(2)</sup> (solid).....	$\text{NaNO}_3$ .....
86.	Silica, or finely powdered whitesand. (See 548, 4),.....	$\text{SiO}_2$ .....
87.	Hydrofluoric acid <sup>(1)</sup> .....	$\text{HF}$ .....
88.	Barium hydrate (in crystals).....	$\text{Ba}(\text{HO})_2 \cdot 8\text{H}_2\text{O}$ .....
89.	Zinc sulphate. (See No. 111).....	$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ .....

\* Nos. 80-89 are certain special

## FOR THE DETECTION OF ACID RADICLES.

the heading of (525).

Proportion by weight of solid to water.	Remarks. [The numbers in this column refer to corresponding numbers in the second column.]
See (527). 1 : 24. See (528, 3). 1 : 24. Solid. 1 : 60. Solid. Solid. Solid. 1 : 24. See (528, 5). In powder. In small pieces. In small crystals. Solid, in pieces. 1 : 12. Crystals. 1 : 12. In pieces as large as a pea.	32. To some of the $\text{PbA}_2$ solution (No. 40) $\text{KHO}$ solution is added until, on warming, the precipitate at first formed is just redissolved : paper dipped into this liquid is a very delicate test for $\text{H}_2\text{S}$ (230). 33. The solution should not contain any free acid : its absence is shown by a single drop of $\text{AmHO}$ giving a reddish-brown precipitate of $\text{Fe}_2\text{H}_2\text{O}_4$ in the solution. 34. $\text{FeSO}_4$ solution rapidly oxidizes in the air, hence the $\text{FeSO}_4$ should be kept in the solid state as small green crystals, which should not show a yellow coating in any part ; these are dissolved when required, by heating them with water. 35. Starch solution rapidly changes, it is best therefore to keep the starch as a powder. Starch solution is made by stirring 2 grms. of this powder, which has been made into a paste with 10 c.c. of cold water, into 100 c.c. of boiling water and cooling. 36. Made by diluting the ordinary solution of indigo in strong $\text{H}_2\text{SO}_4$ with about ten times its volume of water, and cooling the mixture ; a precipitate will separate after a time, from which the solution is separated by decantation or filtration. 37. Should be kept in fine powder : it must not evolve $\text{Cl}$ or $\text{CO}_2$ when warmed with strong $\text{H}_2\text{SO}_4$ . 38. In small pieces, or in powder. 39. The solution is prone to change, and should be made in small quantity only : it must evolve copious red fumes when mixed with $\text{H}_2\text{SO}_4$ . 40. Make the solution with the following proportions : 1 gramme of $\text{Am}_2\text{MoO}_4$ is dissolved in 12.5 c.c. of strong $\text{AmHO}$ which has been previously mixed with an equal quantity of water, the solution is allowed to stand (if necessary) till clear, then poured off into 12.5 c.c. of strong $\text{HNO}_3$ mixed with an equal volume of water : the liquid will become hot, and should be allowed to stand until it is cool before being used.
40 a. The solution of this salt must yield no precipitate with $\text{AgNO}_3$ insoluble in $\text{HNO}_3$ . 41. Must leave no residue on evaporation ; in separate portions no precipitates must be caused by $\text{BaCl}_2$ , $\text{AgNO}_3$ , or $\text{Am}_2\text{C}_2\text{O}_4$ , neither must any precipitate or even dark coloration be produced by addition of $\text{Am}_2\text{S}$ . 42. The proportion by weight should be 1 : 10 of water ; it is best kept in green glass bottles, as it slowly dissolves $\text{Pb}$ from white flint glass. The solution must not become dark in color when mixed with $\text{H}_2\text{S}$ , nor give a gelatinous precipitate ( $\text{Al}_2\text{H}_2\text{O}_4$ when mixed with excess of $\text{AmCl}$ solution). 43. Dry finely powdered $\text{Na}_2\text{CO}_3$ and $\text{K}_2\text{CO}_3$ are intimately mixed in the proportion of 53 : 69 by weight, and kept in a stoppered bottle. 44. Pure powdered $\text{BaCO}_3$ is either purchased, or is made by precipitating $\text{BaCl}_2$ solution completely with $\text{Am}_2\text{CO}_3$ (or $\text{Na}_2\text{CO}_3$ ) solution, and washing the precipitate well by decantation. This powder is then mixed with distilled water to the consistency of thin cream. 45. Dissolve 20 grammes of $\text{NaA}$ in 60 c.c. of distilled water, and add to the solution 40 c.c. of strong $\text{HA}$ . 46. By neutralizing strong hot $\text{Na}_2\text{CO}_3$ solution with $\text{HNO}_3$ and evaporating to dryness. Sold also as Chili saltpetre. 47. Purchased and kept in gutta-percha bottles.	

reagents entered apart.

## PREPARATION OF SATURATED SOLUTIONS.\*

## PREPARATION OF SATURATED SOLUTIONS OF SOLIDS.

527. The method described in (515) may be used, or the substance in powder may be constantly shaken up with the solvent for some time, and the solution then separated from excess of the solid by decantation or filtration. The most ready method of preparing a saturated solution of  $\text{Ca}(\text{HO})_2$ , or of  $\text{CaSO}_4$ , is to pour some of the powder into a Winchester quart, then fill it up to the shoulder with water, and mix the powder thoroughly with the water by shaking the bottle; the shaking is repeated at intervals during half an hour or more; the bottle is then allowed to stand until the excess of powder has settled, and the liquid above it is quite clear, when the liquid is poured off into another "Winchester," leaving the powder behind; fresh water is poured upon this powder and more solution made as above by constant shaking, the bottle being allowed to stand by until the fresh supply is required. By thus making a fresh stock of solution whenever one lot is decanted, time is allowed for the freshly prepared solution to become perfectly clear before it is required for use.

1. *Lime-water* is made in the above manner by shaking slaked lime in powder with common *tap-water*.

2. *Calcium sulphate* solution is prepared by saturating *distilled water* by the above means with gypsum or plaster of Paris in powder.

## PREPARATION OF SATURATED SOLUTIONS OF GASES.

528. The gas is made to bubble in a constant stream from the end of a glass tube which dips nearly to the bottom of the liquid (Fig. 44); the liquid must be kept cold, and be contained in a bottle which is fitted with an accurately ground stopper. In order to ascertain whether

---

\* By a "saturated solution" is meant a solution containing the maximum quantity of the substance which the solvent can dissolve.

the solution is saturated, the bubbles of the gas are occasionally watched to see whether they diminish in size as they rise through the liquid; if they do not, the bottle, after being closed tightly by the thumb or by inserting the stopper, is violently shaken; if this causes a pressure and escape of gas from the bottle into the air on unclosing its mouth, the liquid is saturated; if, on the contrary, it causes a pressure of air into the bottle, the gas must be passed again for some time, and the trial repeated, until on shaking as directed above, an outward pressure is noticed.

*Note*—All the preparations mentioned below should be made in a draught cupboard, as the gases are injurious to the lungs if inhaled.

1. *Sulphuretted Hydrogen Solution* is made by fitting the bent tube *a b* (Fig. 44) by means of the india-rubber joint *c*, upon either of the apparatus described in (500) for the preparation of  $H_2S$  gas; the gas is thus made to bubble through the distilled water (best recently boiled to expel air, and cooled) until it is saturated. The solution should not be made in large quantities at a time, as it gradually decomposes, depositing sulphur and losing its smell; it is then unfit for use.



2. *Sulphurous Acid Solution* is prepared by passing sulphur dioxide gas into distilled water until it is saturated. The gas is made by heating scraps of copper with strong  $H_2SO_4$  in a glass flask fitted as is shown in A (Fig. 45). The gas passes from the flask through some water contained in a washing-bottle as shown at B (Fig. 45), and from this bottle into the distilled water (Fig. 44) through the tube *a b*, which is connected with the outlet tube of the washing bottle by means of the india-rubber joint *c*. The solution must be kept in a well-stoppered bottle.

3. *Chlorine-water*.—Chlorine gas is made in the apparatus shown in Fig. 45, by gently heating manganic oxide

(in lumps or powder) in the flask A with some strong commercial hydrochloric acid, previously mixed with a third its volume of water. The gas bubbles through a little water in the washing-bottle B, and thence through the bent tube *a b* into the distilled water (Fig. 44).

FIG. 45.





FIG. 46.




This solution must be kept in a dark place, or in a bottle which is coated with black paper, as it undergoes alteration by light.

4. *Hydrofluosilicic Acid* is made by passing silicium fluoride gas into water. The gas is prepared in a flask fitted as shown in A, Fig. 45. Into this flask there is first poured an intimate mixture of 50 grammes of dry, pure, finely-powdered fluor-spar; 300 grammes of strong  $\text{H}_2\text{SO}_4$  are then poured into the flask through the funnel-tube, and the acid is mixed with the powder by shaking it round in the flask; the gas is caused to be evolved by gently heating the flask, and is made to pass first through the bottle B (Fig. 45), which must be empty and dry inside; thence it escapes through a bent tube *a b* (Fig. 46), which is fitted upon the outlet tube of B by an india-rubber joint: the end of the tube, *a b*, is made to dip into mercury contained in a small beaker *d* (Fig. 46), which stands in a large beaker containing 400 c.c. of water. As soon as the silicium fluoride gas comes into the water, after escaping from the mercury, it is

decomposed into hydrofluosilicic acid, which dissolves in the water, and silica, which remains suspended in the water as a gelatinous mass; the silica very soon closes the end of the delivery-tube *a b*, unless it is kept immersed in the mercury. As the current of gas slackens the heat is raised, until white fumes of  $\text{H}_2\text{SO}_4$  begin to appear in the preparation flask; the process is then arrested, and the gelatinous silica is separated from the solution by squeezing the latter through fine muslin and afterwards filtering it if not quite clear. The silica may be dried, heated strongly in a porcelain dish, and put by in a stoppered bottle as reagent No. 86 (526), for which it serves admirably.

5. *Nitrogen-tetroxide Solution*.—Lead nitrate in dry pieces is heated in a test-tube fitted with cork and delivery-tube (Fig. 47), and the red fumes which escape are passed into dilute  $\text{H}_2\text{SO}_4$ . Care must be taken to maintain the heat uniformly after the fumes commence to be evolved, else the liquid may be sucked back into the hot tube by the contraction of the gas on cooling; also the delivery-tube must be removed from the liquid as soon as the heating is stopped.

FIG. 47.





## 529.—SOLUTIONS FOR THE

*Note.*—Each bottle should bear on its label the number, name, and ties should be kept near the solutions containing the powdered solid these bottles should also carry on their label the name, number, and

Reference number.	Name. [The numbers in brackets refer to similarly numbered remarks in the last column.]	Formula.
100. s.....	Potassium chloride.....	KCl.....
101. s.....	Ammonium chloride.....	AmCl.....
102. s.....	Sodium chloride.....	NaCl.....
103. s.....	Magnesium sulphate.....	MgSO <sub>4</sub> .7H <sub>2</sub> O.....
104. s.....	Barium chloride.....	BaCl <sub>2</sub> .2H <sub>2</sub> O.....
105. s.....	Strontium nitrate.....	Sr(NO <sub>3</sub> ) <sub>2</sub> .4H <sub>2</sub> O.....
106. s.....	Calcium chloride.....	CaCl <sub>2</sub> .6H <sub>2</sub> O.....
107. s.....	{ Alum, or Aluminium potassium sulphate }.....	AlK(SO <sub>4</sub> ) <sub>2</sub> .12H <sub>2</sub> O.....
108. s.....	Ferric chloride (α)*.....	Fe <sub>2</sub> Cl <sub>3</sub> .....
109. s.....	Ferrous sulphate (α)*.....	FeSO <sub>4</sub> .7H <sub>2</sub> O.....
110. s.....	{ Chrome alum, or Chromium potassium sulphate }.....	CrK(SO <sub>4</sub> ) <sub>2</sub> .12H <sub>2</sub> O.....
111. s.....	Zinc sulphate.....	ZnSO <sub>4</sub> .7H <sub>2</sub> O.....
112. s.....	Manganese chloride (α)*.....	MnCl <sub>2</sub> .....
113. s.....	Nickel sulphate.....	NiSO <sub>4</sub> .7H <sub>2</sub> O.....
114. s.....	Cobalt nitrate.....	Co(NO <sub>3</sub> ) <sub>2</sub> .6H <sub>2</sub> O.....
115. s.....	Mercuric chloride.....	HgCl <sub>2</sub> .....
116. s.....	Lead acetate (α)*.....	PbAc <sub>2</sub> .3H <sub>2</sub> O.....
117. s.....	Bismuth nitrate (α)*.....	Bi(NO <sub>3</sub> ) <sub>3</sub> .5H <sub>2</sub> O.....
118. s.....	Copper sulphate (α)*.....	CuSO <sub>4</sub> .6H <sub>2</sub> O.....
119. s.....	Cadmium sulphate (α)*.....	CdSO <sub>4</sub> .4H <sub>2</sub> O.....
120. s.....	{ Arsenious oxide (α) (†) } Solution in dilute HCl.....	As <sub>2</sub> O <sub>3</sub> .....
120. s.....	{ Arsenious oxide (α) (†) } Solution in water.....	—
121. s.....	Sodium arsenate.....	Na <sub>2</sub> HAsO <sub>4</sub> .12H <sub>2</sub> O.....
122. s.....	Antimonious chloride (α)*, †.....	SbCl <sub>3</sub> .....
123. s.....	Stannous chloride (α)*.....	SnCl <sub>2</sub> .2H <sub>2</sub> O.....
124. s.....	Stannic chloride (α)*.....	SnCl <sub>4</sub> .....
125. s.....	Silver nitrate.....	AgNO <sub>3</sub> .....
126. s.....	Mercurous nitrate (α)*.....	Hg <sub>2</sub> (NO <sub>3</sub> ) <sub>2</sub> .H <sub>2</sub> O.....

\* α signifies that the solid substance is not  
† Solid Sb<sub>2</sub>O<sub>3</sub> or K(SbO)T (Tartar emetic)

## REACTIONS OF THE METALS.

formula of the substance. Small (4 or 6 ounce) wide-mouthed bot-  
of each of those substances whose number has a small (s) affixed to it;  
formula of the substance.

Weight of solid in grammes to be dissolved in one "Winches- ter."	Proportion by weight of solid to water.	Remarks.  [These numbers correspond to the small reference numbers in brackets in the second column.]
100 .....	1 : 25	48. The solution must be made acid with a little $\text{H}_2\text{SO}_4$ , and some clean iron nails kept in it.
200 .....	1 : 12	
50 .....	1 : 50	49. A little $\text{H}\bar{\text{A}}$ must be added to this solution to make it clear.
50 .....	1 : 50	
25 .....	1 : 100	
50 .....	1 : 50	50. 25 grammes of $\text{Bi}(\text{NO}_3)_3$ must be dissolved by heat- ing with 25 c.c. of strong $\text{HCl}$ diluted with 25 of water; this solution is cooled and poured into the bottle, and the measure made up by pouring in dilute $\text{HCl}$ containing one-twentieth of its volume of strong $\text{HCl}$ .
200 (in crystals) .....	1 : 12	
200 .....	1 : 12	51. Dissolve 10 grammes of $\text{As}_2\text{O}_3$ by heating it with 50 c.c. of strong $\text{HCl}$ mixed with 50 c.c. of water, and dilute to the quart.†
25 .....	1 : 100	
25 .....	1 : 100	52. Boil excess of $\text{As}_2\text{O}_3$ for several minutes with 200 c.c. of water, filter, and dilute to the quart.†
50 .....	1 : 50	
25 .....	1 : 100	53. 25 grains of crystallized $\text{SbCl}_3$ are dissolved in 25 c.c. of strong $\text{HCl}$ mixed with 25 c.c. of water, and diluted to the quart; with $\text{HCl}$ mixed with four times its measure of water.
50 .....	1 : 50	
50 .....	1 : 50	54. Boil 25 grammes of $\text{SnCl}_2$ with 50 c.c. of strong $\text{HCl}$ mixed with 50 c.c. of water, as soon as it has dissolved to a clear solution dilute to the quart.† Scraps of $\text{Sn}$ must be kept in the bottle.
25 .....	1 : 100	
25 .....	1 : 100	55. Heat 25 grammes of $\text{SnCl}_2$ with 25 c.c. of strong $\text{HCl}$ and 100 c.c. of water; whilst constantly stirring, add $\text{KClO}_3$ to the hot solution until the liquid turns yellow and $\text{Cl}$ is smelt, boil off the $\text{Cl}$ and dilute to the quart.†
25 .....	1 : 100	
25 .....	1 : 100	56. Dissolve by warming 25 grammes of $\text{Hg}_2(\text{NO}_3)_2$ with 6 c.c. of strong $\text{HNO}_3$ diluted with 114 c.c. of water, then pour water into this solution until it is diluted to a quart.† Keep $\text{Hg}$ in the bottom of the bottle.
25 .....	1 : 100	

in the crystalline condition.  
may be used for the blowpipe reaction.

† Winchester quart.

## 530.—SOLUTIONS FOR THE REACTIONS OF THE ACID RADICLES.

[Note.—Refer to note at the head of the preceding Table (529), which is also applicable to this. Many of the solutions in this list are already prepared amongst the reagents (524, 525, 526), and the solutions for reactions for metals (529).]

Reference number.	Name. [The small numbers in brackets refer to corresponding numbers in the sixth column.]	Formula.	Weight of solid in grammes to be dissolved in one Winchester quart.	Proportion by weight of solid to water.	Remarks.
130.	Sodium sulphate.....	$\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ .....	25	1 : 100	
131. S.	Sodium carbonate ( <sup>1</sup> ).....	$\text{Na}_2\text{CO}_3$ .....	—	—	57. The $\text{Na}_2\text{CO}_3$ is required only in the solid form, and can be taken from the reagent bottle on the working bench, No. 18 (524).
132. S.	Marble (calcium carbonate) ( <sup>2</sup> ).....	$\text{CaCO}_3$ .....	—	—	The marble is kept in small pieces as large as a pea; it is found in reagent bottle No. 79 (526).
133. S.	Ferrous sulphide ( <sup>3</sup> ).....	$\text{FeS}$ .....	—	—	
134.	Ammonium sulphide ( <sup>4</sup> ).....	$\text{Am}_2\text{S}$ .....	100	1 : 25	58. The $\text{FeS}$ is kept as a solid in small pieces as large as a pea.
135.	Acid sodium sulphite.....	$\text{NaHSO}_3$ .....	100	1 : 25	
136. S.	Sodium hypochlorite ( <sup>5</sup> ).....	$\text{NaOCl}$ .....	100	1 : 25	The $\text{Am}_2\text{S}$ can be taken from reagent bottle No. 7, on the working bench.
137. S.	Potassium nitrate ( <sup>6</sup> ).....	$\text{KNO}_3$ .....	100	1 : 25	
138. S.	Potassium nitrite ( <sup>7</sup> ).....	$\text{KNO}_2$ .....	50	1 : 50	59. Strong "Liquor Sodæ Chlorinatæ," diluted with an equal measure of water.
139. S.	Potassium chlorate ( <sup>8</sup> ).....	$\text{KClO}_3$ .....	50	1 : 50	
140. S.	Sodium chloride ( <sup>9</sup> ).....	$\text{NaCl}$ .....	50	1 : 50	60. No. 102, paragraph 529.
141. S.	Potassium iodide.....	$\text{KI}$ .....	12	1 : 200	61. No. 121, paragraph 529.
142.	Sodium phosphate.....	$\text{N H}_2\text{PO}_4 \cdot 12\text{H}_2\text{O}$ .....	50	1 : 50	62. No. 19, paragraph 524.
143.	Sodium arsenate ( <sup>10</sup> ).....	$\text{Na}_2\text{HAsO}_4 \cdot 10\text{H}_2\text{O}$ .....	50	1 : 50	
144. S.	Borax ( <sup>11</sup> ).....	$\text{K}_2\text{B}_4\text{O}_7$ .....	25	1 : 100	63. The solution is made by dissolving 100 grammes of the thick syrup, sold as "soluble glass," in water, and diluting to a quart.* The solid substance to be used is sand finely ground.
145.	Potassium bichromate.....	$\text{K}_2\text{Cr}_2\text{O}_7$ .....	—	—	
146. S.	Sodium silicate ( <sup>12</sup> ).....	$\text{Na}_2\text{SiO}_3$ .....	—	—	64. No. 35, paragraph 525.
147. S.	Silica (white sand).....	$\text{SiO}_2$ .....	—	—	65. No. 42, paragraph 525.
148.	Fluor-spar (calcium fluoride).....	$\text{CaF}_2$ .....	—	—	66. No. 4, paragraph 524.
149. S.	Hydrofluosilicic acid ( <sup>13</sup> ).....	$\text{H}_2\text{SiF}_6$ .....	50	1 : 50	
150.	Potassium cyanide ( <sup>14</sup> ).....	$\text{KCN}$ .....	25	1 : 100	67. No. 30, paragraph 525.
151.	Potassium ferrocyanide.....	$\text{K}_4\text{Fe(CN)}_6 \cdot 3\text{H}_2\text{O}$ .....	25	1 : 100	
152.	Potassium ferricyanide.....	$\text{K}_3\text{Fe(CN)}_6$ .....	25	1 : 25	68. 120 grms. of $\text{H}_2\text{T}$ are dissolved by heat in 500 c.c. of water, the hot solution is exactly neutralized with solid $\text{Na}_2\text{CO}_3$ , and then diluted to a quart.*
153. S.	Potassium sulphocyanide.....	$\text{Am}_2\text{CS}_2 \cdot 2\text{H}_2\text{O}$ .....	100	1 : 10	69. A little of the solid substance (see No. 41, 525) is dissolved, when required, in water.
154. S.	Tartaric acid ( <sup>15</sup> ).....	$\text{H}_2\text{T}$ .....	—	—	
155. S.	Sodium tartrate ( <sup>16</sup> ).....	$\text{Na}_2\text{T} \cdot 2\text{H}_2\text{O}$ .....	—	—	
	Sodium acetate ( <sup>17</sup> ).....	$\text{NaAc} \cdot 3\text{H}_2\text{O}$ .....	—	—	

\* Winchester quart.

## 531. CHEMICALS REQUIRED FOR SECTION I.

In the following list those substances which are used for Section I only are marked by letters of the alphabet. The bottles containing these substances should be arranged apart in alphabetical order. Against others a number is placed; this indicates that the substance is used for analytical purposes, and will therefore be found, together with any necessary descriptive remarks, in one of the foregoing lists (524, 525, 526, 529, 530). [See note at head of Table (532).]

Letter or No. of reference.	Name.	Formula.	Remarks.
A.	Mercuric oxide.....	HgO	The orange-colored <i>precipitated</i> oxide is best.
20.	Potassium chlorate.....	KClO <sub>3</sub>	
B.	Manganese dioxide.....	MnO <sub>2</sub>	In powder. No. 66 would serve the purpose, but it is unnecessary that the substance should be pure.
C.	Wood-charcoal.....	C	In pieces as large as a hazelnut.
57.	Lime-water.....	Ca(OH) <sub>2</sub>	
D.	Sulphur.....	S	Roll sulphur broken into pieces as large as a pea.
21.	{ Litmus and turmeric }	—	—
	{ papers..... }		
59.	Potassium dichromate.....	K <sub>2</sub> Cr <sub>2</sub> O <sub>7</sub>	
E.	Phosphorus.....	P	In small pieces as large as half a pea, kept in a stoppered bottle under water.
F.	Granulated zinc.....	Zn	No. 46 would serve the purpose, but it is not necessary that this zinc should be free from As.
G.	Hydrochloric acid.....	HCl	Strong, commercial.
H.	Marble.....	CaCO <sub>3</sub>	In pieces not larger than a hazelnut.
I.	Copper.....	Cu	In any form— <i>e. g.</i> , turnings, clippings, filings, etc.
J.	Nitric acid.....	HNO <sub>3</sub>	Strong, commercial.
K.	{ Ammonium chloride, }	NH <sub>4</sub> Cl	{ Solid, commercial. The solid (No. 101) will also serve; but for this purpose it need not be pure.
	{ or Sal-ammoniac..... }		
32.	Slaked lime.....	Ca(OH) <sub>2</sub>	
L.	Oxalic acid.....	H <sub>2</sub> C <sub>2</sub> O <sub>4</sub> .2H <sub>2</sub> O	Solid, in crystals.
M.	Sulphuric acid.....	H <sub>2</sub> SO <sub>4</sub>	Strong, commercial.
N.	{ Sodium hydrate, or }	NaHO	{ Strong solution of the commercial solid.
	{ caustic soda..... }		
O.	"Turkey red".....	—	In strips.
P.	{ Sodium chloride, or }	NaCl	{ In pieces as large as a hazelnut broken from a lump of rock salt or of fused table salt.
	{ common salt..... }		
3.	Nitric acid.....	HNO <sub>3</sub>	
48.	Silver nitrate.....	AgNO <sub>3</sub>	
6.	Ammonia.....	NH <sub>3</sub> .HO	
9.	Ammonium oxalate.....	(NH <sub>4</sub> ) <sub>2</sub> C <sub>2</sub> O <sub>4</sub>	
17.	Barium chloride.....	BaCl <sub>2</sub>	
7.	Ammonium sulphide.....	Am <sub>2</sub> S	
39.	Potassium nitrate.....	KNO <sub>3</sub>	

### 532. CHEMICALS REQUIRED FOR THE EXPERIMENTS ON ANALYTICAL OPERATIONS (SECTION III).

All the substances, with those exceptions\* only which are named in this list, are required either for analytical purposes (see lists in paragraphs 524, 525, 526, 529, 530), where they may be found by the number placed in the first column; or they are already entered on the list of substances required for experiments on the gases (531), where they may be found by the reference-letter in the first column. A letter (s) affixed to the number of reference indicates that the substance is required in the solid condition (see note at heading of 529).

*Note.*—All numbers below 22 refer to bottles standing on the shelves above the working bench.

If a substance is required for more than one experiment, its entry is not repeated.

Number or letter of reference.	Name.	Formula.
39.	Potassium nitrate.....	$\text{KNO}_3$ .
118. s.	Copper sulphate (solid).....	$\text{CuSO}_4 \cdot 6\text{H}_2\text{O}$ .
79.	Marble.....	$\text{CaCO}_3$ .
2.	Hydrochloric acid.....	$\text{HCl}$ .
45.	Copper.....	$\text{Cu}$ .
3.	Nitric acid.....	$\text{HNO}_3$ .
17.	Barium chloride.....	$\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$ .
8.	Ammonium carbonate.....	$\text{Am}_2\text{CO}_3$ .
161. *	Filter-paper.....	In sheet.
1.	Sulphuric acid.....	$\text{H}_2\text{SO}_4$ .
A.	Mercuric oxide.....	$\text{HgO}$ .
K.	Ammonium chloride.....	$\text{NH}_4\text{Cl}$ .
100. s.	Potassium chloride.....	$\text{KCl}$ .
18.	Sodium carbonate.....	$\text{Na}_2\text{CO}_3$ .
112. s.	Manganese chloride.....	$\text{MnCl}_2$ .
19.	Borax.....	$\text{Na}_2\text{B}_4\text{O}_7$ .
62. s.	Ferrous sulphate.....	$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ .
102.	Sodium chloride.....	$\text{NaCl}$ .
160. *	Wood-charcoal.....	C.
116. s.	Lead acetate (solid).....	$\text{PbAc}_2 \cdot 3\text{H}_2\text{O}$ .
—	Red cabbage leaves.....	—
2.	Hydrochloric acid.....	$\text{HCl}$ .
6.	Ammonia.....	$\text{NH}_4\text{HO}$ .
5.	Ammonium chloride.....	$\text{NH}_4\text{Cl}$ .
21.	{ Litmus-paper.....	—
	{ Turmeric-paper.....	—

\* The exceptions are Nos. 160 and 161, which numbers refer to paragraph 533; also the red cabbage leaves, which are readily obtainable.

**533. SUNDRY OTHER REQUISITES NOT INCLUDED IN THE PRECEDING LISTS.**

Reference number.	Name and description.
160.	<i>Wood-charcoal.</i> —This may be purchased in small pieces commonly used for fuel. If required for blowpipe experiments (38), the charcoal should be tolerably free from cracks and from bark. It may be sawn into convenient shape and size by a fine-toothed saw.
161.	<i>Filter-paper</i> may be bought in sheets which are cut into squares of the required size: these are folded and cut as directed in experiment 40 (p. 58). Ready cut circular filters can also be purchased, which only require to be folded.
162.	<i>Corks.</i> —These should be obtained as free as possible from holes or cracks; they must be sound and easily softened by pressure. All sizes between $1\frac{1}{2}$ inch and $\frac{1}{2}$ inch in diameter should be kept in stock. For wash-bottles and other permanently fitted apparatus in which the cork is not subjected to much heating, the india-rubber stoppers are much to be preferred to corks, since they are almost imperishable, and retain their elasticity very much longer than corks do.

**SUBSTANCES TO BE GIVEN FOR ANALYSIS BY THE PRECEDING TABLES AND DIRECTIONS.**

**534.** It must be understood that the substances mentioned below are merely brought forward as examples of what may be given to the student for analysis; the teacher will use his own judgment in adopting or extending the list.

**535. SUBSTANCES FOR ANALYSIS WHILST TRYING THROUGH THE REACTIONS FOR METALS AND ACID RADICLES.**

If the student, after trying through the reactions for each of the metallic groups, intends to analyze substances containing one member only, or any two or more members, of that group, it is a good plan to give him first a few substances containing only one member and to

gradually increase the number present in those subsequently given. It is well to give these substances in the dissolved state\* as a rule, as they are intended mainly to afford practice in separation and detection according to the group-tables, and the time spent by the student in preparing the solution is therefore wasted. Occasionally, however, a solid substance may be given in which the metal or acid radicle present is to be detected by blowpipe tests or other tests made on the solid substance.

As examples of the substances to be given, in the order best suited to the gradual advance in difficulty, the following selected for Groups V, IV, and III B will serve:

<i>Group V.</i>	<i>Group IV.</i>	<i>Group III b.</i>
Solution of NaCl	Solution of $\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$	Solution of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$
" $\text{NH}_4\text{Cl}$	" $\text{Sr}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$	" $\text{MnCl}_2$
" KCl	" $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$	" $\text{NiSO}_4 \cdot 7\text{H}_2\text{O}$
" $\text{NaCl} + \text{KCl}$	" $\text{Sr}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O} +$	" $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$
" $\text{NaCl} + \text{NH}_4\text{Cl} +$	$\text{CaCl}_2 \cdot 6\text{H}_2\text{O} +$	" $\text{MnCl}_2 + \text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$
KCl	" $\text{BaCl}_2 \cdot 2\text{H}_2\text{O} +$	" $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O} +$
" $\text{NaCl} + \text{NH}_4\text{Cl}$	$\text{Sr}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O} +$	" $\text{NiSO}_4 \cdot 7\text{H}_2\text{O} +$
" $\text{MgSO}_4 \cdot 7\text{H}_2\text{O} +$	$\text{CaCl}_2 \cdot 6\text{H}_2\text{O} +$	$\text{ZnSO}_4 \cdot 7\text{H}_2\text{O} +$
$\text{NaCl} + \text{KCl}$	" $\text{BaCl}_2 \cdot 2\text{H}_2\text{O} +$	etc., etc., etc.
<i>Solids:</i> — $\text{NaCl}$ , $\text{NH}_4\text{Cl}$ , KCl, $\text{AmCl} + \text{NCl}$ , etc.	" $\text{Sr}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$	<i>Solids:</i> — $\text{MnCl}_2$ , $\text{NiSO}_4 \cdot 7\text{H}_2\text{O}$ $\text{Co}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$ , $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ , etc.

536. Substances are given to be tested for the members of an acid-radicle group in the same order (535),\* the first substances containing one member only, those subsequently given containing two or more members of the group.

### 537. SUBSTANCES FOR ANALYSIS BY THE DIRECTIONS CONTAINED IN SECTION V.

Any one of the solutions named in paragraphs 529 and 530 may be given for analysis by Section V, some of them being neutral, some acid, and others alkaline

\* The solutions already made for the reactions, paragraphs 529 and 530, will serve very well for this purpose, being given either singly or mixed.

in reaction. The following more difficult solutions with acid or alkaline reaction may also be added to the list:

Acid.	Alkaline.
Ca <sub>3</sub> (PO <sub>4</sub> ) <sub>2</sub> ("Bone-ash") dissolved in dilute HCl.	KSbO <sub>3</sub> dissolved in water.
BaCO <sub>3</sub> dissolved in dilute HCl.	Na <sub>2</sub> SiO <sub>3</sub> " " (solution of "soluble glass").
Mu(Bi <sub>2</sub> ) <sub>2</sub> * " " HCl.	Na <sub>2</sub> SnO <sub>3</sub> " " ("preparing salt" of the dyer).
BaCrO <sub>4</sub> " " HCl.	

Any of the solid salts which were dissolved in order to prepare the solutions in paragraphs 529 and 530, may be given for analysis as solid substances by (331 *et seq.*).

As examples of substances possessing metallic appearance (366), the following may be mentioned as suitable for analysis:

Zn, in pieces or filings.	NiAs (Kupfernickel).
Pb, " "	Graphite (Black lead).
FeS <sub>2</sub> (Iron Pyrites).	Iron filings.

As insoluble substances which require to be examined by (367 *et seq.*), the following may be given for analysis:

BaSO <sub>4</sub> .	CaF <sub>2</sub> .
PbSO <sub>4</sub> .	FeCr <sub>2</sub> O <sub>4</sub> (Chrome Iron Ore).
AgCl.	S (as pieces of roll sulphur).
SnO <sub>2</sub> (Tinestone).	C (as powdered wood-charcoal or plumbago).

### 538. SOLID SUBSTANCES TO BE ANALYZED BY THE PRELIMINARY EXAMINATIONS.

For the Preliminary Examinations for Metals (387):

Simple.			Complex.	
NH <sub>4</sub> Cl	Pb(NO <sub>3</sub> ) <sub>2</sub>	ZnSO <sub>4</sub> ·7H <sub>2</sub> O	AmCl + NaCl	AlK(SO <sub>4</sub> ) <sub>3</sub> ·12H <sub>2</sub> O
Hg <sub>2</sub> I <sub>2</sub>	NaCl	MnCl <sub>2</sub>	HgCl <sub>2</sub> + BaCl <sub>2</sub> ·2H <sub>2</sub> O	SnO <sub>2</sub> + Sr(NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O
As <sub>2</sub> O <sub>3</sub>	BaCl <sub>2</sub> ·2H <sub>2</sub> O	Co(NO <sub>3</sub> ) <sub>2</sub> ·6H <sub>2</sub> O	MnCl <sub>2</sub> + KCl	CaCO <sub>3</sub> + ZnSO <sub>4</sub> ·7H <sub>2</sub> O
KNO <sub>3</sub>	Sr(NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O	NiSO <sub>4</sub> ·7H <sub>2</sub> O	NiSO <sub>4</sub> ·7H <sub>2</sub> O + ZnSO <sub>4</sub> ·7H <sub>2</sub> O	Cr <sub>2</sub> O <sub>3</sub> + MnCl <sub>2</sub> + NaCl.
Cr <sub>2</sub> O <sub>3</sub>	CuSO <sub>4</sub> ·6H <sub>2</sub> O (See note.)	SnO <sub>2</sub>		

\* Made by mixing hot solutions of Na<sub>2</sub>B<sub>4</sub>O<sub>7</sub> and Mg(NO<sub>3</sub>)<sub>2</sub>.



*Note.*—Best finely powdered, since its color is then almost destroyed.

For the Preliminary Examination for Acid Radicles (410):

Simple.			Complex.
CaCO <sub>3</sub> Na <sub>2</sub> SO <sub>4</sub> ·7H <sub>2</sub> O CaCl <sub>2</sub> ·O NaCl	FeS KNO <sub>3</sub> KClO <sub>3</sub> KI	Na <sub>2</sub> S <sub>2</sub> O <sub>3</sub> ·5H <sub>2</sub> O KBr NaA	CaCO <sub>3</sub> + KNO <sub>3</sub> CaCl <sub>2</sub> O + CaF <sub>2</sub> Na <sub>2</sub> CO <sub>3</sub> + KI.

A few of the above-mentioned solids may then be examined by both Preliminary Examinations for both Metals and Acid Radicles.

### 539. SUBSTANCES TO BE ANALYZED BY THE GENERAL COURSE (380 *et seq.*).

The following lists are so arranged that the analysis of the substances contained in them is more difficult in each column proceeding from left to right, and usually also in proceeding in one and the same column from top to bottom. For examples of Alkaline Solutions, see (537):

### SUBSTANCES FOR ANALYSIS BY THE GENERAL COURSE.

1. Simple soluble substances.	2. Complex soluble substances.	3. Complex soluble substances.
BaCl <sub>2</sub> ·2H <sub>2</sub> O MnCl <sub>2</sub> HgCl <sub>2</sub> MgSO <sub>4</sub> ·7H <sub>2</sub> O KCl NiSO <sub>4</sub> ·7H <sub>2</sub> O *CaCO <sub>3</sub> Pb(NO <sub>3</sub> ) <sub>2</sub> *As <sub>2</sub> O <sub>3</sub> Fe <sub>2</sub> Cl <sub>3</sub> NH <sub>4</sub> Cl NaCl etc., etc.	CaCl <sub>2</sub> ·6H <sub>2</sub> O + BaCl <sub>2</sub> ·2H <sub>2</sub> O + Sr(NO <sub>3</sub> ) <sub>2</sub> ·4H <sub>2</sub> O AlK(SO <sub>4</sub> ) <sub>2</sub> ·12H <sub>2</sub> O + CrK(SO <sub>4</sub> ) <sub>2</sub> MnCl <sub>2</sub> + ZnSO <sub>4</sub> ·7H <sub>2</sub> O + CuSO <sub>4</sub> ·6H <sub>2</sub> O Pb(NO <sub>3</sub> ) <sub>2</sub> + HgCl <sub>2</sub> + NaCl *As <sub>2</sub> O <sub>3</sub> + CaCO <sub>3</sub> + BaCO <sub>3</sub> *ZnO + MgCO <sub>3</sub> + KClO <sub>3</sub> *MgCO <sub>3</sub> + BaCO <sub>3</sub> + PbCO <sub>3</sub> etc., etc.	<div style="display: flex; align-items: center;"> <div style="font-size: 2em; margin-right: 5px;">{</div> <div> Fe<sub>2</sub>Cl<sub>3</sub> MnCl<sub>2</sub> ZnSO<sub>4</sub>·7H<sub>2</sub>O KCl As<sub>2</sub>O<sub>3</sub> CrK(SO<sub>4</sub>)<sub>2</sub>·12H<sub>2</sub>O Co(NO<sub>3</sub>)<sub>2</sub>·6H<sub>2</sub>O Hg<sub>2</sub>(NO<sub>3</sub>)<sub>2</sub>·4H<sub>2</sub>O Sr(NO<sub>3</sub>)<sub>2</sub>·4H<sub>2</sub>O KNO<sub>3</sub> Bi(NO<sub>3</sub>)<sub>3</sub>·5H<sub>2</sub>O etc., etc. </div> </div>

\* To be given in the solid state.

4. Substances yielding a phosphate precipi- tate in Group III.	5. Substances partly or entirely insoluble.	
$\left\{ \begin{array}{l} \text{CaCl}_2, 6\text{H}_2\text{O} \\ \text{Fe}_2\text{Cl}_6 \\ \text{N H}_2\text{PO}_4, 12\text{H}_2\text{O} \end{array} \right.$  $\left\{ \begin{array}{l} \text{BaCl}_2, 2\text{H}_2\text{O} \\ \text{MgCl}_2 \\ \text{Fe}_2\text{Cl}_6 \\ \text{Na}_2\text{H}_2\text{PO}_4, \text{H}_2\text{O} \end{array} \right.$  $\left\{ \begin{array}{l} \text{Co(NO}_3)_2, 6\text{H}_2\text{O} \\ \text{CaCl}_2, 6\text{H}_2\text{O} \\ \text{MnCl}_2 \\ \text{CrK SO}_4, 12\text{H}_2\text{O} \\ \text{Na}_2\text{H}_2\text{PO}_4, \text{H}_2\text{O} \\ \text{etc., etc.} \end{array} \right.$	$\text{BaSO}_4$ $\text{SiO}_2 + \text{S}$ $\text{BaSO}_4 + \text{CaF}_2 + \text{SnO}_2$  $\left\{ \begin{array}{l} \text{BaCl}_2, 2\text{H}_2\text{O} \\ \text{AlK(SO}_4)_3, 12\text{H}_2\text{O} \\ \text{AgNO}_3 \end{array} \right.$  $\left\{ \begin{array}{l} \text{SiO}_2 \\ \text{S} \\ \text{C} \end{array} \right.$  $\text{PbSO}_4 + \text{BaSO}_4$ $\text{PbCrO}_4$ (ignited)	<i>Metallic Substances (467 et seq.)</i> Iron pyrites ( $\text{FeS}_2$ ) Iron filings ( $\text{Fe}$ ) Zinc clippings ( $\text{Zn}$ ) Brass filings ( $\text{Cu} + \text{Zn}$ ) German silver ( $\text{Cu} + \text{Zn} + \text{Ni}$ ) Bronze ( $\text{Cu} + \text{Sn}$ ) Type-metal ( $\text{Sb} + \text{Sn} + \text{Pb}$ )  <hr/> <i>Silicates (480)—</i> Fire-clay ( $\text{Al}_2\text{O}_3, 2\text{SiO}_2$ ) Brown-clay ( " + $\text{Fe}$ ) Window-glass ( $\text{Na}, \text{Ca}, \text{SiO}_2$ ) Flint-glass ( $\text{Pb}, \text{K}, \text{SiO}_2$ )  <hr/> <i>Cyanogen Compounds (482)—</i> Any of the foregoing which has been mixed with $\text{KC}_y$ , $\text{KC}_y\text{S}$ , $\text{K}_4\text{FeCy}_6, 3\text{H}_2\text{O}$ , $\text{K}_4\text{Fe}_2\text{Cy}_{12}$ , $\text{K}_4\text{Co}_2\text{Cy}_{12}$

## APPENDIX I.

---

### REACTIONS FOR THE RARER ELEMENTS.

THIS Appendix contains some of the most important reactions for many of the rarer elements. The arrangement here used is that adopted in Section IV, those elements being placed together which are precipitated in the same analytical group: the groups are arranged in the order in which they occur in the General Table. A scheme for the detection of these elements follows (553), showing in which group they are precipitated in the General Table, and by which reactions they will be most readily found.

---

#### GROUP I.—SILVER GROUP.

In this group are included Tl and Wo: the former is partially precipitated as chloride by HCl, the latter completely as tungstic acid.

Thallium is only partially precipitated in Group I, since its chloride is not quite insoluble in water: it belongs also to Group III B, being entirely precipitated by  $\text{Am}_2\text{S}$ .

---

#### 540. THALLIUM (Tl).—Use $\text{Tl}_2\text{SO}_4$ solution.

Tl occurs in small quantity in many natural sulphides, often also in the ashes of plants and in mineral waters.

Tl yields both thallic and thallous salts, but the former are very unstable, changing even when their solutions are heated into thallous salts.

*HCl*: a white precipitate, which rapidly settles, does not blacken in the light, and is soluble in aqua regia. It is soluble in a large quantity of water, and therefore does not form in dilute solutions.

**KI:** a yellow precipitate; almost insoluble in water, more soluble in KI solution. In a solution containing Fe, any ferric salt must first be reduced by  $\text{H}_2\text{SO}_2$  before adding KI.

**$\text{PtCl}_4$ :** orange-red precipitate, slightly soluble in water.

**$\text{Am}_2\text{S}$ :** black precipitate, which is easily coagulated by heat, is insoluble in  $\text{AmHO}$ , alkaline sulphides, and in  $\text{KCy}$ : it is readily oxidized by the air to  $\text{Ti}_2\text{SO}_4$ , and is easily soluble in mineral acids.  $\text{Ti}$  is entirely precipitated by  $\text{H}_2\text{S}$  from a solution in which  $\text{H}\bar{\text{A}}$  is the only free acid present, but free mineral acids prevent the precipitation entirely.

**Flame coloration.**—Thallium compounds impart to the Bunsen flame an intense *green* color, which, however, rapidly disappears. The spectrum (p. 360) is very characteristic, consisting of one bright emerald-green line. Thallium may usually be easily detected by the spectroscopic in solution, or better in any of its precipitates mentioned above.

#### 541. TUNGSTATES.—Use $\text{Na}_2\text{WO}_4$ solution.

$\text{Wo}$  usually occurs in the form of a tungstate. The insoluble tungstates yield soluble alkaline tungstates on being fused with alkaline carbonates. From a solution of alkaline tungstate  $\text{HCl}$  precipitates the tungstic acid entirely.

**$\text{HCl}$**  ( $\text{HNO}_3$  or  $\text{H}_2\text{SO}_4$ ), white precipitate ( $\text{H}_2\text{WO}_4$ ), becoming yellow on boiling: insoluble in excess of acid, but soluble in  $\text{AmHO}$ . A piece of  $\text{Zn}$  dropped into the acid liquid containing the precipitate yields a deep blue color.

**$\text{SnCl}_2$**  (a neutral solution made by dissolving  $\text{SnCl}_2$  crystals in water and filtering): *yellow* precipitate, becoming *blue* on addition of  $\text{HCl}$  and heating.  $\text{Am}_2\text{S}$  yields no precipitate in a solution of alkaline tungstate; but if, after adding  $\text{Am}_2\text{S}$ , the liquid is made acid with  $\text{HCl}$ , brown  $\text{WS}_3$  is precipitated.

**Microcosmic bead:**—

In outer flame—*Colorless* or *yellow*.

In inner flame—*Blue*: if a little  $\text{FeSO}_4$  is fused into the bead the color changes to *blood-red*. These colors are best seen when the bead is perfectly cold.

#### GROUP II A.—COPPER GROUP.

In this group are included  $\text{Pd}$  ( $\text{Os}$ ,  $\text{Rh}$ ,  $\text{Ru}$ ): they are precipitated as sulphides by  $\text{H}_2\text{S}$  from acid solutions, and the sulphides are insoluble in  $\text{Am}_2\text{S}$ , and in caustic alkali solution.

**542. PALLADIUM (Pd).—Use  $\text{PdCl}_2$  solution.**

Pd occurs as a metal in native platinum, also in gold and silver. Palladium solutions are reddish-brown, or yellow if dilute: addition of water precipitates a basic salt, unless sufficient free acid is present to prevent it.

$\text{H}_2\text{S}$ : a black pp., in neutral, alkaline, and acid solutions: this pp. is insoluble in  $\text{Am}_2\text{S}$ , but soluble in boiling  $\text{HCl}$  or in aqua regia.

$\text{AmHO}$ : flesh-colored pp. ( $\text{PdCl}_2 \cdot 2\text{NH}_3$ ); soluble in excess of  $\text{AmHO}$  to a colorless liquid, from which  $\text{HCl}$  precipitates yellow crystalline paladammonium-chloride ( $\text{N}_2\text{H}_4\text{Pd}''\text{Cl}_2$ ).

$\text{HgCy}_2$ : yellowish-white gelatinous pp. ( $\text{PdCy}_2$ ), slightly soluble in  $\text{HCl}$ , easily soluble in  $\text{AmHO}$ . Very characteristic reaction.

$\text{KI}$ : black pp. ( $\text{PdI}_2$ ). Very characteristic.

**GROUP II B.—ARSENIC GROUP.**

In this group are included Mo, Se, Te (Ir).

**543. MOLYBDATES ( $\text{Mo}$ ).—Use  $\text{Am}_2\text{MoO}_4$  solution.**

Mo occurs as a molybdate; also as sulphide, which may be readily converted into  $\text{MoO}_3$  by ignition in the air or heating with  $\text{HNO}_3$ . Unignited  $\text{MoO}_3$  dissolves in acids; ignited  $\text{MoO}_3$  is insoluble in acids, but easily soluble in alkalis.

$\text{HCl}$  ( $\text{HNO}_3$  or  $\text{H}_2\text{SO}_4$ ), added in small quantity to an aqueous solution of a molybdate, yields a precipitate which is readily soluble in excess of the acid.

$\text{H}_2\text{S}$ , added in *very small* quantity to the acidified solution gives a *blue* liquid, in larger quantities a *brown* pp. ( $\text{MoS}_3$ ); the precipitation becomes complete only when the solution is heated and  $\text{H}_2\text{S}$  passed for some time: the pp. is soluble in solutions of alkaline sulphides and hydrates, from which it is reprecipitated by acid added in excess.

$\text{Zn}$  or  $\text{SnCl}_2$ , added to a solution of a molybdate in  $\text{HCl}$ , colors it *brown*, *green*, or *blue*, according to its state of concentration.

$\text{KC}_2\text{S}$ , added to a solution acidified with  $\text{HCl}$ , gives no coloration; but on dropping in a piece of  $\text{Zn}$  a beautiful *crimson* color is produced, which, when the liquid is shaken with ether, is taken up by the ether.

$\text{Na}_2\text{HPO}_4$ , added in very small quantity to an acidified solution (best with  $\text{HNO}_3$ ) of a molybdate, gives, on gently warming the liquid, a yellow precipitate readily soluble in excess of alkali-hydrate solution.

*Borax bead*: outer flame, *yellow*; inner, *dark-brown*.

*Microcosmic bead*: outer and inner flames, *green*.

**544. SELENIUM (Se).**—Use a metallic selenide, an alkaline selenite and selenate.

Se occurs as metallic selenides (*e. g.*, of Fe, Cu, Ag). A selenide heated in an open tube evolves a smell of decaying horse-radish, and produces a gray or red sublimate of Se: the smell is very characteristic.

*a. Selenites:*

$H_2S$  gives in acid solutions, if cold a *yellow*, if hot a *reddish-yellow* pp.: soluble in  $Am_2S$ .

$BaCl_2$  in neutral solutions a white pp.: soluble in HCl or  $HNO_3$ .

$SnCl_2$  or  $H_2SO_3$  gives in the presence of free HCl a red, or in warm solutions a gray pp. of Se.

Cu in a hot HCl solution becomes covered with a black film: the liquid on standing with the Cu for some time becomes colored red with Se.

*b. Selenates:*

HCl produces no change in the cold; but on boiling Cl is given off and the selenate reduced to selenite, to which the above tests under *a.* may be applied.

$BaCl_2$ : a white pp. ( $BaSeO_4$ ); insoluble in cold HCl; dissolved by boiling with HCl with evolution of Cl and reduction to  $BaSeO_3$ .

*c. Selenium in any form of combination* is detected by the following reactions:

Heated on charcoal in the inner blowpipe flame a smell of rotten horse-radish is perceived.

Fused on charcoal with  $Na_2CO_3$  in the inner blowpipe flame a fused mass is obtained, which when moistened on Ag yields a black stain, and on addition of HCl evolves  $H_2Se$ .

**545. TELLURIUM (Te).**—Use a metallic telluride, an alkaline tellurite and tellurate.

Te occurs united with metals (*e. g.*, Au, Ag, Bi, Cu, Pb).

A telluride heated in an open glass tube gives white fumes and a sublimate which differs from that given by Sb, by being fusible before the blowpipe.

*a. Tellurites:*

$H_2O$ : on dilution with water tellurous acid is precipitated from the acid solution.

$H_2S$ : in acid solutions a brown pp. ( $TeS_2$ ); easily soluble in  $Am_2S$ .

$H_2SO_3$ ,  $SnCl_2$ , or Zn precipitates black Te.

*b. Tellurates.*

HCl produces no change in the cold, but on boiling Cl is evolved, and the tellurate is reduced tellurite; the solution then gives the above reactions under (a).

*c. Tellurium in any form of combination*, if fused with  $\text{Na}_2\text{CO}_3$  on charcoal in the inner blowpipe flame, gives sodium telluride; a solution of which gives a black stain on Ag, and on being acidified deposits black Te and evolves  $\text{TeH}_2$ .

## GROUP III.—IRON AND ZINC GROUPS.

In this group are included U, In, Ti, Be, Tl [V], (Zr, Ce, Ta, Nb, La, Di, Y, E, Th).

Tl is often partially precipitated as chloride in Group I, and its reactions are given under that group. V is not precipitated by  $\text{Am}_2\text{S}$  unless acid is added in excess after  $\text{Am}_2\text{S}$ .

**546. URANIUM (U).—Use  $(\text{UO}_2)''(\text{NO}_3)_2$ .**

U occurs in nature principally as *pitchblende* (oxide), also as *uranite* (a hydrated uranium-calcium phosphate) and *chalcolite* (hydrated uranium-copper phosphate).

$\text{AmHO}$  ( $\text{KHO}$ ,  $\text{NaHO}$ ): yellow pp., insoluble in excess.

$\text{Am}_2\text{S}$  gives in neutral solutions a *dingy-yellow* or *brown* pp. of uranium oxysulphide, which is soluble in  $\text{Am}_2\text{CO}_3$  (diff. from  $\text{ZnS}$ ,  $\text{MnS}$ ,  $\text{FeS}$ , etc.). The pp. settles slowly unless  $\text{AmCl}$  is added; it is soluble in acids, even in  $\text{H}\bar{\text{A}}$ : on heating with  $\text{Am}_2\text{S}$  in excess it is changed into uranous oxide and sulphur.

$\text{H}_2\text{S}$  produces no pp. in acid solutions.

$\text{Am}_2\text{CO}_3$  ( $\text{KHCO}_3$ ,  $\text{NaHCO}_3$ ): yellow pp., easily soluble in excess; from this solution the uranium is reprecipitated by addition of  $\text{NaHO}$  or  $\text{KHO}$ , or by boiling. (Diff. from Fe.)

$\text{K}_2\text{FeCy}_6$  gives in acid solutions a reddish-brown pp. The pp. is distinguished from the similar one produced in a copper solution by dissolving in excess of  $\text{AmHO}$  to a *yellow* fluid.

$\text{BaCO}_3$ : complete precipitation even in the cold.

Zn changes the *yellow* color of acid solutions to *green*.

*Borax* and *Microcosmic beads*:

Outer flame, *yellow*; inner flame, *green*.

**547. INDIUM (In).**

*Alkali-hydrates* precipitate a hydrate resembling  $\text{Al}_2\text{H}_6$ ; insoluble in excess. Indium solutions are also precipitated by alkaline carbonates, by  $\text{Na}_2\text{HPO}_4$ , by boiling their neutral solutions with  $\text{Na}\bar{\text{A}}$ , by  $\text{BaCO}_3$ , and by alkaline oxalates.

$\text{H}_2\text{S}$  gives no pp. in strong acid solutions, in dilute and feebly acid solutions a little sulphide separates, in a solution containing no free acid but  $\text{H}\bar{\text{A}}$  the metal is entirely precipitated as yellow sulphide: the pp. is insoluble in cold, soluble in boiling,  $\text{Am}_2\text{S}$ ; from the boiling solution white sulphide separates on cooling.

$\text{Am}_2\text{S}$ , added after  $\text{H}_2\text{T}$  and excess of  $\text{AmHO}$ , gives a white pp., becoming yellow on treatment with  $\text{H}\bar{\text{A}}$ .

*Flame coloration*: bluish-violet. The spectrum shows two very characteristic blue lines, which are brilliant, but very rapidly disappear when the chloride is employed: see p. 860.

**548. BERYLLIUM (Be).**—Occurs as silicate in phenakite, and as silicate with Al-silicate in beryl and emerald.

$\text{Am}_2\text{S}$  ( $\text{AmHO}$ ,  $\text{KHO}$ ,  $\text{NaHO}$ ): flocculent hydrate is precipitated resembling  $\text{Al}_2\text{H}_6$  in its appearance, and in being soluble in  $\text{KHO}$ . Differs from  $\text{Al}_2\text{H}_6$  in being precipitated from its solution in  $\text{KHO}$  by dilution and long boiling, and by being soluble when freshly precipitated if it is long boiled with  $\text{AmCl}$  solution.

*Alkaline carbonates* precipitate a carbonate soluble in excess, especially in  $\text{Am}_2\text{CO}_3$ ; from these solutions the carbonate is reprecipitated, on diluting and boiling for some time, with especial ease from the  $\text{Am}_2\text{CO}_3$  solution (diff. from Al).

$\text{BaCO}_3$  precipitates Be solutions completely:  $\text{H}_2\text{C}_2\text{O}_4$  and alkaline oxalates produce no pp. Moistened with  $\text{Co}(\text{NO}_3)_2$  solution and heated on charcoal in the outer blowpipe flame, a gray mass is obtained (diff. from Al).

**549. TITANIUM (Ti.).**—Occurs as  $\text{TiO}_2$  (with traces of Fe, Mn, Cr) in rutile, anatase, and brookite: also as  $\text{TiO}_2$  in combination with Fe in titaniferous iron ore.

Ignited  $\text{TiO}_2$  is insoluble in water and most acids: it is easily soluble in  $\text{HF}$ , less readily in boiling strong  $\text{H}_2\text{SO}_4$ : it is also rendered soluble in cold water by fusion with  $\text{KHSO}_4$ .  $\text{TiO}_2$  differs from  $\text{SiO}_2$  in not being volatilized when heated in a platinum dish with  $\text{HF}$  and strong  $\text{H}_2\text{SO}_4$ .

By dilution and long boiling, hydrated  $\text{TiO}_2$  is precipitated as a white flocculent pp. from solutions in  $\text{H}_2\text{SO}_4$  (or  $\text{HCl}$ ), and from the aqueous solution of the fusion with  $\text{KHSO}_4$ :



## 352 APPENDIX I.—VANADIUM, LITHIUM. [550, 551.]

the pp. is metatitanic acid. It cannot be filtered off unless an acid or  $\text{AmCl}$  is added.

$\text{AmHO}$  ( $\text{KHO}$ ,  $\text{NaHO}$ ,  $\text{Am}_2\text{S}$ , or  $\text{BaCO}_3$ ), white flocculent pp., insoluble in excess: if precipitated and washed in the cold the pp. dissolves in  $\text{HCl}$  and in dilute  $\text{H}_2\text{SO}_4$ .

$\text{Zn}$  or  $\text{Sn}$  gives in acid solutions a *blue*, or if dilute a *rose-colored*, liquid.

$\text{Na}_2\text{S}_2\text{O}_3$ , on boiling, precipitates  $\text{Ti}$  solution entirely (diff. from  $\text{Fe}$ ).

### Microcosmic bead:

In the outer flame: *yellow* whilst hot; *colorless*, cold.

In the inner flame: *yellow*, hot; *violet*, cold. The production of these colors is much aided by adding a fragment of  $\text{Sn}$ . Addition of a small quantity of  $\text{FeSO}_4$  gives in the inner flame a *blood-red* bead.

### 550. VANADIUM (V.)

Occurs in vanadinite [ $3\text{Pb}_3(\text{PO}_4)_2 \cdot \text{PbCl}_2$ ], and in certain  $\text{Fe}$  and  $\text{Cu}$  ores.

Vanadium is known in several stages of oxidation; it will usually occur in analysis as vanadic acid or a vanadate, which in acid solution is of a *yellow* or *reddish* color.

$\text{H}_2\text{S}$  ( $\text{H}_2\text{SO}_3$  or  $\text{H}_2\text{C}_2\text{O}_4$ ) reduces acid solutions of a vanadate, and causes the color of the solution to change to *blue*: with  $\text{H}_2\text{S}$  a deposit of sulphur also forms.

$\text{Am}_2\text{S}$ , if added in excess, gives a brown liquid, in which an excess of acid causes a *brown* pp. of  $\text{V}_2\text{S}_5$ ; excess of  $\text{Am}_2\text{S}$  dissolves this to a reddish-brown liquid.

$\text{Zn}$  gives in a very dilute solution acidified with  $\text{H}_2\text{SO}_4$ , and gently warmed, a *blue* color, which changes into *green* and then *lavender-blue*.

Solid  $\text{AmCl}$ , added until the solution is saturated, precipitates  $\text{Am}_2\text{VO}_4$ , which is insoluble in saturated  $\text{AmCl}$  solution. Very characteristic reaction.

$\text{H}_2\text{O}_2$  when shaken up with an acid solution imparts a red color, which is not removed by shaking with ether. This is a very delicate test.

### Borax bead:

In outer flame; *colorless* or *yellow* if much  $\text{V}$  is present.

In inner flame; *green* hot and cold; if much  $\text{V}$  is present, *brown* hot, and *green* cold.

## GROUP V.—POTASSIUM GROUP.

In this group are included  $\text{Li}$ ,  $\text{Cs}$ ,  $\text{Rb}$ .

### 551. LITHIUM ( $\text{Li}$ ).

$\text{Li}$  occurs frequently in mineral waters and in the ashes of plants, also in small quantities in the minerals

lepidolite, triphane, and petallite. It is allied to the Barium Group by the difficult solubility of its carbonate and phosphate; and it differs from K and  $\text{NH}_4$  by not being precipitated by  $\text{PtCl}_4$  or by  $\text{H}_2\text{T}$ : from Na it is readily distinguished by the flame coloration and spectroscopy.  $\text{LiCl}$  is separated from  $\text{KCl}$  and  $\text{NaCl}$  by its insolubility in a mixture of absolute alcohol and ether.

$\text{Na}_2\text{HPO}_4$  added to the not too dilute solution, made decidedly alkaline with  $\text{NaHO}$ , gives, on boiling, a white crystalline pp. ( $\text{Li}_3\text{PO}_4$ ) which quickly settles. Traces of Li may be precipitated by adding  $\text{Na}_2\text{HPO}_4$ , then  $\text{NaHO}$  until the liquid remains alkaline, evaporating to dryness and washing the residue with dilute  $\text{AmHO}$ . This pp. differs from the phosphates of Ba, Sr, Ca, and Mg by fusing in the blowpipe flame and being absorbed by the charcoal support, also by its diluted solution in  $\text{HCl}$  giving no pp. on addition of  $\text{AmHO}$  in excess in the cold, but a white crystalline pp. on boiling.

*Flame coloration*: this is *carmine-red*; the spectrum (see p. 360) consists of two lines, an intense carmine-red line ( $\alpha$ ) and a feeble orange-yellow line ( $\beta$ ). This coloration is concealed by that of Na, which, however, does not interfere with the lithium spectrum: the Na coloration is also removed if the flame is viewed through the indigo-prism, whereas that of Li can pass through the thinner parts of the prism; it differs from the K coloration, however, in being unable to penetrate the thick layers of indigo solution, or in being much weakened by its passage. As silicate, Li only gives the flame coloration after fusion with  $\text{CaSO}_4$ ; the phosphate requires to be first moistened with  $\text{HCl}$ .

## 552. RUBIDIUM (Rb), and CÆSIUM (Cs).

These metals occur in small quantities in some mineral waters and in a few minerals. Their compounds resemble those of K in being precipitated by  $\text{PtCl}_4$  and by  $\text{H}_2\text{T}$ , and in giving a similar flame coloration. Characteristic differences are, the much greater insolubility in water of the platino-chlorides, which enables the  $\text{K}_2\text{PtCl}_6$  to be dissolved away from  $\text{Rb}_2\text{PtCl}_6$  and  $\text{Cs}_2\text{PtCl}_6$  by boiling the pp. with successive small quantities of water: the alums also show a similar difference in solubility in cold water.

Cs and Rb are readily distinguished from one another and from other elements by their spectra (see p. 360). For Cs the blue lines ( $\alpha$ ,  $\beta$ ) are especially distinct and characteristic: in the Rb spectrum the indigo-blue lines ( $\alpha$ ,  $\beta$ ) are very distinct, but the red lines ( $\gamma$ ,  $\delta$ ) are most characteristic. For the spectroscopic test the chlorides are the most suitable.

## 553.—GENERAL GROUP TABLE FOR

Group I. Reagent HCl.	Group II. Reagent H <sub>2</sub> S.
<p><i>The group pp. may contain in addition to AgCl, Hg<sub>2</sub>Cl<sub>2</sub>, PbCl<sub>2</sub>:</i></p> <p><i>TlCl:</i> which will usually be readily detected by a spectroscopic examination of the group pp. It may be removed from the group pp. by boiling with a little water and pp<sup>d</sup> from the cold filtrate by KI (Pb is also thus pp<sup>d</sup>): the pp. is yellow, and gives the thallium spectrum.</p> <p><i>H<sub>2</sub>WO<sub>4</sub>:</i> a white pp., becoming yellow on boiling. Its presence is confirmed by dropping a piece of Zn into a portion of the acid liquid and pp. when a <i>deep blue</i> color will be obtained.</p> <p>Also by fusion of the pp. in a microcosmic bead, which will be <i>colorless</i> or <i>yellow</i> in the outer flame, <i>blue</i> (or with FeSO<sub>4</sub>, <i>blood-red</i>) in the inner flame.</p>	<p><i>The group pp. may contain as sulphides, in addition to Hg, Pb, Bi, Cu, Cd:</i></p> <p><i>Pd (black-brown), Tl (black; pp<sup>d</sup> with As<sub>2</sub>S<sub>3</sub>) [Os, Rh, Ru.*]</i> } Insoluble in Am<sub>2</sub>S.</p> <p>—</p> <p><i>And in addition to SnS, SnS<sub>2</sub>, Sb<sub>2</sub>S<sub>3</sub>, As<sub>2</sub>S<sub>3</sub>, Au<sub>2</sub>S<sub>3</sub>, PtS<sub>2</sub>:</i></p> <p><i>[Ir-sulphide*] Mo- " (brown) Se- " (red-yellow) Te- " (black)</i> } Soluble in Am<sub>2</sub>S.</p> <p>—</p> <p>The <i>yellow</i> color of a solution containing V changes to <i>blue</i>.</p> <p>=====</p> <p><i>Tl</i> is readily found by the spectroscope in the group pp.</p> <p><i>Pd</i> will remain in solution in excess of AmHO with Cd and Cu in Table II A; from this solution it is pp<sup>d</sup> by adding HCl: its presence may be further confirmed by the HgCy<sub>2</sub> or KI tests.</p> <p>—</p> <p>The examination of the sulphides in Group II B, when the above elements may be present is best commenced by fusion with Na<sub>2</sub>CO<sub>3</sub> and NaNO<sub>3</sub>. From the fused mass, water dissolves sodium-arsenate, -molybdate, -selenate, and -tellurate, leaving SnO<sub>2</sub>, sodium-antimonate, Au, Pt, and Ir undissolved; the elements may then be detected by special tests. For the separation and detection of the platinum metals a larger work must be consulted.</p>

\* Only completely pp<sup>d</sup>, if the liquid has been

## PRECIPITATION OF RARER ELEMENTS.

Group III. Reagents AmCl, AmHO, and Am <sub>2</sub> S.	Precipitated from the filtrate from Group III by excess of HCl.	Group V.
<p><i>The group pp. may contain as sulphides, in addition to Fe, Zn, Mn, Ni, Co:</i></p> <p><i>U (black-brown).</i>  <i>In ( " " ).</i>  <i>Ti ( " " ).</i></p> <hr/> <p><i>And as hydrates in addition to Al, Cr (and phosphates, etc.):</i></p> <p><i>Ti (white, flocculent).</i>  <i>Be (colorless, gelatinous).</i>  <i>[Zr] ( " " ).</i>  <i>[Ta, Nb, Cs, La, Di, Y, E, Th.]</i></p> <hr/> <p><i>Ti and In will usually be readily detected by a spectroscopic examination of the pp.</i></p> <p><i>Ti may be first separated and detected by dissolving some of the group pp. in boiling dilute HCl, and reducing any Fe present in the solution by H<sub>2</sub>SO<sub>4</sub>; then precipitating Ti by addition of KI, and testing the pp. by the spectroscope.</i></p> <hr/> <p><i>A more complete scheme for the analysis of this group pp. will be found in par. 554 (p. 356.)</i></p>	<p><i>As sulphides:</i></p> <p><i>V (black).</i>  <i>W (trace, brown).</i>  <i>Ni (trace, black).</i></p> <hr/> <p><i>This pp. is dried and then fused with a mixture of Na<sub>2</sub>CO<sub>3</sub> and KNO<sub>3</sub>: on boiling the mass with water, NiO remains undissolved; it is filtered off, and its presence confirmed by fusion in a borax bead.</i></p> <p><i>The aqueous solution may contain alkaline vanadate and tungstate. Vis separated by saturating the liquid with AmCl, and its presence confirmed in the pp. by the borax bead, and by dissolving the pp. in HCl and employing the reactions with H<sub>2</sub>O<sub>2</sub> and with Zn.</i></p> <p><i>The filtrate (after concentration, if necessary) is acidified with HCl, and the presence of W confirmed by the Zn reaction and the microcosmic bead.</i></p>	<p><i>The filtrate after the separation of all the group precipitates, may contain besides Mg, K, and Na:</i></p> <p><i>Li, Cs, Eb.</i></p> <hr/> <p><i>These metals are readily detected by their very characteristic spectra (p. 360).</i></p> <p><i>LiCl may also be detected by its behavior with Na<sub>2</sub>HPO<sub>4</sub>.</i></p> <p><i>Before proceeding to the spectroscopic examination, it is well to convert them into chlorides if they are not already in that form.</i></p> <p><i>If the chlorides are evaporated to dryness, and extracted with absolute ether and alcohol, to which a few drops of HCl have been added, LiCl, RbCl, CsCl dissolve, whilst the greater part of the KCl and NaCl remain undissolved.</i></p> <p><i>The solution is evaporated to dryness, dissolved in a little HCl and Rb and Cs are ppd. by PtCl<sub>4</sub>; this pp. is boiled with small quantities of water until it no longer gives the K spectrum, the spectra of Rb and Cs will then be seen, if present. Li is found in the filtrate after adding PtCl<sub>4</sub>.</i></p>

warmed, and H<sub>2</sub>S passed for a long time.



## APPENDIX II.

### USE OF THE SPECTROSCOPE FOR ANALYSIS.

555. When the light from a bright gas or candle-flame is allowed to pass through a narrow slit, and is then looked at through an appropriately cut glass prism, a series of colors is seen following one another without break or interruption; such a band of colors has received the name of a "continuous spectrum." The direct-vision spectroscope\* is a little instrument furnished at one end with the requisite slit, whose width may be varied by means of a projecting screw-collar: at the other end will be found a draw-tube containing lenses for focussing, and the body of the instrument incloses the glass prisms. On looking at a bright gas or candle-flame through this spectroscope a "continuous spectrum" will be seen, whose margins may be sharply focussed by means of the draw-tube. A similar spectrum is seen when any brightly glowing *non-volatile* substance is viewed through the spectroscope.

Many substances, however, when strongly heated volatilize; and their *glowing vapors* emit light whose color is in many cases peculiar and characteristic; mention has been already made of this fact in par 32, and its importance as a means of detecting certain elements was there stated.

When such a vapor is viewed through the spectroscope, a "discontinuous (or 'line') spectrum" is seen, which consists of one or more colored "lines" or "bands," whose position and color are characteristic of that particular vapor. Thus Na vapor yields one bright yellow line, Li one intense carmine line, whilst K gives two lines, one red and the other violet. These lines are readily seen by holding in the Bunsen flame a loop of platinum-wire, which has been previously dipped into strong NaCl, LiCl, or KCl solution, and viewing the colored flame through the spectroscope. But not only does this method of analysis serve to detect the elements when occurring separately, it also detects each of them when two or more are present together, since the spectrum of each is then seen without any interference from the others: thus, although the *flame coloration* of K or of Li is entirely masked by that of Na when the chlorides of both metals are held in the flame together, the two *spectra* are distinctly seen side by side when the flame is looked at through the spectroscope; and both Li and K are thus readily recognized when present with Na.


---

\* This is the cheapest and most handy form of spectroscope, and may be purchased of the makers, Messrs. Browning, of the Strand, London.

The delicacy of this method of detection is extreme; thus the ~~presence~~ of a gramme of Na can be distinguished by the appearance of its yellow spectral line.

The accompanying chart (p. 360) gives the relative position of the most important lines in thirteen different spectra. It is meant to serve merely as a guide to the beginner: when the student has accustomed himself to the appearance of the spectra, reference to the chart will become unnecessary.

It is important to remember that the coloration of the Bunsen flame can only be produced by substances which are volatile at the temperature of the flame, and that the coloration due to any element is more intense the more volatile is the compound employed. Thus KCl and KNO<sub>3</sub> are readily volatilized in the flame, and color it strongly; K<sub>2</sub>SO<sub>4</sub> is somewhat less volatile, and the coloration it yields is correspondingly feeble; K<sub>2</sub>CO<sub>3</sub> is still less volatile, and imparts little color to the flame; whilst K<sub>2</sub>SiO<sub>3</sub> is non-volatile, and yields no flame coloration whatever. Hence in testing for an element by the spectroscopy, care must be taken to insure its conversion, if present, into a form of combination which is volatile in the Bunsen flame. As a rule, chlorides and nitrates of the metals are the most volatile salts; sulphates are less volatile, carbonates still less so, and silicates are usually non-volatile. It is in many cases sufficient to moisten the substance upon the platinum-wire loop with strong H<sub>2</sub>SO<sub>4</sub> or HCl before introducing it into the flame; the sulphates of Ba, Sr, and Ca should be first reduced to sulphides by heating them in the inner blowpipe flame, the sulphides are then readily converted into chlorides by moistening them with HCl. K<sub>2</sub>SiO<sub>3</sub> in a natural silicate yields no K flame coloration; by fusion of the powdered mineral with gypsum powder, however, the K<sub>2</sub>SiO<sub>3</sub> is converted into K<sub>2</sub>SO<sub>4</sub>, and the sulphate yields the K coloration readily:



Owing to the invariable presence of small quantities of Na in atmospheric dust, and to the extreme delicacy of the spectroscopic reaction for Na, the Bunsen flame always shows the Na line when looked at through the spectroscopy, and the brightness of the line may be increased by striking the table, or in any other way raising dust, in the neighborhood of the flame. This is in many respects advantageous; since, by rendering the Na line distinct, by regulating the width of the slit and the position of the focussing lens, the spectroscopy may be placed in adjustment before introducing into the flame the substance whose spectrum is to be observed. The Na line further serves as a fixed point from which the position of the lines of other elements may be judged; and in order to assist the student in so using it, the position of the Na line is indicated in all the other spectra on the accompanying chart by means of a dotted line.

**556.** When working with the spectroscopy the Bunsen flame should be so placed as to have a dark background, in order that light *from the flame only* may enter the slit: if not working in a darkened room, a piece of black velvet or cloth may be hung behind the flame; this is, however, not indispensable; it will usually

be sufficient to avoid the entrance of direct or brightly reflected light into the instrument. It is best to view first either the sky or a bright flame—such as that obtained by closing the air-holes at the foot of the Bunsen burner, and to turn the spectroscope into such a position that the continuous spectrum becomes horizontal with its red end to the left hand; the instrument is next directed on a portion of the non-luminous Bunsen flame about two-thirds of the height from the flame's base, and held either by a clamp or very steadily by the hand, in this position. The Na line is then focussed as directed above, and, whilst carefully watching the flame through the spectroscope, the substance is introduced into the outer part of the front of the flame at about one-third the flame's height from its base: the observation of the flame is continued for a short time after moistening the substance with acid as directed above. The careful observation of the flame during the introduction of the substance into the flame is necessary, since some elements (*e. g.*, Tl), when present in small quantity, give a spectrum which appears only as a momentary flash: other substances appear only as the substance attains the full temperature of the flame, and their spectra are therefore seen only after the substance has been heated in the flame for some time.

The regulation of the width of the slit must depend upon circumstances. A wide slit admits more light and increases the brightness of the lines; a narrow slit causes the lines to be more widely separated, and thus prevents neighboring lines from blending, and generally enables their position to be more accurately noted. It is well to use a narrow slit, unless working with a faint flame coloration, or unless lines of feeble intensity have to be looked for.

The spectroscope is more especially useful in qualitative analysis:

1st. In detecting readily K and Li, and other substances whose flame colorations are masked by that of Na, when occurring with Na.

2d. It also rapidly distinguishes Sr from Ca; since as shown in the chart the Sr spectrum contains a characteristic blue line  $\delta$ , as well as certain red lines situated to the left hand of all those in the Ca spectrum; the green line  $\beta$  of Ca is also very distinctive of that element. Ba is specially characterized by the four green lines,  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ .


3d. The spectroscope is of great value for detecting the rarer metals, Li, In, Tl, Cs, Rb, which usually occur in comparatively small quantity in nature: the last four of these metals were first searched after and discovered on account of their very marked spectra having been seen by the spectroscope.

**557. Spectrum Chart.**—In using the chart below, the color of a line in any spectrum may be found by referring to the head of the chart, and seeing in which of the bracketted spaces it falls: above each bracket is placed the initial letter of the name of the color: the colors run from left to right in the following order: Red, Orange, Yellow, Green, Blue, Indigo, Violet.

The chart is a reduction from Bunsen's map: the relative position of the lines is given correctly: those in each spectrum which



are most striking or characteristic are made thicker than the rest, and are denoted by Greek letters in the order of their relative importance. Many of the thin lines will not be seen by means of a small pocket spectroscope.



\* This spectrum is seen when boric acid or a borate is fused with a little  $\text{Na}_2\text{CO}_3$  on a loop of platinum wire, and the flame coloration obtained from this bead is examined.

## APPENDIX III.

### 558. LIST OF CHEMICAL ELEMENTS, WITH THEIR SYMBOLS AND ATOMIC WEIGHTS.

*Note.*—The words in brackets are the Latin names of the elements from which the symbols have been derived.

Name.	Symbol.	Atomic Weight.	Name.	Symbol.	Atomic Weight.
Aluminium.....	Al.	27.5	Molybdenum.....	Mo.	96
Antimony (stibium).....	Sb.	122	Nick-el.....	Ni.	59
Arsenicum.....	As.	75	Niobium.....	Nb.	94
Barium.....	Ba.	137	Nitrogen.....	N.	14
Bismuth.....	Bi.	210	Osmium.....	Os.	199
Boron.....	B.	11	Oxygen.....	O.	16
Bromine.....	Br.	80	Palladium.....	Pd.	106
Cadmium.....	Cd.	112	Phosphorus.....	P.	31
Cæsium.....	Cs.	133	Platinum.....	Pt.	197
Calcium.....	Ca.	40	Potassium (kalium).....	K.	39
Carbon.....	C.	12	Rhodium.....	Rh.	104
Cerium.....	Ce.	92	Rubidium.....	Rb.	85
Chlorine.....	Cl.	35.5	Ruthenium.....	Ru.	104
Chromium.....	Cr.	52.5	Selenium.....	Se.	79.5
Cobalt.....	Co.	59	Silicium.....	Si.	28
Copper (cuprum).....	Cu.	63.5	Silver (argentum).....	Ag.	108
Didymium.....	D.	96	Sodium (natrium).....	Na.	23
Erbium.....	E.	112	Strontium.....	Sr.	87.5
Fluorine.....	F.	19	Sulphur.....	S.	32
Gluccium.....	G.	9.5	Tantalum.....	Ta.	182
Gold (aurum).....	Au.	197	Tellurium.....	Te.	129
Hydrogen.....	H.	1	Thallium.....	Tl.	204
Iodine.....	I.	127	Thorium.....	Th.	238
Iridium.....	Ir.	197	Tin (stannum).....	Sn.	118
Iron (ferrum).....	Fe.	56	Titanium.....	Ti.	50
Lanthanum.....	La.	92	Tungsten (Wolframium).....	W.	184
Lead (plumbum).....	Pb.	207	Uranium.....	U.	120
Lithium.....	Li.	7	Vanadium.....	V.	51
Magnesium.....	Mg.	24	Yttrium.....	Y.	62
Manganese.....	Mn.	55	Zinc.....	Zn.	65
Mercury (hydrargyrum).....	Hg.	200	Zirconium.....	Zr.	89

### 559. THERMOMETRIC SCALES.

There are two different thermometric scales in use in this country, the Centigrade and Fahrenheit; of these two the former is rapidly becoming universal for scientific purposes. The two scales are mutually convertible by the following formulæ, in which F.° rep-

resents a temperature on the Fahrenheit scale, C.° a temperature on the Centigrade scale:

$$\begin{aligned} \frac{5}{9} (F.^{\circ} - 32) &= C.^{\circ} \\ \frac{9}{5} C.^{\circ} + 32 &= F.^{\circ} \end{aligned}$$

The temperatures occasionally referred to in this treatise are given on the Centigrade scale.

### WEIGHTS AND MEASURES.

**560.** The corresponding values of the French and English weights and measures are here given; the use of the French or decimal system is strongly recommended by its extreme simplicity, since the smaller denominations are obtained by taking a tenth, hundredth, thousandth, etc., of the unit chosen, and are designated by the Latin prefixes deci-, centi-, milli-, etc., whilst the higher denominations are 10 times, 100 times, 1000, etc., times the unit, and are named by the Greek prefixes deca-, hecto-, kilo-, etc.; examples of this will be found in the tables given below.

**561.** The starting-point of the French system is the "metre" (= 39.37 inches); this is the "unit of length." The "unit of measure" is the "litre," which is one cubic decimetre: the "unit of weight" is the gramme,\* which is the weight of 1 cubic centimetre of distilled water at 4° C.

The chief conveniences arising from the use of this system are:

1st. That all the different denominations can be written as one, since they are either multiples by ten or are decimal fractions of the unit. Thus 5 decagrammes, 3 grammes, 4 decigrammes, 8 milligrammes, would be written 53.408 grammes.

2d. That since 1 cubic centimetre of water at 4° C. weighs 1 gramme, we may obtain the *weight* of water to be used from the *measure* by simply converting the measure into cubic centimetres; the number thus obtained will represent at once the corresponding weight of water in grammes. Of course this conversion is strictly accurate only when the water is measured at 4° C., but for ordinary purposes the error introduced when the water is at the temperature of the air is too small to be of any importance in the preparation of solutions.

The weights and measures most frequently used for chemical purposes are the gramme, the millimetre,† the litre, and the cubic centimetre,‡ which is  $\frac{1}{1000}$ th of a litre.

MEASURES OF LENGTH.				ENGLISH.			
	Metre.	Inches.		Mile.	Furlong.	Yards.	Feet. Inches.
Millimetre	— 0.001 —	— .003937 —					.63937
Centimetre	— 0.01 —	— .39371 —					.39371
Decimetre	— 0.1 —	— 3.93708 —					3.9371
Metre	— 1.0 —	— 39.37079 —				3	3.371
Decametre	— 10.0 —	— 393.70790 —				10	2 9.7
Hectometre	— 100.0 —	— 3937.07900 —				109	1 1
Kilometre	— 1000.0 —	— 39370.79000 —			4	213	4 10.2
Myriometre	— 10000.0 —	— 393707.90000 —		6	1	156	0 6
		1 inch = .0254 metre.					
		1 foot = .3048 "					

\* Frequently written "gram" in English, for the sake of brevity.

† Usually written (m.m.).

‡ Usually abbreviated (c.c.).

## MEASURES OF CAPACITY.

1 Litre = 1 cubic decimetre.

	Litre.	Cubic inches.	Pints.
{ Millilitre, or Cubic centimetre (c.c.) }	.001	.06108	0.00176
Centilitre	.01	.61027	0.10761
Decilitre	.1	6.1027	0.17608
Litre	1.0	61.027	1.76077
Decalitre	10.0	610.27	17.60773
Hectolitre	100.0	6102.7	176.07734
Kilolitre	1000.0	61027.0	1760.77341
Myriolitre	10000.0	610270.0	17607.73414

1 cubic inch = .01639 litre.

1 cubic foot = 28.31531 litres.

1 gallon = 4.54336 "

## MEASURES OF WEIGHT.

1 gramme = the weight of 1 cubic centimetre (c.c.) of water at 4° C.

	Gramme.	Grains.	Avoirdupois.		
Milligramme	.001	0.01543			
Centigramme	.01	0.15432			
Decigramme	.1	1.54323			
Gramme	1.0	15.43235			
Decagramme	10.0	154.32349	lbs.	oz.	drms.
Hectogramme	100.0	1543.23488	— 0	0	5.65
Kilogramme	1000.0	15432.34880	— 0	3	8.5
Myriogramme	10000.0	154323.48800	— 2	3	5
			— 22	1	2

1 grain = 0.0649 gramme.

1 oz. (Troy) = 31.1035 grammes.

1 lb. (Avoirdupois) = 453.593 "

## ENGLISH WEIGHTS AND MEASURES.

## APOTHECARIES' WEIGHT.


lb.	oz.	drms.	scruples.	grains.
1	12	96	288	5760
	1	8	24	480
		1	3	60
			1	20

## AVOIRDUPOIS WEIGHT.

lb.	oz.	drms.	grains.
1	16	256	7000
	1	16	437.5
		1	27.343

## IMPERIAL MEASURE.

	gallon.	pinta.	fluid oz.	fluid drms.
	1	8	160	1280
		1	20	160
			1	8
1 gallon				70,000 grains of water at 16.7° C.
1 fluid ounce	—	$\frac{1}{8}$ pint	—	437.5 "
1 gallon	—		277.280 cubic inches.	
1 fluid ounce	—		1.733 "	



# INDEX.

---

	PAGE
Acid, test for an, . . . . .	78
"    radicles, detection of in simple substances, . . . . .	207
"    "    detection of, in complex substances, . . . . .	265
"    "    reactions of, . . . . .	146
Acids, removal from clothes, . . . . .	84
Acetates, reactions of, . . . . .	183
Acetic acid, . . . . .	183
Alkaline reaction, . . . . .	74
Aluminium, reactions of, . . . . .	104
Ammonia gas, preparation of, . . . . .	31
"    "    properties of, . . . . .	82
"    "    tests for, . . . . .	38
Ammonium, reactions of, . . . . .	92
Analysis, course of, . . . . .	78
"    introduction to, . . . . .	80
"    of a simple liquid, . . . . .	194
"        "    solid, . . . . .	188
"        a complex liquid, . . . . .	237
"        "    solid, . . . . .	238
"        an alkaline liquid, . . . . .	214
"        cyanogen compounds, . . . . .	291
"        insoluble substances, . . . . .	218, 286
"        metallic substances, . . . . .	217, 281
"        silicates, . . . . .	290
"        substances for, . . . . .	341
Analytical classification, . . . . .	81
"        "    table of, . . . . .	186
"        groups, . . . . .	82
"        reactions, . . . . .	89, 146
Antimony, reactions of, . . . . .	135
Apparatus, list of students', . . . . .	808
"    list of general, . . . . .	806
"    cleaning of, . . . . .	49
Arsenic group, . . . . .	127
"    "    separation and detection of, . . . . .	139
"    "    table of differences, . . . . .	139
"    "    table for analysis, . . . . .	199, 256
"    "    reactions of, . . . . .	128
Atomic weights, list of, . . . . .	861

	PAGE
Barium group, . . . . .	100
" " separation and detection of, . . . . .	102
" " table of differences, . . . . .	102
" " table for analysis, . . . . .	206, 261
Barium, reactions of, . . . . .	100
Bending glass tube, . . . . .	44
Beryllium, detection in analysis, . . . . .	355, 356
" reactions of, . . . . .	351
Bismuth, reactions of, . . . . .	122
Blowpipe, use of, . . . . .	42, 66
Borates, reactions of, . . . . .	169
Borax bead, . . . . .	67
Boric acid, reactions of, . . . . .	170
Boring corks, . . . . .	45
Bromides, reactions of, . . . . .	159
Bunsen burner, . . . . .	40
 Cadmium, reactions of, . . . . .	 125
Cæsium, detection in analysis, . . . . .	355
" reactions of, . . . . .	353
Calcium, reactions of, . . . . .	101
" sulphate, preparation of, . . . . .	332
Carbon dioxide, preparation of, . . . . .	25
" " from combustion, . . . . .	30
" " from respiration, . . . . .	29
" " properties of, . . . . .	26-29
" " tests for, . . . . .	29
" " in the air, . . . . .	31
Carbon monoxide, preparation of, . . . . .	33
" " properties of, . . . . .	35
" " tests for, . . . . .	35
Carbonates, reactions of, . . . . .	143
Carbonic acid, reactions of, . . . . .	149
Centigrade scale, . . . . .	361
Charcoal, wood, . . . . .	341
" combustion in oxygen, . . . . .	20
" ignition on, . . . . .	70
Chemicals, list for Section I, . . . . .	389
" " " III, . . . . .	340
Chlorates, reactions of, . . . . .	156
Chloric acid, reactions of, . . . . .	157
Chloride group, separation and detection of, . . . . .	164
Chlorides, reactions of, . . . . .	157
Chlorine, preparation and properties of, . . . . .	36
" tests for, . . . . .	36
" water, preparation of, . . . . .	333
Chromium, reactions of, . . . . .	106
Chromates, reactions of, . . . . .	170
Classification of metals, . . . . .	82
" " table of, . . . . .	186

	PAGE
Cleaning apparatus, . . . . .	49
" platinum, . . . . .	50
Cobalt glass, . . . . .	306
" reactions of, . . . . .	114
Combustion in oxygen, of charcoal, . . . . .	20
" " phosphorus, . . . . .	21
" " sulphur, . . . . .	21
Contractions, list of, . . . . .	82, 87
Copper group, separation and detection, . . . . .	126
" " table of differences, . . . . .	125
" " for analysis, . . . . .	199, 255
" reactions of, . . . . .	123
Corks, . . . . .	341
" boring of, . . . . .	45
Cutting glass tube, . . . . .	43
Cyanides, reactions of, . . . . .	176
Decantation, process of, . . . . .	60
Detection of gold and platinum, . . . . .	285
Differences, table of: Group I, . . . . .	144
" " " II A, . . . . .	125
" " " II B, . . . . .	139
" " " III A, . . . . .	108
" " " III B, . . . . .	116
" " " IV, . . . . .	102
" " " V, . . . . .	96
Distillation of water, . . . . .	87, 313
" nitric acid, . . . . .	38
Distilled water, tests for, . . . . .	88, 331
Drawing out glass tube, . . . . .	44
Drying precipitates, . . . . .	62
Elements, list of chemical, . . . . .	361
Entry in note-book, rules for, . . . . .	84
" of results, examples of, . . . . .	222, 296
Equation, chemical, . . . . .	85
Evaporation, process of, . . . . .	55
Fahrenheit scale, . . . . .	361
Ferricyanides, reactions of, . . . . .	179
Ferrocyanides, " . . . . .	179
Filter-paper, . . . . .	341
Filtration, process of, . . . . .	58
Flame colorations, . . . . .	69
Flask, tubulated, . . . . .	312
Fluorides, reactions of, . . . . .	173
Fluosilicates, " " . . . . .	175
Fusion, process of, . . . . .	66
General table for groups, . . . . .	249
" " detection of rarer elements, . . . . .	354



	PAGE
Glass tube, bending, . . . . .	44
"    cutting, . . . . .	43
"    drawing out, . . . . .	44
Gold, reactions of, . . . . .	145
"    detection of, . . . . .	285
"    separation from platinum, . . . . .	285
Gramme, . . . . .	362
Group III, precipitation of, . . . . .	273-276
"    tables for analysis of, . . . . .	276-281
"    table for rarer elements, . . . . .	356
Heating glass and porcelain, . . . . .	51
Hydriodic acid, reactions of, . . . . .	163
Hydrobromic acid, " . . . . .	161
Hydrochloric acid gas, preparation of, . . . . .	36
"    "    tests for, . . . . .	36
"    acid, reactions of, . . . . .	159
Hydrocyanic acid, . . . . .	178
Hydrofluoric acid, . . . . .	175
Hydrofluosilicic acid, . . . . .	176
"    "    preparation of, . . . . .	334
Hydrogen, collection of, . . . . .	22
"    combustion of, . . . . .	24
"    explosion of, . . . . .	25
"    preparation of, . . . . .	22
"    sulphide, reactions of, . . . . .	150
"    test for, . . . . .	25
Hydrosulphuric acid, reactions of, . . . . .	150
Hypochlorites, reactions of, . . . . .	152
Hypochlorous acid, " . . . . .	152
Hyposulphites, " . . . . .	152
Hyposulphurous acid, " . . . . .	151
Ignition, process of, . . . . .	64
"    on charcoal, . . . . .	70
"    tubes, preparation of, . . . . .	46
Indigo prism, . . . . .	306
Indium, detection in analysis, . . . . .	355, 356
"    reactions of, . . . . .	351
Insoluble substances, analysis of, . . . . .	218, 286
Iodides, reactions of, . . . . .	161
Iron group, . . . . .	108
"    separation and detection of, . . . . .	109
"    table of differences, . . . . .	108
"    table for analysis, . . . . .	257
Iron, reactions of, . . . . .	105
Labelling bottles, . . . . .	324
Labels, varnishing of, . . . . .	325
Lead, reactions of, . . . . .	121
Lime-water, preparation of, . . . . .	332

	PAGE
Liquid reagents, dilution of, . . . . .	818
Lithium, detection in analysis, . . . . .	855
"    reactions of, . . . . .	852
Litmus-paper, use of, . . . . .	78
Litre, the, . . . . .	862
Magnesium, reactions of, . . . . .	94
Manganese, " . . . . .	111
Mercuricum, " . . . . .	118
Mercuriosum, " . . . . .	148
Metallic substances, analysis of, . . . . .	217, 281
Metals, examination for, . . . . .	196, 249
Molybdates, detection in analysis, . . . . .	354
"    reactions of, . . . . .	348
Mortar, agate, . . . . .	810
Neutralizing, process of, . . . . .	74
Neutral reaction, . . . . .	78
Nickel, reactions of, . . . . .	112
Nitrates, " . . . . .	154
Nitric acid, reactions of, . . . . .	155
Nitric oxide, preparation of, . . . . .	30
Nitrites, reactions of, . . . . .	158
Nitrogen tetroxide solution, preparation of, . . . . .	896
Nitrous acid, reactions of, . . . . .	158
Notation, chemical, . . . . .	84
Note-book, rules for entry in, . . . . .	84
"    examples of entry, . . . . .	222, 296
Organic acid radicles, reactions of, . . . . .	176
Oxalates, reactions of, . . . . .	180
Oxalic acid, " . . . . .	181
Oxygen gas, collection of, . . . . .	18
"    "    experiments with, . . . . .	20, 21
"    "    preparation of, . . . . .	17
"    "    test for, . . . . .	21
Palladium, detection in analysis of, . . . . .	854
"    reactions of, . . . . .	848
Phosphates, " . . . . .	166
"    table for analysis of, . . . . .	279
Phosphoric acid, reactions of, . . . . .	167
Phosphorus, combustion in oxygen, . . . . .	21
Platinum, detection of, . . . . .	286
"    reactions of, . . . . .	145
"    separation from gold, . . . . .	286
Platinum wire, mounting, . . . . .	47
"    "    cleaning, . . . . .	50
Potassium group, . . . . .	89
"    "    separation and detection, . . . . .	97

	PAGE
Potassium group, table of differences, . . . . .	96
"    "    table of analysis, . . . . .	99, 264
Potassium, reactions of, . . . . .	90
Precipitates, drying of, . . . . .	62
"    "    removal from filter, . . . . .	62
"    "    washing, . . . . .	61
Precipitation, process of, . . . . .	56
"    "    of Group III, remarks on, . . . . .	273
"    "    "    rules for, . . . . .	275
Preliminary examination for metals, . . . . .	190, 194, 240
"    "    for acid radicles, . . . . .	209, 246
"    "    of cyanogen compounds, . . . . .	292
"    "    of insoluble substances, . . . . .	218, 287
"    "    of metallic substances, . . . . .	217, 281
Prussic acid, reactions of, . . . . .	178
Rarer elements, reactions of, . . . . .	346
"    "    tables for detection of, . . . . .	354, 356
Reactions, entry of, . . . . .	84
Reagents, addition of, in excess, . . . . .	293
"    preparation of, . . . . .	318
"    list of special, . . . . .	326
"    "    general, for metals, . . . . .	328
"    "    "    acid radicles, . . . . .	330
Residues, silver, . . . . .	316
"    platinum, . . . . .	317
Rose burner, . . . . .	41
Rubidium, detection in analysis, . . . . .	355
"    reactions of, . . . . .	353
Rules for working, . . . . .	87
Saturated solutions, preparation of, . . . . .	332
Selenates, detection in analysis, . . . . .	354
"    reactions of, . . . . .	349
Selenites, "    "    . . . . .	349
Selenium, "    "    . . . . .	349
Silicates, analysis of, . . . . .	290
"    reactions of, . . . . .	172
Silicic acid, "    "    . . . . .	172
Silver group, . . . . .	143
"    "    separation and detection of, . . . . .	144
"    "    table of differences, . . . . .	144
"    "    table of analysis, . . . . .	254
"    reactions of, . . . . .	143
Simple substances, analysis of, . . . . .	185
Sodium, reactions of, . . . . .	93
Solubility, table of, . . . . .	271
Solution, process of, . . . . .	53
Solution of solid reagents, . . . . .	319
Solutions for reactions of metals, . . . . .	336
"    "    "    "    acid radicles, . . . . .	338

	PAGE
Spectroscope, its use in analysis, . . . . .	357
Spectrum chart, . . . . .	360
Spirit-lamp, . . . . .	42
Steam-oven, . . . . .	310
Stirring rods, preparation of, . . . . .	46
Strontium, reactions of, . . . . .	101
Sublimation, process of, . . . . .	65
Substances for analysis, . . . . .	341
Sulphates, reactions of, . . . . .	147
Sulphides, " " . . . . .	149
Sulphites, " " . . . . .	151
Sulphocyanides, reactions of, . . . . .	180
Sulphur, combustion in oxygen of, . . . . .	20
Sulphuretted hydrogen tube, . . . . .	45
"    "    apparatus, . . . . .	306
"    "    reactions of, . . . . .	160
"    "    passing, . . . . .	309
"    "    solution, . . . . .	333
Sulphurous acid, reactions of, . . . . .	161
"    "    preparation of, . . . . .	333
Sulphuric acid, reactions of, . . . . .	147
Symbols, chemical, . . . . .	84
"    list of, . . . . .	361
Table for analysis of Group I, . . . . .	254
"    "    II A, . . . . .	255
"    "    III A, . . . . .	257
"    "    III B, . . . . .	259
"    "    III A-B, . . . . .	276, 277, 278, 356
"    "    IV, . . . . .	262
"    "    V, . . . . .	99, 264
"    "    phosphates, . . . . .	279
Tartaric acid, reactions of, . . . . .	183
Tartrates, reactions of, . . . . .	181
Tellurates, detection in analysis, . . . . .	354
"    reactions of, . . . . .	350
Tellurites, " " . . . . .	349
Tellurium, " " . . . . .	350
Test papers, use of, . . . . .	72
Thallium, detection in analysis, . . . . .	354, 355, 356
"    reactions of, . . . . .	346
Theiosulphates, reactions of, . . . . .	151
Theiosulphuric acid, reactions of, . . . . .	152
Thermometric scales, conversion of, . . . . .	361
Tin, reactions of, . . . . .	137
Titanium, detection in analysis, . . . . .	355, 356
"    reactions of, . . . . .	351
Tungstates, detection in analysis, . . . . .	354
"    reactions of, . . . . .	347
Turmeric-paper, use of, . . . . .	75

	PAGE
Uranium, detection in analysis, . . . . .	355, 356
"    reactions of, . . . . .	350
Vanadium, detection in analysis, . . . . .	355
"    reactions of, . . . . .	352
Varnishing labels, . . . . .	325
Wash-bottle, . . . . .	47
Washing gases, . . . . .	83
"    precipitates, . . . . .	61
Water-bath, . . . . .	311
Water, distillation of, . . . . .	37, 313
"    distilling apparatus, . . . . .	313
Weights and measures, French and English, . . . . .	362, 363
Zinc group, . . . . .	109
"    "    separation and detection of, . . . . .	117
"    "    table of differences, . . . . .	116
"    "    "    for analysis, . . . . .	260
Zinc, reactions of, . . . . .	110

# CATALOGUE OF BOOKS

PUBLISHED BY

## HENRY C. LEA.

(LATE LEA & BLANCHARD.)

The books in the annexed list will be sent by mail, post-paid, to any Post Office in the United States, on receipt of the printed prices. No risks of the mail, however, are assumed, either on money or books. Gentlemen will therefore, in most cases, find it more convenient to deal with the nearest bookseller.

Detailed catalogues furnished or sent free by mail on application. An illustrated catalogue of 64 octavo pages, handsomely printed, mailed on receipt of 10 cents. Address,

HENRY C. LEA,  
Nos. 706 and 708 Sansom Street, Philadelphia.

### PERIODICALS,

*Free of Postage.*

<b>A</b> <b>AMERICAN JOURNAL OF THE MEDICAL SCIENCES.</b>	} For five Dollars per annum, in advance
Edited by Isaac Hays, M.D., published quarterly, about 1100 large 8vo. pages per annum,	
<b>MEDICAL NEWS AND LIBRARY</b> , monthly, 384 large 8vo. pages per annum,	

OR,

<b>A</b> <b>AMERICAN JOURNAL OF THE MEDICAL SCIENCES,</b>	} For six Dollars per annum, in advance.
Quarterly,	
<b>MEDICAL NEWS AND LIBRARY</b> , monthly,	
<b>MONTHLY ABSTRACT OF MEDICAL SCIENCE</b> , 48 pages per month, or nearly 600 pages per annum.	
In all, about 2100 large 8vo. pages per annum,	

**M**EDICAL NEWS AND LIBRARY, monthly, in advance, \$1 00.

**M**ONTHLY ABSTRACT OF MEDICAL SCIENCE, in advance, \$2 50.

**O**BSTETRICAL JOURNAL. With an American Supplement, edited by J. V. INGHAM, M. D. \$5 00 per annum, in advance. Single Numbers, 50 cents. Is published monthly, each number containing ninety-six octavo pages.

- ASHTON (T. J.)** ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE RECTUM AND ANUS. With remarks on Habitual Constipation. Second American from the fourth London edition, with illustrations. 1 vol. 8vo. of about 300 pp. Cloth, \$3 25.
- ASHWELL (SAMUEL).** A PRACTICAL TREATISE ON THE DISEASES OF WOMEN. Third American from the third London edition. In one 8vo. vol. of 528 pages. Cloth, \$3 50.
- ASHHURST (JOHN, Jr.)** THE PRINCIPLES AND PRACTICE OF SURGERY. FOR THE USE OF STUDENTS AND PRACTITIONERS. In 1 large 8vo. vol. of over 1000 pages, containing 533 wood-cuts. Cloth, \$6 50; leather, \$7 50.
- ATTFIELD (JOHN).** CHEMISTRY; GENERAL, MEDICAL, AND PHARMACEUTICAL. Seventh edition, revised by the author. In 1 vol. 12mo. Cloth, \$2 75; leather, \$3 25.
- BROWNE (EDGAR A.)** HOW TO USE THE OPHTHALMOSCOPE. Elementary instruction in Ophthalmoscopy for the Use of Students. In one small 12mo. vol., many illust. Cloth, \$1. (*Now ready.*)
- BLOXAM (C. L.)** CHEMISTRY, INORGANIC AND ORGANIC. With Experiments. In one handsome octavo volume of 700 pages, with 300 illustrations. Cloth, \$4 00; leather, \$5 00.
- BRINTON (WILLIAM).** LECTURES ON THE DISEASES OF THE STOMACH. From the second London ed. 1 vol. 8vo. Cloth, \$3 25.
- BIGELOW (HENRY J.)** ON DISLOCATION AND FRACTURE OF THE HIP, with the Reduction of the Dislocations by the Flexion Method. In one 8vo. vol. of 150 pp., with illustrations. Cloth, \$2 50.
- BASHAM (W. R.)** RENAL DISEASES; A CLINICAL GUIDE TO THEIR DIAGNOSIS AND TREATMENT. With illustrations. 1 vol. 12mo. Cloth, \$2 00.
- BUMSTEAD (F. J.)** THE PATHOLOGY AND TREATMENT OF VENEREAL DISEASES. Third edition, revised and enlarged, with illustrations. 1 vol. 8vo. of over 700 pages. Cloth, \$5; leather, \$6.
- **AND CULLERIER'S ATLAS OF VENEREAL.** See "CULLERIER."
- BARLOW (GEORGE H.)** A MANUAL OF THE PRACTICE OF MEDICINE. 1 vol. 8vo., of over 600 pages. Cloth, \$2 50.
- BAIRD (ROBERT).** IMPRESSIONS AND EXPERIENCES OF THE WEST INDIES. 1 vol. royal 12mo. Cloth, 75 cents.
- BARNES (ROBERT).** A PRACTICAL TREATISE ON THE DISEASES OF WOMEN. In one handsome 8vo. vol. of about 800 pages, with 169 illustrations. Cloth, \$5; leather, \$6.
- BRYANT (THOMAS).** THE PRACTICE OF SURGERY. In one handsome octavo volume, of over 1000 pages, with many illustrations. Cloth, \$6 25; leather, \$7 25.
- BRISTOWE (JOHN SYER).** A MANUAL OF THE PRACTICE OF MEDICINE. A new work, edited with additions by James H. Hutchinson, M.D. In one handsome 8vo. volume of over 1100 pages. Cloth, \$5 50; leather, \$6 50. (*Just issued*)
- BOWMAN (JOHN E.)** A PRACTICAL HAND-BOOK OF MEDICAL CHEMISTRY. Sixth American, from the fourth London edition. With numerous illustrations. 1 vol. 12mo. of 350 pp. Cloth, \$2 25.
- **INTRODUCTION TO PRACTICAL CHEMISTRY, INCLUDING ANALYSIS.** Sixth American, from the sixth London edition, with numerous illustrations. 1 vol. 12mo. of 350 pp. Cloth, \$2 25.
- BELLAMY'S MANUAL OF SURGICAL ANATOMY.** With numerous illustrations. In one royal 12mo. vol. Cloth, \$2 25. (*Lately issued.*)

- BURNETT (CHARLES H.) THE EAR: ITS ANATOMY, PHYSIOLOGY, AND DISEASES.** A Practical Treatise for the Use of Students and Practitioners. In one handsome 8vo. vol., with many illustrations. (*In press.*)
- BLANDFORD (G. FIELDING). INSANITY AND ITS TREATMENT.** With an Appendix of the laws in force in the United States on the Confinement of the Insane, by Dr. Isaac Ray. In one handsome 8vo. vol., of 471 pages. Cloth, \$3 25.
- CARTER (R. BRUDENELL). A PRACTICAL TREATISE ON DISEASES OF THE EYE.** With additions and test-types, by John Green, M.D. In one handsome 8vo. vol. of about 500 pages, with 124 illustrations. Cloth, \$3 75.
- CHAMBERS (T. K.) A MANUAL OF DIET IN HEALTH AND DISEASE.** In one handsome octavo volume of 310 pages. Cloth, \$2 75. (*Just issued.*)
- **RESTORATIVE MEDICINE.** An Harveian Annual Oration delivered at the Royal College of Physicians, London, June 21, 1871. In one small 12mo. volume. Cloth, \$1 00.
- COOPER (B. B.) LECTURES ON THE PRINCIPLES AND PRACTICE OF SURGERY.** In one large 8vo. vol. of 750 pages. Cloth, \$2 00.
- CARPENTER (WM. B.) PRINCIPLES OF HUMAN PHYSIOLOGY.** From the Eighth English Edition. In one large vol. 8vo., of 1083 pages. With 373 illustrations. Cloth, \$5 50; leather, raised bands, \$6 50. (*Just issued.*)
- **PRIZE ESSAY ON THE USE OF ALCOHOLIC LIQUORS IN HEALTH AND DISEASE.** New edition, with a Preface by D. F. Condie, M.D. 1 vol. 12mo. of 178 pages. Cloth, 60 cents.
- CLELAND (JOHN). A DIRECTORY FOR THE DISSECTION OF THE HUMAN BODY.** In one small royal 12mo. vol. Cloth, \$1 25. (*Just issued.*)
- CENTURY OF AMERICAN MEDICINE.—A HISTORY OF MEDICINE IN AMERICA, 1776-1876.** By E. H. Clarke, M.D., Late Prof. of Materia Medica in Harvard Univ.; Henry J. Bigelow, M.D., Prof. of Surgery in Harvard Univ.; Samuel D. Gross, M.D., D.C.L. Oxon., Prof. of Surgery in Jefferson Med. Coll., Philada.; T. Gaillard Thomas, Prof. of Obstetrics, etc., in Coll. of Phys. and Surgeons, N. Y.; J. S. Billings, M.D., U.S.A., Librarian of National Medical Library, Washington. In one handsome royal 12mo. volume of 366 pages. Cloth, \$2 25.
- CHRISTISON (ROBERT). DISPENSATORY OR COMMENTARY ON THE PHARMACOPŒIAS OF GREAT BRITAIN AND THE UNITED STATES.** With a Supplement by R. E. Griffith. In one 8vo. vol. of over 1000 pages, containing 213 illustrations. Cloth, \$4.
- CHURCHILL (FLEETWOOD). ON THE THEORY AND PRACTICE OF MIDWIFERY.** With notes and additions by D. Francis Condie, M.D. With about 200 illustrations. In one handsome 8vo. vol. of nearly 700 pages. Cloth, \$4; leather, \$5.
- **ESSAYS ON THE PUERPERAL FEVER, AND OTHER DISEASES PECULIAR TO WOMEN.** In one neat octavo vol. of about 450 pages. Cloth, \$2 50.
- CONDIE (D. FRANCIS). A PRACTICAL TREATISE ON THE DISEASES OF CHILDREN.** Sixth edition, revised and enlarged. In one large 8vo. vol. of 800 pages. Cloth, \$5 25; leather, \$6 25.
- CLOWES (FRANK). AN ELEMENTARY TREATISE ON PRACTICAL CHEMISTRY AND QUALITATIVE INORGANIC ANALYSIS.** Especially adapted for Laboratory Use. From the Second English Edition. In one royal 12mo. vol. (*In press.*)



- CULLERIER (A.) AN ATLAS OF VENEREAL DISEASES.** Translated and edited by FREEMAN J. BUMSTEAD, M.D. A large imperial quarto volume, with 26 plates containing about 150 figures, beautifully colored, many of them the size of life. In one vol., strongly bound in cloth, \$17.
- Same work, in five parts, paper covers, for mailing, \$3 per part.
- CYCLOPEDIA OF PRACTICAL MEDICINE.** By DUNGLISON, Forbes, Tweedie, and Conolly. In four large super-royal octavo volumes, of 3254 double-columned pages, leather, raised bands, \$15. Cloth, \$11.
- CAMPBELL'S LIVES OF LORDS KENYON, ELLENBOROUGH, AND TENTERDEN.** Being the third volume of "Campbell's Lives of the Chief Justices of England." In one crown octavo vol. Cloth, \$2.
- DALTON (J. C.) A TREATISE ON HUMAN PHYSIOLOGY.** Sixth edition, thoroughly revised, and greatly enlarged and improved, with 316 illustrations. In one very handsome 8vo. vol. of 830 pp. Cloth, \$5 50; leather, \$6 50. (*Just issued.*)
- DAVIS (F. H.) LECTURES ON CLINICAL MEDICINE.** Second edition, revised and enlarged. In one 12mo. vol. Cloth, \$1 75.
- DON QUIXOTE DE LA MANCHA.** Illustrated edition. In two handsome vols. crown 8vo. Cloth, \$2 50; half morocco, \$3 70.
- DEWEES (W. P.) A TREATISE ON THE DISEASES OF FEMALES.** With illustrations. In one 8vo. vol. of 536 pages. Cloth, \$3.
- DEBUIIT (ROBERT). THE PRINCIPLES AND PRACTICE OF MODERN SURGERY.** A revised American, from the eighth London edition. Illustrated with 432 wood engravings. In one 8vo. vol. of nearly 700 pages. Cloth, \$4; leather, \$5.
- DUNGLISON (ROBLEY). MEDICAL LEXICON; a Dictionary of Medical Science.** Containing a concise explanation of the various subjects and terms of Anatomy, Physiology, Pathology, Hygiene, Therapeutics, Pharmacology, Pharmacy, Surgery, Obstetrics, Medical Jurisprudence, and Dentistry. Notices of Climate and of Mineral Waters; Formulæ for Official, Empirical, and Dietetic Preparations, with the accentuation and Etymology of the Terms, and the French and other Synonymes. In one very large royal 8vo. vol. New edition. Cloth, \$6 50; leather, \$7 50. (*Just issued.*)
- **HUMAN PHYSIOLOGY.** Eighth edition, thoroughly revised. In two large 8vo. vols. of about 1500 pp., with 532 illus. Cloth, \$7.
- **NEW REMEDIES, WITH FORMULÆ FOR THEIR PREPARATION AND ADMINISTRATION.** Seventh edition. In one very large 8vo. vol. of 770 pages. Cloth, \$4.
- DE LA BECHE'S GEOLOGICAL OBSERVER.** In one large 8vo. vol. of 700 pages, with 300 illustrations. Cloth, \$4.
- DANA (JAMES D.) THE STRUCTURE AND CLASSIFICATION OF ZOOPHYTES.** With illust. on wood. In one imp. 4to. vol. Cloth, \$4.
- ELLIS (BENJAMIN). THE MEDICAL FORMULARY.** Being a collection of prescriptions derived from the writings and practice of the most eminent physicians of America and Europe. Twelfth edition, carefully revised by A. H. Smith, M. D. In one 8vo. volume of 374 pages. Cloth, \$3.
- ERICHSEN (JOHN). THE SCIENCE AND ART OF SURGERY.** A new and improved American, from the sixth enlarged and revised London edition. Illustrated with 630 engravings on wood. In two large 8vo. vols. Cloth, \$9 00; leather, raised bands, \$11 00.
- ENCYCLOPEDIA OF GEOGRAPHY.** In three large 8vo. vols. Illustrated with 83 maps and about 1100 wood-cuts. Cloth, \$5.

**FOTHERGILL'S PRACTITIONER'S HANDBOOK OF TREATMENT.**  
In one handsome 8vo. vol. of about 550 pp. Cloth, \$4. (*Just issued.*)

**FENWICK (SAMUEL). THE STUDENTS' GUIDE TO MEDICAL DIAGNOSIS.** From the Third Revised and Enlarged London Edition. In one vol. royal 12mo. Cloth, \$2 25.

**FLETCHER'S NOTES FROM NINEVEH, AND TRAVELS IN MESO-POTAMIA, ASSYRIA, AND SYRIA.** In one 12mo. vol. Cloth, 75 cts.

**FOX (TILBURY). EPITOME OF SKIN DISEASES, with Formulæ for Students and Practitioners.** In one small 12mo. vol. Cloth, \$1.

**FLINT (AUSTIN). A TREATISE ON THE PRINCIPLES AND PRACTICE OF MEDICINE.** Fourth edition, thoroughly revised and enlarged. In one large 8vo. volume of 1070 pages. Cloth, \$6; leather, raised bands, \$7. (*Just issued.*)

— **A PRACTICAL TREATISE ON THE PHYSICAL EXPLORATION OF THE CHEST, AND THE DIAGNOSIS OF DISEASES AFFECTING THE RESPIRATORY ORGANS.** Second and revised edition. One 8vo. vol. of 595 pages. Cloth, \$4 50.

— **A PRACTICAL TREATISE ON THE DIAGNOSIS AND TREATMENT OF DISEASES OF THE HEART.** Second edition, enlarged. In one neat 8vo. vol. of over 500 pages, \$4 00.

— **ON PHTHISIS: ITS MORBID ANATOMY, ETIOLOGY, &c.** in a series of Clinical Lectures. A new work. In one handsome 8vo. volume. Cloth, \$3 50. (*Just issued.*)

— **A MANUAL OF PERCUSSION AND AUSCULTATION; of the Physical Diagnosis of Diseases of the Lungs and Heart, and of Thoracic Aneurism.** In one handsome royal 12mo. volume. Cloth, \$1 75. (*Just issued.*)

— **MEDICAL ESSAYS.** In one neat 12mo. volume. Cloth, \$1 38.

**FOWNES (GEORGE). A MANUAL OF ELEMENTARY CHEMISTRY.** From the tenth enlarged English edition. In one royal 12mo. vol. of 857 pages, with 197 illustrations. Cloth, \$2 75; leather, \$3 25.

**FULLER (HENRY). ON DISEASES OF THE LUNGS AND AIR PASSAGES.** Their Pathology, Physical Diagnosis, Symptoms, and Treatment. From the second English edition. In one 8vo. vol. of about 500 pages. Cloth, \$3 50.

**GALLOWAY (ROBERT). A MANUAL OF QUALITATIVE ANALYSIS.** From the fifth Eng. ed. In one 12mo. vol. Cloth, \$2 50.

**GLUGE (GOTTLIEB). ATLAS OF PATHOLOGICAL HISTOLOGY.** Translated by Joseph Leidy, M.D., Professor of Anatomy in the University of Pennsylvania, &c. In one vol. imperial quarto, with 320 copperplate figures, plain and colored. Cloth, \$4.

**GREEN (T. HENRY). AN INTRODUCTION TO PATHOLOGY AND MORBID ANATOMY.** Second Amer., from the third Lond. Ed. In one handsome 8vo. vol., with numerous illustrations. Cloth, 2 75. (*Just issued.*)

**GRAY (HENRY). ANATOMY, DESCRIPTIVE AND SURGICAL.** A new American, from the fifth and enlarged London edition. In one large imperial 8vo. vol. of about 900 pages, with 462 large and elaborate engravings on wood. Cloth, \$6; leather, \$7. (*Lately issued.*)

**GRIFFITH (ROBERT E.) A UNIVERSAL FORMULARY, CONTAINING THE METHODS OF PREPARING AND ADMINISTERING OFFICIAL AND OTHER MEDICINES.** Third and Enlarged edition. Edited by John M. Maisch. In one large 8vo. vol. of 800 pages, double columns. Cloth, \$4 50; leather, \$5 50.

**GROSS (SAMUEL D.)** A SYSTEM OF SURGERY, PATHOLOGICAL, DIAGNOSTIC, THERAPEUTIC, AND OPERATIVE. Illustrated by 1403 engravings. Fifth edition, revised and improved. In two large imperial 8vo. vols. of over 2200 pages; strongly bound in leather, raised bands, \$15.

**GROSS (SAMUEL D.)** A PRACTICAL TREATISE ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE URINARY BLADDER, THE PROSTATE GLAND, AND THE URETHRA. Third Edition, thoroughly Revised and Condensed, by Samuel W. Gross, M.D., Surgeon to the Philadelphia Hospital. In one handsome octavo volume, with about two hundred illustrations. Cloth, \$4 50. (*Just issued.*)

— A PRACTICAL TREATISE ON FOREIGN BODIES IN THE AIR PASSAGES. In one 8vo. vol. of 468 pages. Cloth, \$2 75.

**GIBSON'S** INSTITUTES AND PRACTICE OF SURGERY. In two 8vo. vols. of about 1000 pages, leather, \$6 50.

**GOSSELIN (L.)** CLINICAL LECTURES ON SURGERY, Delivered at the Hospital of La Charité Translated from the French by Lewis A. Stimson, M.D., Surgeon to the Presbyterian Hospital, New York. With illustrations. (*Publishing in the Medical News and Library for 1876-7.*)

**HUDSON (A.)** LECTURES ON THE STUDY OF FEVER. 1 vol. 8vo., 316 pages. Cloth, \$2 50.

**HEATH (CHRISTOPHER).** PRACTICAL ANATOMY; A MANUAL OF DISSECTIONS. With additions, by W. W. Keen, M.D. In 1 volume; with 247 illustrations. Cloth, \$3 50; leather, \$4.

**HARTSHORNE (HENRY).** ESSENTIALS OF THE PRINCIPLES AND PRACTICE OF MEDICINE. Fourth and revised edition. In one 12mo. vol. Cloth, \$2 63; half bound, \$2 88. (*Lately issued.*)

— CONSPICUOUS OF THE MEDICAL SCIENCES. Comprising Manuals of Anatomy, Physiology, Chemistry, Materia Medica, Practice of Medicine, Surgery, and Obstetrics. Second Edition. In one royal 12mo. volume of over 1000 pages, with 477 illustrations. Strongly bound in leather, \$5 00; cloth, \$4 25. (*Lately issued.*)

— A HANDBOOK OF ANATOMY AND PHYSIOLOGY. In one neat royal 12mo. volume, with many illustrations. Cloth, \$1 75.

**HAMILTON (FRANK H.)** A PRACTICAL TREATISE ON FRACTURES AND DISLOCATIONS. Fifth edition, carefully revised. In one handsome 8vo. vol. of 830 pages, with 344 illustrations. Cloth, \$5 75; leather, \$6 75. (*Just issued.*)

**HOLMES (TIMOTHY).** SURGERY, ITS PRINCIPLES AND PRACTICE. In one handsome 8vo. volume of 1000 pages, with 411 illustrations. Cloth, \$6; leather, with raised bands, \$7. (*Just ready.*)

**HOBLYN (RICHARD D.)** A DICTIONARY OF THE TERMS USED IN MEDICINE AND THE COLLATERAL SCIENCES. In one 12mo. volume, of over 500 double-columned pages. Cloth, \$1 50; leather, \$2.

**HJUDGE (HUGH L.)** ON DISEASES PECULIAR TO WOMEN, INCLUDING DISPLACEMENTS OF THE UTERUS. Second and revised edition. In one 8vo. volume. Cloth, \$4 50.

— THE PRINCIPLES AND PRACTICE OF OBSTETRICS. Illustrated with large lithographic plates containing 159 figures from original photographs, and with numerous wood-cuts. In one large quarto vol. of 550 double-columned pages. Strongly bound in cloth, \$14.

**HOLLAND (SIR HENRY).** MEDICAL NOTES AND REFLECTIONS. From the third English edition. In one 8vo. vol. of about 500 pages. Cloth, \$3 50.

**HODGES (RICHARD M.)** PRACTICAL DISSECTIONS. Second edition. In one neat royal 12mo. vol., half bound, \$2.

**HUGHES.** SCRIPTURE GEOGRAPHY AND HISTORY, with 12 colored maps. In 1 vol. 12mo. Cloth, \$1.

**HORNER (WILLIAM E.)** SPECIAL ANATOMY AND HISTOLOGY. Eighth edition, revised and modified. In two large 8vo. vols. of over 1000 pages, containing 300 wood-cuts. Cloth, \$6.

**HILL (BERKELEY).** SYPHILIS AND LOCAL CONTAGIOUS DISORDERS. In one 8vo. volume of 467 pages. Cloth, \$3 25.

**HILLIER (THOMAS).** HAND-BOOK OF SKIN DISEASES. Second Edition. In one neat royal 12mo. volume of about 300 pp., with two plates. Cloth, \$2 25

**HALL (MRS. M.)** LIVES OF THE QUEENS OF ENGLAND BEFORE THE NORMAN CONQUEST. In one handsome 8vo. vol. Cloth, \$2 25; crimson cloth, \$2 50; half morocco, \$3.

**JONES (C. HANDFIELD).** CLINICAL OBSERVATIONS ON FUNCTIONAL NERVOUS DISORDERS. Second American Edition. In one 8vo. vol. of 348 pages. Cloth, \$3 25.

**KIRKES (WILLIAM SENHOUSE).** A MANUAL OF PHYSIOLOGY. A new American, from the eighth London edition. One vol., with many illus., 12mo. Cloth, \$3 25; leather, \$3 75.

**KNAPP (F.)** TECHNOLOGY; OR CHEMISTRY, APPLIED TO THE ARTS AND TO MANUFACTURES, with American additions, by Prof. Walter R. Johnson. In two 8vo. vols., with 500 ill. Cloth, \$6.

**KENNEDY'S MEMOIRS OF THE LIFE OF WILLIAM WIRT.** In two vols. 12mo. Cloth, \$2.

**LEA (HENRY C.)** SUPERSTITION AND FORCE; ESSAYS ON THE WAGER OF LAW, THE WAGER OF BATTLE, THE ORDEAL, AND TORTURE. Second edition, revised. In one handsome royal 12mo. vol., \$2 75.

— STUDIES IN CHURCH HISTORY. The Rise of the Temporal Power—Benefit of Clergy—Excommunication. In one handsome 12mo. vol. of 515 pp. Cloth, \$2 75.

— AN HISTORICAL SKETCH OF SACERDOTAL CELIBACY IN THE CHRISTIAN CHURCH. In one handsome octavo volume of 602 pages. Cloth, \$3 75.

**LA ROCHE (R.)** YELLOW FEVER. In two 8vo. vols. of nearly 1500 pages. Cloth, \$7.

— PNEUMONIA. In one 8vo. vol. of 500 pages. Cloth, \$3.

**LINCOLN (D. F.)** ELECTRO-THERAPEUTICS. A Condensed Manual of Medical Electricity. In one neat royal 12mo. volume, with illustrations. Cloth, \$1 50. (*Just issued.*)

**LISHMAN (WILLIAM).** A SYSTEM OF MIDWIFERY. Including the Diseases of Pregnancy and the Puerperal State. Second American, from the Second English Edition. With additions, by J. S. Parry, M.D. In one very handsome 8vo. vol. of 800 pages and 200 illustrations. Cloth, \$5; leather, \$6. (*Just issued.*)

**LAURENCE (J. Z.) AND MOON (ROBERT C.)** A HANDY-BOOK OF OPHTHALMIC SURGERY. Second edition, revised by Mr. Laurence. With numerous illus. In one 8vo. vol. Cloth, \$2 75.

- LEHMANN (C. G.)** PHYSIOLOGICAL CHEMISTRY. Translated by George F. Day, M.D. With plates, and nearly 200 illustrations. In two large 8vo. vols., containing 1200 pages. Cloth, \$6.
- A MANUAL OF CHEMICAL PHYSIOLOGY. In one very handsome 8vo. vol. of 336 pages. Cloth, \$2 25.
- LAWSON (GEORGE).** INJURIES OF THE EYE, ORBIT, AND EYE-LIDS, with about 100 illustrations. From the last English edition. In one handsome 8vo. vol. Cloth, \$3 50.
- LUDLOW (J. L.)** A MANUAL OF EXAMINATIONS UPON ANATOMY, PHYSIOLOGY, SURGERY, PRACTICE OF MEDICINE, OBSTETRICS, MATERIA MEDICA, CHEMISTRY, PHARMACY, AND THERAPEUTICS. To which is added a Medical Formulary. Third edition. In one royal 12mo. vol. of over 800 pages. Cloth, \$3 25; leather, \$3 75.
- LYNCH (W. F.)** A NARRATIVE OF THE UNITED STATES EXPEDITION TO THE DEAD SEA AND RIVER JORDAN. In one large octavo vol., with 28 beautiful plates and two maps. Cloth, \$3.
- Same Work, condensed edition. One vol. royal 12mo. Cloth, \$1.
- LEE (HENRY)** ON SYPHILIS. In one 8vo. vol. Cloth, \$2 25.
- LYONS (ROBERT D.)** A TREATISE ON FEVER. In one neat 8vo. vol. of 362 pages. Cloth, \$2 25.
- MARSHALL (JOHN).** OUTLINES OF PHYSIOLOGY, HUMAN AND COMPARATIVE. With Additions by FRANCIS G. SMITH, M. D., Professor of the Institutes of Medicine in the University of Pennsylvania. In one 8vo. volume of 1026 pages, with 122 illustrations. Strongly bound in leather, raised bands, \$7 50. Cloth, \$6 50.
- MACLISE (JOSEPH).** SURGICAL ANATOMY. In one large imperial quarto vol., with 68 splendid plates, beautifully colored; containing 190 figures, many of them life size. Cloth, \$14.
- MEIGS (CHAS. D.).** ON THE NATURE, SIGNS, AND TREATMENT OF CHILDBED FEVER. In one 8vo. vol. of 365 pages. Cloth, \$2.
- MILLER (JAMES).** PRINCIPLES OF SURGERY. Fourth American, from the third Edinburgh edition. In one large 8vo. vol. of 700 pages, with 240 illustrations. Cloth, \$3 75.
- THE PRACTICE OF SURGERY. Fourth American, from the last Edinburgh edition. In one large 8vo. vol. of 700 pages, with 364 illustrations. Cloth, \$3 75.
- MONTGOMERY (W. F.)** AN EXPOSITION OF THE SIGNS AND SYMPTOMS OF PREGNANCY. From the second English edition. In one handsome 8vo. vol. of nearly 600 pages. Cloth, \$3 75.
- MULLER (J.)** PRINCIPLES OF PHYSICS AND METEOROLOGY. In one large 8vo. vol. with 550 wood-cuts, and two colored plates. Cloth, \$4 50.
- MIRABEAU; A LIFE HISTORY.** In one 12mo. vol. Cloth, 75 cts.
- MACFARLAND'S** TURKEY AND ITS DESTINY. In 2 vols. royal 12mo. Cloth, \$2.
- MARSH (MRS.)** A HISTORY OF THE PROTESTANT REFORMATION IN FRANCE. In 2 vols. royal 12mo. Cloth, \$2.
- NEELIGAN (J. MOORE).** AN ATLAS OF CUTANEOUS DISEASES. In one quarto volume, with beautifully colored plates, &c. Cloth, \$5 50.
- NEILL (JOHN) AND SMITH (FRANCIS G.)** COMPENDIUM OF THE VARIOUS BRANCHES OF MEDICAL SCIENCE. In one handsome 12mo. vol. of about 1000 pages, with 374 wood-cuts. Cloth, \$4; leather, raised bands, \$4 75.

- NIEBUHR (B. G.)** LECTURES ON ANCIENT HISTORY; comprising the history of the Asiatic Nations, the Egyptians, Greeks, Macedonians, and Carthaginians. Translated by Dr. L. Schmitz. In three neat volumes, crown octavo. Cloth, \$5 00.
- ODLING (WILLIAM).** A COURSE OF PRACTICAL CHEMISTRY FOR THE USE OF MEDICAL STUDENTS. In one 12mo. vol. of 261 pp., with 75 illustrations. Cloth, \$2.
- PLAYFAIR (W. S.)** A TREATISE ON THE SCIENCE AND PRACTICE OF MIDWIFERY. In one handsome octavo vol. of 576 pp., with 166 illustrations, and two plates. Cloth, \$4; leather, \$5. (*Just issued.*)
- PAY (F. W.)** A TREATISE ON THE FUNCTION OF DIGESTION, ITS DISORDERS AND THEIR TREATMENT. From the second London ed. In one 8vo. vol. of 246 pp. Cloth, \$2.
- A TREATISE ON FOOD AND DIETETICS, PHYSIOLOGICALLY AND THERAPEUTICALLY CONSIDERED. In one neat octavo volume of about 500 pages. Cloth, \$4 75. (*Just issued.*)
- PARRISH (EDWARD).** A TREATISE ON PHARMACY. With many Formulæ and Prescriptions. Fourth edition. Enlarged and thoroughly revised by Thomas S. Wiegand. In one handsome 8vo. vol. of 977 pages, with 280 illus. Cloth, \$5 50; leather, \$6 50.
- PIEBIE (WILLIAM)** THE PRINCIPLES AND PRACTICE OF SURGERY. In one handsome octavo volume of 780 pages, with 316 illustrations. Cloth, \$3 75.
- PEREIRA (JONATHAN).** MATERIA MEDICA AND THERAPEUTICS. An abridged edition. With numerous additions and references to the United States Pharmacopœia. By Horatio C. Wood, M. D. In one large octavo volume, of 1040 pages, with 236 illustrations. Cloth, \$7 00; leather, raised bands, \$8 00.
- PULSZKY'S** MEMOIRS OF AN HUNGARIAN LADY. In one neat royal 12mo. vol. Cloth, \$1.
- PAGET'S** HUNGARY AND TRANSYLVANIA. In two royal 12mo. vols. Cloth, \$2.
- REMSEN (IRA).** THE PRINCIPLES OF CHEMISTRY. In one handsome 12mo. vol. Cloth, \$1 50. (*Just issued.*)
- ROBERTS (WILLIAM).** A PRACTICAL TREATISE ON URINARY AND RENAL DISEASES. A second American, from the second London edition. With numerous illustrations and a colored plate. In one very handsome 8vo. vol. of 616 pages. Cloth, \$4 50.
- RAMSBOTHAM (FRANCIS H.)** THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDICINE AND SURGERY. In one imperial 8vo. vol. of 650 pages, with 64 plates, besides numerous woodcuts in the text. Strongly bound in leather, \$7.
- RIGBY (EDWARD).** A SYSTEM OF MIDWIFERY. Second American edition. In one handsome 8vo. vol. of 422 pages. Cloth, \$2 50.
- RANKE'S** HISTORY OF THE TURKISH AND SPANISH EMPIRES in the 16th and beginning of 17th Century. In one 8vo. volume, paper, 25 cts.
- HISTORY OF THE REFORMATION IN GERMANY. Parts I., II., III. In one vol. Cloth, \$1.
- SCHAFER (EDWARD ALBERT)** A COURSE OF PRACTICAL HISTOLOGY: A Manual of the Microscope for Medical Students. In one handsome octavo vol. With many illust. Cloth, \$2. (*Just issued.*)

- SMITH (EUSTACE).** ON THE WASTING DISEASES OF CHILDREN. Second American edition, enlarged. In one 8vo. vol. Cloth, \$2 50.
- SARGENT (F. W.)** ON BANDAGING AND OTHER OPERATIONS OF MINOR SURGERY. New edition, with an additional chapter on Military Surgery. In one handsome royal 12mo. vol. of nearly 400 pages, with 184 wood-cuts. Cloth, \$1 75.
- SMITH (J. LEWIS.)** A TREATISE ON THE DISEASES OF INFANCY AND CHILDHOOD. Third Edition, revised and enlarged. In one large 8vo. volume of 724 pages, with illustrations. Cloth, \$5; leather, \$6. (*Just issued.*)
- SHARPEY (WILLIAM) AND QUAIN (JONES AND RICHARD).** HUMAN ANATOMY. With notes and additions by Jos. Leidy, M.D., Prof. of Anatomy in the University of Pennsylvania. In two large 8vo. vols. of about 1300 pages, with 511 illustrations. Cloth, \$6.
- SKEY (FREDERIC C.)** OPERATIVE SURGERY. In one 8vo. vol. of over 650 pages, with about 100 wood-cuts. Cloth, \$3 25.
- SLADE (D. D.)** DIPHTHERIA; ITS NATURE AND TREATMENT. Second edition. In one neat royal 12mo. vol. Cloth, \$1 25.
- SMITH (HENRY H.) AND HORNER (WILLIAM E.)** ANATOMICAL ATLAS. Illustrative of the structure of the Human Body. In one large imperial 8vo. vol., with about 650 beautiful figures. Cloth, \$4 50.
- SMITH (EDWARD).** CONSUMPTION; ITS EARLY AND REMEDIABLE STAGES. In one 8vo. vol. of 254 pp. Cloth, \$2 25.
- STILLE (ALFRED).** THERAPEUTICS AND MATERIA MEDICA. Fourth edition, revised and enlarged. In two large and handsome volumes 8vo. Cloth, \$10; leather, \$12. (*Just issued.*)
- STILLE (ALFRED) AND MAISCH (JOHN M.)** THE NATIONAL DISPENSATORY: Embracing the Chemistry, Botany, Materia Medica, Pharmacy, Pharmacodynamics, and Therapeutics of the Pharmacopœias of the United States and Great Britain. For the Use of Physicians and Pharmacutists. In one handsome 8vo. vol., with numerous illustrations. (*in press.*)
- SCHMITZ AND ZUMPT'S CLASSICAL SERIES.** In royal 18mo.
- CORNELII NEPOTIS LIBER DE EXCELLENTIBUS DUCIBUS EXTERARUM GENTIUM, CUM VITIS CATONIS ET ATTICI.** With notes, &c. Price in cloth, 60 cents; half bound, 70 cts.
- C. I. CÆSARIS COMMENTARI DE BELLO GALLICO.** With notes, map, and other illustrations. Cloth, 60 cents; half bound, 70 cents.
- C. C. SALLUSTII DE BELLO CATILINARIO ET JUGURTHINO.** With notes, map, &c. Price in cloth, 60 cents; half bound, 70 cents.
- Q. CURTII RUFII DE GESTIS ALEXANDRI MAGNI LIBRI VIII.** With notes, map, &c. Price in cloth, 80 cents; half bound, 90 cents.
- P. VIRGILII MARONIS CARMINA OMNIA.** Price in cloth, 85 cents; half bound, \$1.
- M. T. CICERONIS ORATIONES SELECTÆ XII.** With notes, &c. Price in cloth, 70 cents; half bound, 80 cents.
- ECLOGÆ EX Q. HORATII FLACCI POEMATIBUS.** With notes, &c. Price in cloth, 70 cents; half bound, 80 cents.
- ADVANCED LATIN EXERCISES, WITH SELECTIONS FOR READING.** Revised. Cloth, price 60 cents; half bound, 70 cents.

- SWAYNE (JOSEPH GRIFFITHS).** OBSTETRIC APHORISMS. A new American, from the fifth revised English edition. With additions by E. R. Hutchins, M. D. In one small 12mo. vol. of 177 pp., with illustrations. Cloth, \$1 25.
- STURGES (OCTAVIUS).** AN INTRODUCTION TO THE STUDY OF CLINICAL MEDICINE. In one 12mo. vol. Cloth, \$1 25.
- SCHOEDLER (FREDERICK) AND MEDLOCK (HENRY).** WONDERS OF NATURE. An elementary introduction to the Sciences of Physics, Astronomy, Chemistry, Mineralogy, Geology, Botany, Zoology, and Physiology. Translated from the German by H. Medlock. In one neat 8vo. vol., with 679 illustrations. Cloth, \$3.
- STOKES (W.)** LECTURES ON FEVER. In one 8vo. vol. Cloth, \$2.
- SMALL BOOKS ON GREAT SUBJECTS.** Twelve works; each one 10 cents, sewed, forming a neat and cheap series; or done up in 3 vols., cloth, \$1 50.
- STRICKLAND (AGNES).** LIVES OF THE QUEENS OF HENRY THE VIII. AND OF HIS MOTHER. In one crown octavo vol., extra cloth, \$1; black cloth, 90 cents.
- MEMOIRS OF ELIZABETH, SECOND QUEEN REGNANT OF ENGLAND AND IRELAND. In one crown octavo vol., extra cloth, \$1 40; black cloth, \$1 30.
- TANNER (THOMAS HAWKES).** A MANUAL OF CLINICAL MEDICINE AND PHYSICAL DIAGNOSIS. Third American from the second revised English edition. Edited by Tilbury Fox, M. D. In one handsome 12mo. volume of 366 pp. Cloth, \$1 50.
- ON THE SIGNS AND DISEASES OF PREGNANCY. From the second English edition. With four colored plates and numerous illustrations on wood. In one vol. 8vo. of about 500 pages. Cloth, \$4 25.
- TUKE (DANIEL HACK).** INFLUENCE OF THE MIND UPON THE BODY. In one handsome 8vo. vol. of 416 pp. Cloth, \$3 25.
- TAYLOR (ALFRED S.)** MEDICAL JURISPRUDENCE. Seventh American edition. Edited by John J. Reese, M. D. In one large 8vo. volume of 879 pages. Cloth, \$5; leather, \$6. (*Just issued.*)
- PRINCIPLES AND PRACTICE OF MEDICAL JURISPRUDENCE. From the Second English Edition. In two large 8vo. vols. Cloth, \$10; leather, \$12. (*Just issued.*)
- ON POISONS IN RELATION TO MEDICINE AND MEDICAL JURISPRUDENCE. Third American from the Third London Edition. 1 vol. 8vo. of 788 pages, with 104 illustrations. Cloth, \$5 50; leather, \$6 50. (*Just issued.*)
- THOMAS (T. GAILLARD).** A PRACTICAL TREATISE ON THE DISEASES OF FEMALES. Fourth edition, thoroughly revised. In one large and handsome octavo volume of 801 pages, with 191 illustrations. Cloth, \$5 00; leather, \$6 00. (*Just issued.*)
- TODD (ROBERT BENTLEY).** CLINICAL LECTURES ON CERTAIN ACUTE DISEASES. In one vol. 8vo. of 320 pp., cloth, \$2 50.
- THOMPSON (SIR HENRY).** CLINICAL LECTURES ON DISEASES OF THE URINARY ORGANS. Second and revised edition. In one 8vo. volume, with illustrations. Cloth, \$2 25. (*Just issued.*)
- THE DISEASES OF THE PROSTATE, THEIR PATHOLOGY AND TREATMENT. Fourth edition, revised. In one very handsome 8vo. vol. of 355 pp., with 13 plates. Cloth, \$3 75.



**T**HOMPSON (SIR HENRY). THE PATHOLOGY AND TREATMENT OF STRICTURE OF THE URETHRA AND URINARY FISTULÆ. From the third English edition. In one 8vo. vol. of 359 pp., with illustrations. Cloth, \$3 50.

**W**ALSHE (W. H.) PRACTICAL TREATISE ON THE DISEASES OF THE HEART AND GREAT VESSELS. Third American from the third revised London edition. In one 8vo. vol. of 420 pages. Cloth, \$3.

**W**ATSON (THOMAS). LECTURES ON THE PRINCIPLES AND PRACTICE OF PHYSIC. A new American from the fifth and enlarged English edition, with additions by H. Hartshorne, M.D. In two large and handsome octavo volumes. Cloth, \$9; leather, \$11.

**W**ÖHLER'S OUTLINES OF ORGANIC CHEMISTRY. Translated from the 8th German edition, by Ira Remsen, M.D. In one neat 12mo. vol. Cloth, \$3 00. (*Lately issued.*)

**W**ELLS (J. SOELBERG). A TREATISE ON THE DISEASES OF THE EYE. Second American, from the Third English edition, with additions by I. Minis Hays, M.D. In one large and handsome octavo vol., with 6 colored plates and many wood-cuts, also selections from the test-types of Jaeger and Snellen. Cloth, \$5 00; leather, \$6 00.

**W**HAT TO OBSERVE AT THE BEDSIDE AND AFTER DEATH IN MEDICAL CASES. In one royal 12mo. vol. Cloth, \$1.

**W**EST (CHARLES). LECTURES ON THE DISEASES PECULIAR TO WOMEN. Third American from the Third English edition. In one octavo volume of 550 pages. Cloth, \$3 75; leather, \$4 75.

— LECTURES ON THE DISEASES OF INFANCY AND CHILDHOOD. Fifth American from the sixth revised English edition. In one large 8vo. vol. of 670 closely printed pages. Cloth, \$4 50; leather, \$5 50. (*Just issued.*)

— ON SOME DISORDERS OF THE NERVOUS SYSTEM IN CHILDHOOD. From the London Edition. In one small 12mo. volume. Cloth, \$1.

**W**ILLIAMS (CHARLES J. B. and C. T.) PULMONARY CONSUMPTION: ITS NATURE, VARIETIES, AND TREATMENT. In one neat octavo volume. Cloth, \$2 50.

**W**ILSON (ERASMUS). A SYSTEM OF HUMAN ANATOMY. A new and revised American from the last English edition. Illustrated with 397 engravings on wood. In one handsome 8vo. vol. of over 600 pages. Cloth, \$4; leather, \$5.

— ON DISEASES OF THE SKIN. The seventh American from the last English edition. In one large 8vo. vol. of over 800 pages. Cloth, \$5.


Also, A SERIES OF PLATES, illustrating "Wilson on Diseases of the Skin," consisting of 20 plates, thirteen of which are beautifully colored, representing about one hundred varieties of Disease. \$5 50.

Also, the TEXT AND PLATES, bound in one volume. Cloth, \$10.


— THE STUDENT'S BOOK OF CUTANEOUS MEDICINE. In one handsome royal 12mo. vol. Cloth, \$3 50.

**W**INCKEL ON PATHOLOGY AND TREATMENT OF CHILDBED. With Additions by the Author. Translated by Chadwick. In one handsome octavo volume of 484 pages. Cloth, \$4. (*Just issued.*)


**Z**EISSL ON VENEREAL DISEASES. Translated by Sturgis. (*Preparing.*)











the 1990s, the number of people in the world who are undernourished has increased from 600 million to 800 million. The number of people who are malnourished has increased from 1.2 billion to 1.6 billion. The number of people who are obese has increased from 100 million to 300 million.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

The World Bank has estimated that the cost of malnutrition to the world economy is \$100 billion per year. The cost of obesity to the world economy is \$100 billion per year. The cost of undernutrition to the world economy is \$100 billion per year.

DEC 10 1880

DEC 4 1889

JUN 1 1898

JAN 11 1882

MAY 6 1890

MAR 11 1882

OCT 8 1890

MAR 9 1900

DEC 13 1882

NOV 24 1890

OCT 30 1900

SEP 19 1884

APR 1 1892

DEC 14 1901

APR 8 1885

JUN 23 1885

JUL 22 1885

SEP 24 1892

AUG 22 1885

MAR 2 1893

SEP 24 1885

DEC 22 1902

OCT 26 1885

DEC 9 1893

DEC 4 1885

JAN 16 1886

DEC 23 1893

JUN 24 1886

SEP 2 1886

FEB 17 1894

OCT 18 1903

JAN 21 1888

MAR 6 1894

JAN 27 1903

APR 13 1888

OCT 18 1888

APR 25 1894

DEC 8 1888

FEB 12 1889

MAY 17 1889

NOV 28 1896

JUL 13 1889

JAN 26 1897



Chem 668.77.3  
An elementary treatise on practical  
Cabot Science 001357208



3 2044 091 914 713